

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
18 July 2002 (18.07.2002)

PCT

(10) International Publication Number
WO 02/055665 A2

(51) International Patent Classification⁷: C12N
(21) International Application Number: PCT/US02/00311
(22) International Filing Date: 7 January 2002 (07.01.2002)
(25) Filing Language: English
(26) Publication Language: English
(30) Priority Data:
60/260,179 5 January 2001 (05.01.2001) US
(71) Applicant: NEW YORK UNIVERSITY [US/US]; 70
Washington Square South, New York, NY 10012 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LI, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(72) Inventor: PAGANO, Michele; One Washington Square Village, New York, NY 10012 (US).

(74) Agents: REIN, Barry, D. et al.; Pennie & Edmonds LLP, 1155 Avenue of the Americas, New York, NY 10036 (US).

(54) Title: METHODS TO IDENTIFY COMPOUNDS USEFUL FOR THE TREATMENT OF PROLIFERATIVE AND DIFFERENTIATIVE DISORDERS

5KP2** 10 20 30 40 50
Fbp1** ... D E L L G ... S C L P E ... L K V S G ... Y R L A S ... D E S I S W O
Fbp2** ... A R G L C H I A E N ... D A K S ... C A A E L ... Y R V Y S ... D G M L W K
Fbp3** ... L E L S F V ... G D P O T ... L T C C L ... S K W N K V T S ... A C T E L W O
Fbp4** ... O D I N L O V F K Y ... G P L L D ... R A H A S O V E R N N O V F H ... M P D L W A
Fbp5** ... H H V I V C O M F O Y ... S P H O ... R A C A S V I C H R W N E V F H ... I S D L W A
Fbp6** ... I V I V C O M F O Y ... S P H O ... G C L G S T N H Y W N E V R ... N P I L W A
Fbp7** ... O N I L L R F T A O ... S P H O ... I N V S K V S I T T W K K L E D ... D K G A F O
Fbp8** ... L E S K R U F R ... L L D V R S ... L N C H L V I C S L R D D I D ... L L T L W K
Fbp9** ... P E S S T T A S Y ... N A T A L ... L C S A ... V I C H O L F T A S N ... D P L W A
Fbp10** ... Q E V L E Y L O C Q S ... L T A A L ... L O T S T T O R L R E C S Q S ... D E L L W A
Fbp11** ... E V V E R Y L S Y ... U P A K A ... G R A V C V C R L R E C S Q S ... S G K V W K
Fbp12** ... D E V L K F S Y ... U L E Q D ... G R A C V C K R F S E V R ... R V L R T H S V W K
Fbp13** ... L E L W H M C A Y ... G H P O D ... G R C S L V C R A Y E S L S ... D P N L W K
Fbp14** ... T D P L L L S F ... G O V R D ... I N C O Y V S A R L S O S S ... H O P L W A
Fbp15** ... W A W G E K G V L S N U E A L ... D G G ... D O P V W L V I C G S W R R H V G ... A G L C W A
Fbp16** ... P E L L E R V L A A ... U P A A E L ... V A C R L V C L R K E V D ... G A P L W A
Fbp17** ... E V L E H M C S Y ... G M R A ... N V A L G O T C A Y F H E V D ... G E Q W A
Fbp18** ... L H M G N W E Y R F ... S O G W O ... G O L A G V Y N W L W H T N C ... D L L R R O I A W A
Fbp19** ... D H S M V O F S T ... U P T N O ... I T L G O V Y P T L Y M E S E ... D R G A W K
Fbp20** ... L E L L G O F G L ... U P A A G M P F ... G R I C A R V I C H R W N E V R ... D P R A W R
Fbp21** ... P E L L E R V L A A ... U P O E ... G R A A R V I C H R W N E V R ... O P A W W K
Fbp22** ... K E L L R F S F ... D O I V T ... G R I C S O V S M K W S O L Y K ... T O S L W K
Fbp23** ... Y E L L G L A R N ... L L P M ... G R I C A O E S K A L N T I A L ... D S N M O
Fbp24** ... M E L L G L A R N ... L L P M ... G R I C A O E S K A L N T I A L ... D S N M O
Fbp25** ... P E L L E R V L A A ... U P O E ... G R I C A O E S K A L N T I A L ... D S N M O
Fbp25** ... P E L L E R V L A A ... U P O E ... G R I C A O E S K A L N T I A L ... D S N M O

WO 02/055665 A2

(57) Abstract: The present invention relates to the discovery, identification and characterization of nucleotides that encode novel substrate-targeting subunits of ubiquitin ligases. The invention encompasses nucleotides encoding novel substrate-targeting subunits of ubiquitin ligases: FBP1, FBP2, FBP3, FBP4, FBP5, FBP6, FBP7, FBP8, FBP9, FBP10, FBP11, FBP12, FBP13, FBP14, FBP15, FBP16, FBP17, FBP18, FBP19, FBP20, FBP21, FBP22, FBP23, FBP24, and FBP25, transgenic mice, knock-out mice, host cell expression systems and proteins encoded by the nucleotides of the present invention. The present invention relates to screening assays that use the novel substrate-targeting subunits to identify potential therapeutic agents such as small molecules, compounds or derivatives and analogues of the novel ubiquitin ligases which modulates activity of the novel ubiquitin ligases for the treatment of proliferative and differentiative disorders, such as cancer, major opportunistic infections, immune disorders, certain cardiovascular diseases, and inflammatory disorders. The invention further encompasses therapeutic protocols and pharmaceutical compositions designed to target ubiquitin ligases and their substrates for the treatment of proliferative disorders.

**METHODS TO IDENTIFY COMPOUNDS USEFUL FOR
THE TREATMENT OF PROLIFERATIVE AND
DIFFERENTIATIVE DISORDERS**

- 5 This application claims priority under 35 U.S.C. §119(e) to U.S. Application
• No. 60/260,179, filed January 5, 2001, the contents of which are incorporated herein by
reference in their entirety.

1. INTRODUCTION

10 The present invention relates to the discovery, identification and
characterization of nucleotide sequences that encode novel substrate-targeting subunits of
ubiquitin ligases. The invention encompasses nucleic acid molecules comprising nucleotide
sequences encoding novel substrate-targeting subunits of ubiquitin ligases: FBP1, FBP2,
FBP3a, FBP3b, FBP4, FBP5, FBP6, FBP7, FBP8, FBP11, FBP12, FBP13, FBP14, FBP15,
15 FBP17, FBP18, FBP20, FBP21, FBP22, FBP23, AND FBP25, transgenic mice, knock-out
mice, host cell expression systems and proteins encoded by the nucleotides of the present
invention. The present invention relates to screening assays to identify potential therapeutic
agents such as small molecules, compounds or derivatives and analogues of the novel
ubiquitin ligases which modulate activity of the novel ubiquitin ligases for the treatment of
20 proliferative and differentiative disorders, such as cancer, major opportunistic infections,
immune disorders, certain cardiovascular diseases, and inflammatory disorders. The
invention further encompasses therapeutic protocols and pharmaceutical compositions
designed to target ubiquitin ligases and their substrates for the treatment of proliferative
disorders.

25

2. BACKGROUND OF THE INVENTION

2.1 CELL CYCLE REGULATORY PROTEINS

30 The eukaryotic cell cycle is regulated by a family of serine/threonine protein
kinases called cyclin dependent kinases (Cdks) because their activity requires the
association with regulatory subunits named Cyclins (Hunter & Pines, 1994, Cell 79:573).
Cdks also associate with Cdk inhibitors (Ckis) which mediate cell cycle arrest in response to
various antiproliferative signals. So far, based on their sequence homology, two families of
Ckis have been identified in mammalian cells: the Cip/Kip family, which includes p21, p27
35 and p57; and the Ink family, which includes p15, p16, p18, and p20 (Sherr & Roberts, 1999,
Genes & Dev. 13: 1501).

2.2 THE UBIQUITIN PATHWAY

Ubiquitin-mediated proteolysis is an important pathway of non-lysosomal protein degradation which controls the timed destruction of many cellular regulatory proteins including, p27, p53, p300, cyclins, E2F, STAT-1, c-Myc, c-Jun, EGF receptor, 5 IkB α , NFkB and β -catenin (reviewed in Pagano, 1997, FASEB J. 11:1067). Ubiquitin is an evolutionary highly conserved 76-amino acid polypeptide which is abundantly present in all eukaryotic cells. The ubiquitin pathway leads to the covalent attachment of a poly-ubiquitin chain to target substrates which are then degraded by the multi-catalytic proteasome 10 complex (see Pagano, supra, for a recent review). Many of the steps regulating protein ubiquitination are known. Initially the ubiquitin activating enzyme (E1), forms a high energy thioester with ubiquitin which is, in turn, transferred to a reactive cysteine residue of one of many ubiquitin conjugating enzymes (Ubc's or E2's). The final transfer of ubiquitin to an ϵ -amino group of a reactive lysine residue in the target protein occurs in a reaction that 15 may or may not require an ubiquitin ligase (E3) protein. The large number of ubiquitin ligases ensures the high level of substrate specificity.

2.3 THE UBIQUITIN PATHWAY AND THE REGULATION OF THE G1 PHASE BY F BOX PROTEINS

20 Genetic and biochemical studies in several organisms have shown that the G1 phase of the cell cycle is regulated by the ubiquitin pathway. Proteolysis of cyclins, Ck's and other G1 regulatory proteins is controlled in yeast by the ubiquitin conjugating enzyme Ubc3 (also called Cdc34) and by an E3 ubiquitin ligase formed by three subunits: Cdc53, Skp1 and one of many F box proteins (reviewed in E. Patton et al., 1998, TIG. 25 14:6). The F box proteins (FBPs) are so called because they contain a motif, the F box, that was first identified in Cyclin F, and that is necessary for FBP interaction with Skp1 (Bai, et al., 1996, Cell 86:263). In addition, F box proteins also contain either WD-40 domains or Leucine-Rich Repeats (LRR) protein-protein interaction domains. Cdc53 (also called Cul A) and Skp1 appear to participate in the formation of at least three distinct E3, each 30 containing a different F box protein. Because these ligases are similar protein modules composed of Skp1, Cul A, and an F box protein, they have been named SCF. The interaction of the ligase with its substrates occurs via the F box subunit. The three SCFs identified so far in *S. cerevisiae* are: SCF^{Cdc4} (which recruits the Ck's Sic1 and Far1, the replication factor Cdc6, and the transcriptional activator Gcn4, as substrates through the F 35 box protein Cdc4), SCF^{Grr1} (which recruits the G1 cyclins Cln1 and Cln2 as substrates

through the F box protein GRR1), and SCF^{Met30} (which recruits the G1 cyclin Cln3 as a substrate throughout the F box protein MET30; see Pagano and Patton, supra, for recent reviews).

The intracellular level of the human Cki p27, a cell cycle regulated cyclin-dependent kinase (Cdk) inhibitor, is mainly regulated by degradation and it is known that the ubiquitin system controls p27 degradation (Pagano et al., 1995, Science 269:682). Similarly, degradation of other G1 human regulatory proteins (Cyclin E, Cyclin D1, p21, E2F, β -catenin) is controlled by the ubiquitin-pathway (reviewed in M. Pagano, supra). Yet, the specific enzymes involved in the degradation of G1 regulatory proteins have not been identified. A family of 6 genes (*CUL1*, 2, 3, 4a, 4b, and 5) homologous to *S. cerevisiae* cul A have been identified by searching the EST database (Kipreos, et al., 1996, Cell 85:829). Human Skp1 and the F box protein Skp2 (that contains five LRRs) were identified as two proteins associated in vivo with Cyclin A and thus designated as S-phase kinase-associated protein 1 and 2 (Zhang, et al., 1995, Cell 82:915). It has been demonstrated that phosphorylated p27 is specifically recognized by Skp2. Skp1 and Skp2 are also found to associate with Cul-1 and ROC1/Rbx1 to form an SCF ubiquitin ligase complex, SCF^{Skp2} ubiquitin ligase complex. While studies establish that p27 is targeted for degradation by the SCF^{Skp2} ubiquitin ligase complex, key factors involved in the degradation were unknown. It had been hypothesized that Nedd8, a highly conserved ubiquitin-like protein that is ligated to different cullins, is a necessary component for ligation of p27 (Podust, et al., 2000, Proc. Natl. Acad. Sci. USA 97:4579).

The highly conserved Suc1(suppressor of Cdc2 mutation)/Cks(cyclin-dependent kinase subunit) family of cell cycle regulatory proteins binds to some cyclin-dependent kinases and phosphorylated proteins and is essential for cell cycle progression. Suc1 (Hayles, et al., 1986, Mol. Gen. Genet. 202:291) and Cks1 (Hadwiger, et al., 1989, Mol. Cell Biol. 9:2034) were discovered in fission and budding yeast, respectively, as essential gene products that interact with cyclin-dependent kinases. Homologues from different species share extensive sequence conservation, and the two human homologues can functionally substitute for Cks1 in budding yeast (Richardson, et al. 1990, Genes and Dev. 4:1332). Crystal structures of the two human homologues and the fission yeast Suc1 have shown that they share a four-stranded β -sheet involved in binding to a Cdk catalytic subunit (Bourne, et al., 1996, Cell 84:863; Pines, J., 1996, Curr. Biol. 11:1399). In addition, they share a highly conserved phosphate-binding site, positioned on a surface contiguous to the Cdk catalytic site in the Cks-Cdk complex (Bourne, et al., 1996, Cell 84:863).

Cks proteins are involved in several cell cycle transitions, including the G1 to S-phase transition, entry into mitosis and exit from mitosis (Pines, J., 1996, *Curr. Biol.* 11:1399), but the molecular basis for their different actions is not well understood. With the exception of Cln2/Cln3-Cdk1 complexes from budding yeast being activated by Cks1 (Reynard, et al., 2000, *Mol. Cell Biol.* 20:5858), Cks proteins do not directly affect the catalytic activity of the cyclin-dependent kinase. However, Cks proteins can promote multi-site phosphorylations of some substrates by cyclin-dependent kinases. It has been proposed that by simultaneously binding to a partially phosphorylated protein and to a Cdk, Cks proteins increase the affinity of the kinase for the substrate and thus accelerate subsequent multiple phosphorylations (Pines, J., 1996, *Curr. Biol.* 11:1399). Indeed, Cks proteins promote Cdk-catalyzed multiple phosphorylations of subunits of the cyclosome/APC (Patra, D. & Dunphy, W.G., 1998, *Genes Dev.* 12:2549; Shteinberg, M. & Hershko, A., 1999, *Biochem. Biophys. Res. Commun.* 257:12), as well as G2/M regulators such as Cdc25, Myt1 and Wee1 (Patra, et al., 1999, *J. Biol. Chem.* 274:36839).

2.4 DEREGULATION OF THE UBIQUITIN PATHWAY IN CANCER AND OTHER PROLIFERATIVE DISORDERS

Cancer develops when cells multiply too quickly. Cell proliferation is determined by the net balance of positive and negative signals. When positive signals overcome or when negative signals are absent, the cells multiply too quickly and cancer develops.

Ordinarily cells precisely control the amount of any given protein and eliminate the excess or any unwanted protein. To do so, the cell specifically tags the undesired protein with a long chain of molecules called ubiquitin. These molecules are then recognized and destroyed by a complex named proteasome. However, all this mechanism goes awry in tumors leading to the excessive accumulation of positive signals (oncogenic proteins), or resulting in the abnormal degradation of negative regulators (tumor suppressor proteins). Thus, without tumor suppressor proteins or in the presence of too much oncogenic proteins, cells multiply ceaselessly, forming tumors (reviewed by Ciechanover, 1998, *EMBO J.* 17: 7151; Spataro, 1998, *Br. J. Cancer* 77: 448). For example, abnormal ubiquitin-mediated degradation of the p53 tumor suppressor (reviewed by J. Brown and M. Pagano, 1997, *Biochim. Biophys. Acta* 1332: 1), the putative oncogene β -catenin (reviewed by Peifer, 1997, *Science* 275:1752) and the Cki p27 (reviewed in Ciechanover, supra; Spataro, supra; Lloyd, 1999, *Am. J. Pathol.* 154: 313) have been correlated with

tumorigenesis, opening to the hypothesis that some genes encoding ubiquitinating enzymes may be mutated in tumors.

Initial evidence indicates that human F-box proteins play a role in the ubiquitination of G1 regulatory proteins as their homologues do in yeast (see below).

- 5 Unchecked degradation of cell cycle regulatory proteins has been observed in certain tumors and it is possible that deregulated ubiquitin ligase play a role in the altered degradation of cell cycle regulators. A well understood example is that of Mdm2, a ubiquitin ligase whose overexpression induces low levels of its substrate, the tumor suppressor p53.

10

3. SUMMARY OF THE INVENTION

The present invention relates to novel F box proteins and therapeutic protocols and pharmaceutical compositions designed to target the novel F box proteins and their interactions with substrates for the treatment of proliferative and differentiative
15 disorders. The present invention also relates to screening assays to identify substrates of the novel F box proteins and to identify agents which modulate or target the novel ubiquitin ligases and interactions with their substrates. The invention further relates to screening assays based on the identification of novel substrates of known F box proteins, such as the two novel substrates of the known F box protein Skp2, E2F and p27. The screening assays
20 of the present invention may be used to identify potential therapeutic agents for the treatment of proliferative or differentiative disorders and other disorders that related to levels of expression or enzymatic activity of F box proteins.

The invention is based in part, on the Applicants' discovery, identification and characterization of nucleic acids comprising nucleotide sequences that encode novel
25 ubiquitin ligases with F box motifs. These twenty-six novel substrate-targeting subunits of ubiquitin ligase complexes, FBP1, FBP2, FBP3a, FBP3b, FBP4, FBP5, FBP6, FBP7, FBP8, FBP9, FBP10, FBP11, FBP12, FBP13, FBP14, FBP15, FBP16, FBP17, FBP18, FBP19, FBP20, FBP21, FBP22, FBP23, FBP24, and FBP25, described herein, were first identified based on their interaction with components of the ubiquitin ligase complex (FBP1, FBP2,
30 FBP3a, FBP4, FBP5, FBP6 and FBP7) or by sequence comparison of these proteins with nucleotide sequences present in DNA databases (FBP3b, FBP8, FBP9, FBP10, FBP11, FBP12, FBP13, FBP14, FBP15, FBP16, FBP17, FBP18, FBP19, FBP20, FBP21, FBP22, FBP23, FBP24, and FBP25). These novel substrate-targeting subunits of ubiquitin ligase complexes each contain an F box motif through which they interact with the other
35 components of the ubiquitin ligase complex. In addition, some of these FBPs contain WD-

40 domains and LRRs (which appear to be involved in their interaction with substrates), while other FBPs contain potential protein-protein interaction modules not yet identified in FBPs, such as leucine zippers, ring fingers, helix-loop-helix motifs, proline rich motifs and SH2 domains. The invention is also based, in part, on the Applicants' discovery and
5 identification of FBP specific substrates p27 and β -catenin and on methods to identify novel FBP substrates. Some of the genes encoding the novel F box proteins were also mapped to chromosome sites frequently altered in breast, prostate and ovarian cancer, nasopharyngeal and small cell lung carcinomas, gastric hepatocarcinomas, Burkitt's lymphoma and parathyroid adenomas. Finally, the invention is also based, in part, on the Applicants'
10 generation of transgenic mice expressing wild type or dominant negative versions of FBP proteins and on the generation of FBP knock-out mice.

The invention encompasses the following nucleotide sequences, host cells expressing such nucleotide sequences, and the expression products of such nucleotide sequences: (a) nucleotide sequences that encode mammalian FBP1, FBP2, FBP3a, FBP3b,
15 FBP4, FBP5, FBP6, FBP7, FBP8, FBP11, FBP12, FBP13, FBP14, FBP15, FBP17, FBP18, FBP20, FBP21, FBP22, FBP23, and FBP25, including the human nucleotides, and their gene products; (b) nucleotides that encode portions of the novel substrate-targeting subunits of ubiquitin ligase complexes, and the polypeptide products specified by such nucleotide sequences, including but not limited to F box motifs, the substrate binding domains; WD-40
20 domains; and leucine rich repeats, *etc.*; (c) nucleotides that encode mutants of the novel ubiquitin ligases in which all or part of the domain is deleted or altered, and the polypeptide products specified by such nucleotide sequences; (d) nucleotides that encode fusion proteins containing the novel ubiquitin ligases or one of its domains fused to another polypeptide.

The invention further encompasses agonists and antagonists of the novel
25 substrate-targeting subunits of ubiquitin ligase complexes, including small molecules, large molecules, mutants that compete with native F box binding proteins, and antibodies as well as nucleotide sequences that can be used to inhibit ubiquitin ligase gene expression (*e.g.*, antisense and ribozyme molecules, and gene regulatory or replacement constructs) or to enhance ubiquitin ligase gene expression (*e.g.*, expression constructs that place the ubiquitin
30 ligase gene under the control of a strong promoter system), and transgenic animals that express a ubiquitin ligase transgene or knock-outs that do not express the novel ubiquitin ligases.

Further, the present invention also relates to methods for the use of the genes and/or gene products of novel substrate-targeting subunits of ubiquitin ligase complexes for
35 the identification of compounds which modulate, *i.e.*, act as agonists or antagonists, of

ubiquitin ligase activity. Such compounds can be used as agents to control proliferative or differentiative disorders, *e.g.* cancer. In particular, the present invention encompasses methods to inhibit the interaction between β -catenin and FBP1 or p27 and Skp2. In fact, agents able to block these interactions can be used to modulate cell proliferation and/or growth.

Still further, the invention encompasses screening methods to identify derivatives and analogues of the novel substrate-targeting subunits of ubiquitin ligase complexes which modulate the activity of the novel ligases as potential therapeutics for proliferative or differentiative disorders. The invention provides methods of screening for proteins that interact with novel components of the ubiquitin ligase complex, including FBP1, FBP2, FBP3a, FBP3b, FBP4, FBP5, FBP6, FBP7, FBP8, FBP9, FBP10, FBP11, FBP12, FBP13, FBP14, FBP15, FBP16, FBP17, FBP18, FBP19, FBP20, FBP21, FBP22, FBP23, FBP24, and FBP25 or derivatives, fragments or domains thereof, such as the F box motif. In accordance with the invention, the screening methods may utilize known assays to identify protein-protein interactions including phage display assays or the yeast two-hybrid assay system or variations thereof.

In addition, the present invention is directed to methods that utilize FBP gene sequences and/or FBP gene product sequences for the diagnostic evaluation, genetic testing and/or prognosis of an FBP-related disorder, such as a proliferative disorder. For example, the invention relates to methods for diagnosing FBP-related disorders, *e.g.*, proliferative disorders, wherein such methods can comprise measuring FBP gene expression in a patient sample, or detecting an FBP mutation that correlates with the presence or development of such a disorder, in the genome of a mammal suspected of exhibiting such a disorder. In particular, the invention encompasses methods for determining if a subject (*e.g.*, a human patient) is a risk for a disorder characterized by one or more of: (i) a mutation of an FBP gene encoding a protein represented in part A of Figures 3-28, or a homologues thereof; (ii) the mis-expression of an FBP gene; (iii) the mis-expression of an FBP protein.

The invention is illustrated by way of working examples which demonstrate the identification and characterization of the novel substrate-targeting subunits of ubiquitin ligase complexes. The working examples of the present invention further demonstrate the identification of the specific interaction of (i) FBP1 with β -catenin and (ii) the known FBP, Skp2, with the cell-cycle regulatory proteins E2F and p27 and the cell cycle protein Cks1. These interactions suggest that β -catenin is a specific substrate of FBP1, while E2F and p27 are substrates of Skp2 and Cks1 is a mediator for Skp2 and p27. In fact, the working examples of the present invention further demonstrate that β -catenin is a specific substrate

of FBP1, while p27 is substrates of Skp2 and Cks1 binds to both p27 and Skp2. The identification of proteins interacting with the novel FBPs will be possible using the methods described herein or with a different approach.

5 3.1 DEFINITIONS

As used herein, the term "F-box motif" refers to a stretch of approximately 40 amino acid that was identified as being necessary for the interaction of F-box containing proteins with Skp1. The consensus sequence of an F-box motif is described in Bai et al., 1996, Cell 86:263-274, incorporated herein by reference in its entirety.

10 As used herein the term "F-box protein" (FBP) refers to peptide, polypeptide or protein which contains an F-box motif.

Although, FBPs are substrate-targeting subunits of ubiquitin ligase complexes, as used herein the term "ubiquitin ligase" refers to a peptide, polypeptide or protein that contains an F-box motif and interacts with Skp1.

15 As used herein, the term "functionally equivalent to an FBP gene product" refers to a gene product that exhibits at least one of the biological activities of the endogenous FBP gene product. For example, a functionally equivalent FBP gene product is one that is capable of interacting with Skp1 so as to become associated with a ubiquitin ligase complex. Such a ubiquitin ligase complex may be capable of ubiquitinating a specific
20 cell-cycle regulatory protein, such as a cyclin or cki protein.

As used herein, the term "to target" means to inhibit, block or prevent gene expression, enzymatic activity, or interaction with other cellular factors.

As used herein, the term "therapeutic agent" refers to any molecule, compound or treatment that alleviates or assists in the treatment of a proliferative disorder
25 or related disorder.

As used herein, the terms "WD-40 domain", "Leucine Rich Repeat", "Leucine Zipper", "Ring finger", "Helix-loop-helix motif", "Proline rich motif", and "SH2 domain" refer to domains potentially involved in mediating protein-protein interactions. The "WD-40 domain" refers to a consensus sequence of forty amino acid repeats which is
30 rich in tryptophan and aspartic acid residues and is commonly found in the beta subunits of trimeric G proteins (see Neer et al., 1994 Nature 371:297-300 and references therein, which are incorporated herein by reference in their entirety). An "LRR" or a "Leucine Rich Repeat" is a leucine rich sequence also known to be involved in mediating protein-protein interactions (see Kobe & Deisenhofer, 1994, Trends. Biochem. Sci. 19:415-421 which are
35 incorporated herein by reference in their entirety). A "leucine zipper" domain refers to a

domain comprising a stretch of amino acids with a leucine residue in every seventh position which is present in a large family of transcription factors (see Landshultz et al., 1988, Science 240:1759-64; see also Sudol et al., 1996, Trends Biochem. 21:1-3, and Koch et al., 1991, Science 252:668-74).

5

4. BRIEF DESCRIPTION OF THE FIGURES

Figure 1. Alignment of the conserved F-box motif amino acid residues in the human F-box proteins FBP1 (SEQ ID NO:15), FBP2 (SEQ ID NO:16), FBP3a (SEQ ID NO:17), FBP3b (SEQ ID NO:78), FBP4 (SEQ ID NO:18), FBP5 (SEQ ID NO:19), FBP6 (SEQ ID NO:20), FBP7 (SEQ ID NO:21), Skp2 (SEQ ID NO:22), FBP8 (SEQ ID NO:61), FBP9 (SEQ ID NO:62), FBP10 (SEQ ID NO:63), FBP11 (SEQ ID NO:64), FBP12 (SEQ ID NO:65), FBP13 (SEQ ID NO:79), FBP14 (SEQ ID NO:66), FBP15 (SEQ ID NO:67), FBP16 (SEQ ID NO:68), FBP17 (SEQ ID NO:69), FBP18 (SEQ ID NO:70), FBP19 (SEQ ID NO:71), FBP20 (SEQ ID NO:72), FBP21 (SEQ ID NO:73), FBP22 (SEQ ID NO:74), FBP23 (SEQ ID NO:75), FBP24 (SEQ ID NO:76), FBP25 (SEQ ID NO:77). Alignment of the F-boxes of a previously known FBP, Skp2, with the F-boxes of FBPs identified through a two-hybrid screen (designated by the pound symbol) or BLAST searches (designated by a cross) was performed using the Clustal W method (MacVector(tm)) followed by manual re-
adjustment. Identical residues in at least 15 F-boxes are shaded in dark gray, while similar residues are shaded in light gray. One asterisk indicates the presence in the cDNA of a STOP codon followed by a polyA tail, while potential full length clones are designated with two asterisks. The asterisks on the bottom of the figure indicate the amino acid residues mutated in FBP3a (see Figure 29).

25

FIG. 2. Schematic representation of FBPs. Putative protein-protein interaction domains in human FBPs are represented (see key-box for explanation). FBPs identified by a two-hybrid screen are designated by the pound symbol, FBPs identified through BLAST searches by a cross. The double slash indicates that the corresponding cDNAs are incomplete at the 5' end; the asterisks indicate the presence in the cDNA of a STOP codon followed by a polyA tail.

30

FIG. 3 A-B. A. Amino acid sequence of human F-box protein FBP1 (SEQ ID NO:2). B. Corresponding cDNA (SEQ ID NO:1).

35

FIG. 4 A-B. **A.** Amino acid sequence of human F-box protein FBP2 (SEQ ID NO:4). **B.** Corresponding cDNA (SEQ ID NO:3).

FIG. 5 A-B. **A.** Amino acid sequence of human F-box protein FBP3a (SEQ ID NO:6). **B.** Corresponding cDNA (SEQ ID NO:5).

FIG. 6 A-B. **A.** Amino acid sequence of human F-box protein FBP3b (SEQ ID NO:24). **B.** Corresponding cDNA (SEQ ID NO:23).

FIG. 7 A-B. **A.** Amino acid sequence of human F-box protein FBP4 (SEQ ID NO:8). **B.** Corresponding cDNA (SEQ ID NO:7).

FIG. 8 A-B. **A.** Amino acid sequence of human F-box protein FBP5 (SEQ ID NO:10). **B.** Corresponding cDNA (SEQ ID NO:9).

FIG. 9 A-B. **A.** Amino acid sequence of human F-box protein FBP6 (SEQ ID NO:12). **B.** Corresponding cDNA (SEQ ID NO:11).

FIG. 10 A-B. **A.** Amino acid sequence of human F-box protein FBP7 (SEQ ID NO:14). **B.** Corresponding cDNA (SEQ ID NO:13).

FIG. 11 A-B. **A.** Amino acid sequence of human F-box protein FBP8 (SEQ ID NO:26). **B.** Corresponding cDNA (SEQ ID NO:25).

FIG. 12 A-B. **A.** Amino acid sequence of human F-box protein FBP9 (SEQ ID NO:28). **B.** Corresponding cDNA (SEQ ID NO:27).

FIG. 13 A-B. **A.** Amino acid sequence of human F-box protein FBP10 (SEQ ID NO:30). **B.** Corresponding cDNA (SEQ ID NO:29).

FIG. 14 A-B. **A.** Amino acid sequence of human F-box protein FBP11 (SEQ ID NO:32). **B.** Corresponding cDNA (SEQ ID NO:31).

FIG. 15 A-B. **A.** Amino acid sequence of human F-box protein FBP12 (SEQ ID NO:34). **B.** Corresponding cDNA (SEQ ID NO:33).

FIG. 16 A-B. **A.** Amino acid sequence of human F-box protein FBP13 (SEQ ID NO:36). **B.** Corresponding cDNA (SEQ ID NO:35).

5 FIG. 17 A-B. **A.** Amino acid sequence of human F-box protein FBP14 (SEQ ID NO:38). **B.** Corresponding cDNA (SEQ ID NO:37).

FIG. 18 A-B. **A.** Amino acid sequence of human F-box protein FBP15 (SEQ ID NO:40). **B.** Corresponding cDNA (SEQ ID NO:39).

10 FIG. 19 A-B. **A.** Amino acid sequence of human F-box protein FBP16 (SEQ ID NO:42). **B.** Corresponding cDNA (SEQ ID NO:41).

15 FIG. 20 A-B. **A.** Amino acid sequence of human F-box protein FBP17 (SEQ ID NO:44). **B.** Corresponding cDNA (SEQ ID NO:43).

FIG. 21 A-B. **A.** Amino acid sequence of human F-box protein FBP18 (SEQ ID NO:46). **B.** Corresponding cDNA (SEQ ID NO:45).

20 FIG. 22 A-B. **A.** Amino acid sequence of human F-box protein FBP19 (SEQ ID NO:48). **B.** Corresponding cDNA (SEQ ID NO:47).

FIG. 23 A-B. **A.** Amino acid sequence of human F-box protein FBP20 (SEQ ID NO:50). **B.** Corresponding cDNA (SEQ ID NO:49).

25 FIG. 24 A-B. **A.** Amino acid sequence of human F-box protein FBP21 (SEQ ID NO:52). **B.** Corresponding cDNA (SEQ ID NO:51).

30 FIG. 25 A-B. **A.** Amino acid sequence of human F-box protein FBP22 (SEQ ID NO:54). **B.** Corresponding cDNA (SEQ ID NO:53).

FIG. 26 A-B. **A.** Amino acid sequence of human F-box protein FBP23 (SEQ ID NO:56). **B.** Corresponding cDNA (SEQ ID NO:55).

35 FIG. 27 A-B. **A.** Amino acid sequence of human F-box protein FBP24 (SEQ ID NO:58). **B.** Corresponding cDNA (SEQ ID NO:57).

FIG. 28A-B. **A.** Amino acid sequence of human F-box protein FBP25 (SEQ ID NO:60). **B.** Corresponding cDNA (SEQ ID NO:59).

FIG. 29. FBPs interact specifically with Skp1 through their F-box. The
 5 cDNAs of FBPs (wild type and mutants) were transcribed and translated in vitro (IVT) in the presence of 35S- methionine. Similar amounts of IVT proteins (indicated at the top of each lane) were subjected to a histidine-tagged pull-down assay using Nickel-agarose beads to which either His-tagged-Skp1 (lanes 1, 3, 4, 6-10, 12, 15, 17, 19 and 21), His-tagged-Elongin C (lanes 2, 5, 11, 14, 16, 18, 19 and 22), or His-tagged p27 (lane 12) were pre-
 10 bound. Bound IVT proteins were analyzed by SDS-PAGE and autoradiography. The arrows on the left side of the panels point to the indicated FBPs. The apparent molecular weights of the protein standards are indicated on the right side of the panels.

FIG. 30. FBP1, FBP2, FBP3a, FBP4 and FBP7 form novel SCFs with
 15 endogenous Skp1 and Cull1 in vivo. HeLa cells were transfected with mammalian expression plasmids encoding Flag-tagged versions of FBP1 (lane 1), (Δ F)FBP1 (lane 2), FBP4 (lane 3), FBP7 (lane 5), FBP2 (lane 7), (Δ F)FBP2 (lane 8), FBP3a (lane 9), (Δ F)FBP3a (lane 10), or with an empty vector (lanes 4 and 6). Cells were lysed and extracts were subjected to immunoprecipitation with a rabbit anti-Flag antibody (lanes 1-8).
 20 Immunoprecipitates were then immunoblotted with a mouse anti-Cul1 monoclonal antibody, a rabbit anti-Skp1 polyclonal antibody or a rabbit anti-Cul2 polyclonal antibody, as indicated. The last lane contains 25 μ g of extracts from non-transfected HeLa cells; lane 9 contains recombinant Cul1, Skp1, or Cul2 proteins used as markers. The slower migrating bands detected with the antibodies to Cul1 and Cul2 are likely generated by the
 25 covalent attachment of a ubiquitin-like molecule to these two cullins, as already described for the yeast cullin Cdc53 and mammalian Cul4a.

FIG. 31. FBP1, FBP2, FBP3a, FBP4 and FBP7 associate with a ubiquitin
 ligase activity. HeLa cells were transfected with mammalian expression plasmids encoding
 30 human Skp1, Cul1 and Flag-tagged versions of FBP1 (lane 3), (Δ F)FBP1 (lane 4), FBP2 (lanes 2 and 5), (Δ F)FBP2 (lane 6), FBP7 (lane 7), FBP3a (lanes 8 and 13), (Δ F)FBP3a (lane 9), a non relevant Flag-tagged protein (Irf3, lane 10), FBP4 (lanes 11 and 12) or with an empty vector (lane 1). Cells were lysed and extracts were subjected to
 35 immunoprecipitation with a rabbit anti-Flag antibody. Immunoprecipitates were incubated in the presence of purified recombinant E1 and Ubc4 (lanes 1-11) or Ubc2 (lanes (12 and

13) and a reaction mix containing biotinylated ubiquitin. Reaction in lane 2 contained also NEM. Ubiquitinated proteins were visualized by blotting with HRP-streptavidin. The bracket on the left side of the panels marks a smear of ubiquitinated proteins produced in the reaction, the asterisk indicates ubiquitin conjugated with E1 that were resistant to
 5 boiling.

FIG. 32. Subcellular localization of FBPs. HeLa cells were transfected with mammalian expression plasmids encoding Flag-tagged versions of FBP1 (a-b), FBP2 (c-d), FBP3a (e-f), FBP4 (g-h), (DF)FBP2 (i-j), or (Δ F)FBP3a (k-l). After 24 hours, cells were
 10 subjected to immunofluorescence with a rabbit anti-Flag antibody (a, c, e, g, i, k) to stain FBPs and bisbenzimidazole (b, d, f, h, j, l) to stain nuclei.

FIG. 33. Abundance of FBP transcripts in human tissues. Membranes containing electrophoretically fractionated poly(A)⁺ mRNA from different human tissues
 15 were hybridized with specific probes prepared from FBP1, FBP2, FBP3a, FBP4, SKP2, and β -ACTIN cDNAs. The arrows on the left side of the figure point to the major transcripts as described in the text.

FIG. 34 A-E. FISH localization of FBP genes. Purified phage DNA
 20 containing a genomic probe was labeled with digoxigenin dUTP and detected with Cy3-conjugated antibodies. The signals corresponding to the locus of the genomic probe (red) are seen against the DAPI-Actinomycin D stained normal human chromosomes (blue-white). Panel **A** shows localization of FBP1 to 10q24, **B** shows localization of FBP2 to 9q34, **C** shows localization of FBP3a to 13q22, **D** shows localization of FBP4 to 5p12, and
 25 **E** shows localization of FBP5 to 6q25-26. Arrows point to FBP-specific FISH signals.

FIG. 35A-C. FBP1 associates with β -catenin. **A.** Extracts from baculovirus-infected insect cells expressing either β -catenin alone (lane 1) or in combination with Flag-tagged FBP1 (lane 2) were immunoprecipitated (IP) with a rabbit anti-Flag antibody (α -Flag), followed by immunoblotting with anti-Flag ($m\alpha$ -Flag) and anti- β -catenin mouse antibodies, as indicated. Lanes 3 and 4 contain 25 μ g of extracts from infected insect cells immunoblotted with the same antibodies. **B.** Extracts from baculovirus-infected insect cells expressing cyclin D1, Flag-FBP1 in the absence (lanes 1-3) or in the presence of Skp1 (lanes 4-6) were immunoprecipitated with normal rabbit IgG (r-IgG, lanes 1 and 4), rabbit
 30 anti-Flag antibody @ α -Flag, lanes 2 and 5), or rabbit anti-cyclin D1 antibody @ α -D1, lanes
 35

3 and 6). Immunoprecipitates were then immunoblotted with anti-Flag (α -Flag) and cyclin D1 (α -D1) mouse antibodies, as indicated. The last lane contains 25 μ g of a representative extract from infected insect cells immunoblotted with the same antibodies. C. 293 cells were transfected with mammalian expression plasmids encoding HA-tagged β -catenin alone or in combination with either Flag-tagged FBP1 or Flag-tagged (Δ F)FBP1. Cells were lysed and extracts were subjected to immunoprecipitation with a rabbit anti-Flag antibody @ α -Flag, lanes 4-6) and immunoblotted with rat anti-HA (α -HA) and mouse anti-Flag (α -Flag) antibodies, as indicated. The first three lanes contain 25 μ g of extracts from transfected 293 cells immunoblotted with the same antibodies. Transfecting high levels of β -catenin expression vector, the associations of β -catenin with FBP1 and (Δ F)FBP1 could be determined independently of β -catenin levels.

FIG. 36 A-B. Stabilization of β -catenin by a dominant negative (Δ F)FBP1 mutant. A. Human 293 cells were transfected with mammalian expression plasmids encoding HA-tagged β -catenin alone or in combination with either Flag-tagged (Δ F)FBP1 or Flag-tagged (Δ F)FBP2. Cells were lysed and extracts were subjected to immunoblotting with rat anti-HA and rabbit anti-Flag @ α -Flag) antibody, as indicated. B. Pulse chase analysis of β -catenin turnover rate. HA-tagged β -catenin in combination with either an empty vector, FBP1, or (Δ F)FBP1 was co-transfected in 293 cells. 24 hours later cells were labeled with 35S-methionine for 30 minutes and chased with medium for the indicated times. Extracts were then subjected to immunoprecipitation with a rat anti-HA antibody.

FIG. 37A-C. Binding of phosphorylated p27 to Skp2. A. A panel of in vitro translated [35S]FBPs were used in binding reactions with beads coupled to the phospho-peptide NAGSVEQT*PKKPGLRRRQT, corresponding to the carboxy terminus of the human p27 with a phosphothreonine at position 187 (T*). Beads were washed with RIPA buffer and bound proteins were eluted and subjected to electrophoresis and autoradiography (Upper Panel). Bottom Panel: 10% of the in vitro translated [35S]FBP inputs. B. HeLa cell extracts were incubated with beads coupled to the phospho-p27 peptide (lane 2), an identical except unphosphorylated p27 peptide (lane 1) or the control phospho-peptide AEIGVGAY*GTVYKARDPHS, corresponding to an amino terminal peptide of human Cdk4 with a phosphotyrosine at position 17 (Y*) (lane 3). Beads were washed with RIPA buffer and bound proteins were immunoblotted with antibodies to the proteins indicated on the left of each panel. A portion of the HeLa extract (25 μ g) was used as a control (lane 4). The slower migrating band in Cull1 is likely generated by the covalent

attachment of a ubiquitin-like molecule, as already described for other cullins 48. C. One μ l of in vitro translated [35S] wild type p27 (WT, lanes 1-4) or p27(T187A) mutant (T187A, lanes 5-6) were incubated for 30 minutes at 30°C in 10 μ l of kinase buffer. Where indicated, ~2.5 pmole of recombinant purified cyclin E/Cdk2 or ~1 pmole Skp2 (in
 5 Skp1/Skp2 complex) were added. Samples were then incubated with 6 μ l of Protein-A beads to which antibodies to Skp2 had been covalently linked. Beads were washed with RIPA buffer and bound proteins subjected to electrophoresis and autoradiography. Lanes 1-6: Skp2-bound proteins; Lanes 7 and 8: 7.5% of the in vitro translated [35S] protein inputs.

10

FIG. 38. In vivo binding of Skp2 to p27. Extracts from HeLa cells (lanes 1-2 and 5-6) or IMR90 fibroblasts (lanes 9-10) were immunoprecipitated with different affinity purified (AP) antibodies to Skp2 or with purified control IgG fractions. Lane 1: extract immunoprecipitated with a goat IgG (G-IgG); lane 2: with an AP goat antibody to an
 15 N-terminal Skp2 peptide (G- α -Skp2); lanes 5 and 9: with a rabbit IgG (R-IgG); lanes 6 and 10: with an AP rabbit antibody to Skp2 (R- α -Skp2). Immunoprecipitates were immunoblotted with antibodies to the proteins indicated on the left of each panel. Lanes 1-4 in the bottom panel were immunoblotted with a phospho-site p27 specific antibody. Lanes 3, 7, and 11 contain 25 μ g of cell extracts; Lanes 4, 8, and 12 contain the relevant
 20 recombinant proteins used as markers. The altered migration of some markers is due to the presence of tags on the recombinant proteins.

FIG. 39 A-B. Skp2 and cyclin E/Cdk2 complex are rate-limiting for p27 ubiquitination in G1 extracts. A. In vitro ubiquitin ligation (lanes 1-12 and 17-20) and
 25 degradation (lanes 13-16) of p27 were carried out with extracts from asynchronously growing (Asyn. ext., lanes 2-3) or G1-arrested (G1 ext., lanes 4-20) HeLa cells. Lane 1 contains no extract. Recombinant purified proteins were supplemented as indicated. Reactions were performed using wild-type p27 (lanes 1-18) or p27(T187A) mutant (T187A, lanes 19-20). Lanes 1-8, 9-12, and 17-20 are from three separate experiments. The bracket
 30 on the left side of the panels marks a ladder of bands >27,000 corresponding to polyubiquitinated p27. The asterisk indicates a non-specific band present in most samples. B. Immunoblot analysis of levels of Skp2 and p27 in extracts from asynchronous (lane 1) or G1-arrested (lane 2) HeLa cells.

35

FIG. 40 A-C. Skp2 is required for p27-ubiquitin ligation activity. **A.** Immunodepletion. Extracts from asynchronous HeLa cells were untreated (lane 2) or immunodepleted with pre-immune serum (lane 3), anti-Skp2 antibody pre-incubated with 2 μ g of purified GST (lane 4), or anti-Skp2 antibody pre-incubated with 2 μ g of purified GST-Skp2 (lane 5). Lane 1 contains no extract. Samples (30 μ g of protein) were assayed for p27 ubiquitination in the presence of cyclin E/Cdk2. The bracket on the left side of the panels marks a ladder of bands >27,000 corresponding to polyubiquitinated p27. The asterisk indicates a non-specific band present in all samples. **B.** Reconstitution. The restoration of p27 ubiquitination activity in Skp2-immunodepleted extracts was tested by the addition of the indicated purified proteins. All samples contained 30 μ g of Skp2-depleted extract (Skp2-depl. ext.) and cyclin E/Cdk2. **C.** Immunoprecipitation. Extracts from asynchronous HeLa cells were immunoprecipitated with a rabbit anti-Skp2 antibody (lanes 3 and 5) or pre-immune serum (PI, lanes 2 and 4). Total extract (lane 1) and immuno-beads (lanes 2-5) were added with p27, recombinant purified cyclin E/Cdk2 and ubiquitination reaction mix. Samples in lanes 4 and 5 were supplemented with recombinant purified E1 and Ubc3. All samples were then assayed for p27 ubiquitination.

FIG. 41 A-B. In vivo role of Skp2 in p27 degradation. **A.** Stabilization of p27 by a dominant negative (Δ F)Skp2 mutant in vivo. NIH-3T3 cells were transfected with mammalian expression vectors encoding human p27 alone (lane 2), p27 in combination with either (Δ F)Skp2 (lane 3), or (Δ F)FBP1 (lane 4). Lane 1: untransfected cells. Cells were lysed and extracts were subjected to immunoblotting with antibodies to p27, Skp2 or Flag [to detect Flag-tagged (Δ F)FBP1]. Exogenous human p27 protein migrates more slowly than the endogenous murine p27. **B.** Pulse chase analysis of p27 turnover rate. Human p27 in combination with either an empty vector, or (Δ F)Skp2 was transfected in NIH-3T3 cells. Twenty-four hours later, cells were labeled with [35 S]-methionine for 20 minutes and chased with medium for the indicated times. Extracts were then subjected to immunoprecipitation with a mouse anti-p27 antibody.

FIG. 42. Stabilization of cellular p27 by antisense oligonucleotides targeting SKP2 mRNA. HeLa cells were treated for 16 -18 hours with two different anti-sense oligodeoxynucleotides (AS) targeting two different regions of SKP2 mRNA. Lanes 2, 6, 12 and 16: AS targeting the N-terminal SKP2 region (NT); Lanes 4 and 8: AS targeting the C-terminal SKP2 region (CT); Lanes 1, 3, 5, 7, 11 and 15: control oligodeoxynucleotides pairs (Ctrl). Lanes 1-4, and 5-8 are from two separate experiments. Lanes 11-12 and 15-16:

HeLa cells were blocked in G1/S with either Hydroxyurea or Aphidicolin treatment respectively, for 24 hours. Cells were then transfected with oligodeoxynucleotides, lysed after 12 hours (before cells had re-entered G1) and immunoblotted with antibodies to Skp2 (top panels) and p27 (bottom panels). Lanes 9 and 13: Untransfected HeLa cells; Lanes 10 and 14: Untransfected HeLa cells treated with drugs as transfected cells.

FIG. 43 A-C. Timing of Skp2 action in the process of p27 degradation. **A.** IMR90 fibroblasts were synchronized in G0/G1 by serum deprivation, reactivated with serum, and sampled at the indicated intervals. Protein extracts were analyzed by immunoblot with the antibodies to the indicated proteins. The Skp2 doublet was likely generated by phosphorylation since was consistently observed using a 12.5% gel only when cell lysis was performed in the presence of okadaic acid. **B.** HeLa cells blocked in mitosis with nocodazole were shaken off, released in fresh medium and sampled at the indicated intervals. Protein extracts were analyzed by immunoblotting with the antibodies to the indicated proteins. **C.** Extracts from G1 (3 hours after release from nocodazole block) (lane 1) and S-phase (12 hours after release from the nocodazole block) (lane 2) HeLa cells were either immunoprecipitated with an anti-p27 antibody (top two panels) or with an anti-Skp2 antibody (bottom three panels) and then immunoblotted with the antibodies to the indicated proteins.

Fig. 44. The heat-stable factor is sensitive to trypsin action. Heat-treated Fraction 1 (~ 0.1 mg/ml) was incubated at 37°C for 60 min with 50 mM Tris-HCl (pH 8.0) either in the absence (lane 1) or in the presence of 0.6 mg/ml of TPCK-treated trypsin (Sigma T8642) (lane 2). Trypsin action was terminated by the addition of 2 mg/ml of soybean trypsin inhibitor (STI). In lane 3, STI was added 5 min prior to a similar incubation with trypsin. Subsequently, samples corresponding to ~50 ng of heat-treated Fraction 1 were assayed for the stimulation of p27-ubiquitin ligation.

Fig. 45 A-C. The heat-stable factor is not Nedd8 and is required following the modification of Cul-1 by Nedd8. **A.** Purified Nedd8 does not replace the factor in the stimulation of p27-ubiquitin ligation. Where indicated, ~50 ng of heat-treated Fraction 1 or 100 ng of purified recombinant human Nedd8 were added to the p27-MeUb ligation assay. **B.** Ligation of Nedd8 to Cul-1. Cul-1/ROC1 (3 µl) was incubated with Nedd8 (10 µg) and purified Nedd8-conjugating enzymes (20 µl) in a 100 -µl reaction mixture containing Tris (pH 7.6), MgCl₂, ATP, phosphocreatine, creatine phosphokinase, DTT, glycerol and STI at

concentrations similar to those described for the p27-ubiquitin ligation assay. A control preparation of Cul1/ROC1 was incubated under similar conditions, but without Nedd8 conjugating enzymes. Following incubation at 30°C for 2 hours, samples of control (lane 1) or Nedd8-modified (lane 2) preparations were separated on an 8% polyacrylamide-SDS gel and immunoblotted with an anti-Cul-1 antibody (Zymed). C. SCF^{Skp2} complex containing Nedd8-modified Cul-1 still requires the factor from Fraction 1 for p27-ubiquitin ligation. p27-MeUb ligation was assayed, except that ³⁵S-labeled p27 was replaced by bacterially expressed purified p27 (20 ng), and Cul-1/ROC1 was replaced by 2 µl of the unmodified or Nedd8-modified Cul-1/ROC1 preparations. Following incubation (30°C, 60 min), samples were separated on a 12.5% polyacrylamide-SDS gel, transferred to nitrocellulose and blotted with an anti-p27 monoclonal antibody (Transduction Laboratories). A cross-reacting protein is labeled by an asterisk.

Fig. 46 A, B. Purification of the factor required for p27-ubiquitin ligation and its identification as Cks1. A. Last step of purification by gel filtration chromatography. The peak of active material from the MonoS step was applied to a Superdex 75 HR 10/30 column (Pharmacia) equilibrated with 20 mM Tris-HCl (pH 7.2), 150 mM NaCl, 1 mM DTT and 0.1% Brij-35. Samples of 0.5 ml were collected at a flow rate of 0.4 ml/min. Column fractions were concentrated to a volume of 50 µl by centrifuge ultrafiltration (Centricon-10, Amicon). Samples of 0.004 µl of column fractions were assayed for activity to stimulate p27-ubiquitin ligation. Results were quantified by phosphorimager analysis and were expressed as the percentage of ³⁵S-p27 converted to ubiquitin conjugates. Arrows at top indicate the elution position of molecular mass marker proteins (kDa). B. Silver staining of samples of 2.5 µl from the indicated fractions of the Superdex 75 column, resolved on a 16% polyacrylamide-SDS gel. Numbers on the right indicate the migration position of molecular mass marker proteins (kDa).

Fig. 47. All bacterially expressed Cks/Suc1 proteins stimulate the multi-phosphorylation of the Cdc27 subunit of the cyclosome/APC. Cyclosomes from S-phase HeLa cells were partially purified and incubated with 500 units of Suc1-free Cdk1/cyclin B (Shteinberg, M. & Hershko, A., 1999, Biochem. Biophys. Res. Commun. 257:12; Yudkovsky, et al., 2000, Biochem. Biophys. Res. Commun. 271:299). Where indicated, 10 ng/µl of the corresponding Cks/Suc1 protein was supplemented. The samples were subjected to immunoblotting with a monoclonal antibody directed against human Cdc27 (Transduction Laboratories).

- FIG. 48 A, B. Identification of the factor required for p27-ubiquitin ligation as Cks1. **A.** The ligation of ^{35}S -p27 to MeUb was assayed. Where indicated, Fraction 1 (5 μg protein) or heat-treated Fraction 1 (~ 50 ng) were added. The bracket on the left side of the panels marks a ladder of bands $>27,000$ Da corresponding to polyubiquitinated p27. **B.**
- 5 Cks1, but not other Cks proteins, is required for p27-ubiquitin ligation. Where indicated, the following proteins were added: "Factor", 0.02 μl of pooled fractions # 28-29 from the peak of the Superdex column, which is the last step of purification of the factor required for p27 ubiquitinylation; "Cks1 IVT", 0.3 μl of in-vitro translated Cks1; "Cks2 IVT", 0.3 μl of in vitro-translated Cks2; "Retic. lys.", 0.3 μl of reticulocyte lysate translation mix; Cks1,
 - 10 Cks2 and Suc1, 2 ng of the corresponding bacterially expressed, purified proteins. *In vitro*-translated ^{35}S -labeled Cks1 and Cks2 in lanes 3 and 4 are not visible since they migrated off the gel.

- FIG 49 A-D. Cks1 increases the binding of phosphorylated p27 to Skp2. **A.**
- 15 Cks1 does not affect the phosphorylation of p27 by Cdk2/cyclin E. Purified p27 was phosphorylated with the only difference that the mixtures were incubated at 20°C for the time periods indicated. Where indicated, 2 ng of purified Cks1 was added. Samples of 1 μl were taken for SDS-polyacrylamide gel electrophoresis and autoradiography. **B.** Cks1 acts at a stage subsequent to the phosphorylation of p27. ^{32}P purified p27 was prepared. Where
 - 20 indicated, 0.02 μl of "Factor" (purified as in Fig. 1b, lane 2) or 1 ng of purified recombinant human Cks1 were added. Using this purified system, we have not observed conjugates with MeUb larger than the di-ubiquitinylated form, as opposed to the 4-5 conjugates observed using *in vitro*-translated ^{35}S -p27 (compare with Fig. 1). Possibly, ubiquitin is ligated to only
 - 25 (derived from ubiquitin present in reticulocyte lysates) terminated by MeUb. **C.** Cks1 increases the binding of p27 to Skp2/Skp1, dependent upon phosphorylation of Thr-187. The binding of ^{35}S -labeled wild-type (WT) or Thr-187-Ala mutant p27 (T187A) to Skp2/Skp1 was determined. Where indicated, 1 ng of purified Cks1 was added to the incubation. Inputs show 5% of the starting material. **D.** Cks1 increases the binding of ^{32}P -
 - 30 p27 to Skp2/Skp1. The experiment was similar to that described in 2c, except that ^{35}S -p27 was replaced by ^{32}P -labeled purified p27.

- Fig. 50 A-D. Binding of Cks1 to Skp2 and phosphorylated p27. **A.** Cks1 but not Cks2 binds to Skp2/Skp1. The binding of ^{35}S -labeled Cks1 or Cks2 to Skp2/Skp1
- 35 was assayed by a procedure similar to that described for the binding of p27 to Skp2/Skp1,

except that Cdk2/cyclin E, ATP and the ATP-regenerating system were omitted. Where indicated, 1 μ l of Skp2/Skp1 was added. **B.** Cks1 does not bind to Skp1. The binding of 35 S-Cks1 to His₆-Skp1 or to the Skp2/His₆-Skp1 complex (1 μ l each) was determined as described in 3a, except that Ni-NTA-agarose beads (Quiagen, 10 μ l) were used for precipitation. In both 3a and 3b, inputs show 5% of the starting material. **C.** Cks1 stimulates the binding of Skp2 to p27 phosphopeptide. Sepharose beads to which a peptide corresponding to 19 C-terminal amino acid residues of p27 ("p27 beads"), or to a similar peptide containing phosphorylated Thr187 ("P-p27 beads") were prepared as described in Carrano, et al., 1999, Nat. Cell Biol 1:193. In vitro-translated 35 S-Skp2 (3 μ l) was mixed with 15 μ l of the corresponding beads in the absence (lanes 1 and 3) or in the presence of 10 ng (lane 4) or 100 ng (lanes 2 and 5) of Cks1. Following rotation at 4°C for 2 hours, beads were washed 4 times with RIPA buffer. **D.** Cks1 binds to p27 phosphopeptide. 35 S-Cks1 (2 μ l) was mixed with the indicated beads, and beads were treated as in Fig. 3c. Inputs show 10% of the starting material.

FIG. 51 A-C. Western blot analysis of Skp2/E2F interaction assay. Details of the Western Blot experiments are given in the Example in Section 9.

5. DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to novel F-box proteins and to novel substrates of F-box proteins. The present invention relates to screening assays designed to identify substrates of the novel F-box proteins and to identify small molecules and compounds which modulate the interaction and/or activity of the F-box proteins and their substrates.

The present invention relates to screening assays to identify substrates of the novel F-box proteins and to identify potential therapeutic agents. The present invention further relates to screening assays based on the identification of novel substrates of both novel and known F-box proteins. The screening assays of the present invention may be used to identify potential therapeutic agents which may be used in protocols and as pharmaceutical compositions designed to target the novel ubiquitin ligases and interactions with their substrates for the treatment of proliferative disorders. In one particular embodiment the present invention relates to screening assays and potential therapeutic agents which target the interaction of FBP with novel substrates β -catenin, p27 and E2F as identified by Applicants.

The invention further encompasses the use of nucleotides encoding the novel F-box proteins, proteins and peptides, as well as antibodies to the novel ubiquitin ligases (which can, for example, act as agonists or antagonists), antagonists that inhibit ubiquitin ligase activity or expression, or agonists that activate ubiquitin ligase activity or increase its expression. In addition, nucleotides encoding the novel ubiquitin ligases and proteins are useful for the identification of compounds which regulate or mimic their activity and therefore are potentially effective in the treatment of cancer and tumorigenesis.

In particular, the invention described in the subsections below encompasses FBP1, FBP2, FBP3a, FBP3b, FBP4, FBP5, FBP6, FBP7, FBP8, FBP9, FBP10, FBP11, FBP12, FBP13, FBP14, FBP15, FBP16, FBP17, FBP18, FBP19, FBP20, FBP21, FBP22, FBP23, FBP24, and FBP25 polypeptides or peptides corresponding to functional domains of the novel ubiquitin ligases (*e.g.*, the F-box motif, the substrate binding domain, and leucine-rich repeats), mutated, truncated or deleted (*e.g.* with one or more functional domains or portions thereof deleted), ubiquitin ligase fusion proteins, nucleotide sequences encoding such products, and host cell expression systems that can produce such ubiquitin ligase products.

The present invention provides methods of screening for peptides and proteins that interact with novel components of the ubiquitin ligase complex, including FBP1, FBP2, FBP3a, FBP3b, FBP4, FBP5, FBP6, FBP7, FBP8, FBP9, FBP10, FBP11, FBP12, FBP13, FBP14, FBP15, FBP16, FBP17, FBP18, FBP19, FBP20, FBP21, FBP22, FBP23, FBP24, and FBP25 or derivatives, fragments or analogs thereof. Preferably, the method of screening is a yeast two-hybrid assay system or a variation thereof, as further described below. Derivatives (*e.g.*, fragments) and analogs of a protein can be assayed for binding to a binding partner by any method known in the art, for example, the modified yeast two-hybrid assay system described below, immunoprecipitation with an antibody that binds to the protein in a complex followed by analysis by size fractionation of the immunoprecipitated proteins (*e.g.*, by denaturing or nondenaturing polyacrylamide gel electrophoresis), Western analysis, non-denaturing gel electrophoresis, *etc.*

The present invention relates to screening assays to identify agents which modulate the activity of the novel ubiquitin ligases. The invention encompasses both *in vivo* and *in vitro* assays to screen small molecules, compounds, recombinant proteins, peptides, nucleic acids, antibodies *etc.* which modulate the activity of the novel ubiquitin ligases and thus, identify potential therapeutic agents for the treatment of proliferative or differentiative disorders. In one embodiment, the present invention provides methods of screening for proteins that interact with the novel ubiquitin ligases.

The invention also encompasses antibodies and anti-idiotypic antibodies, antagonists and agonists, as well as compounds or nucleotide constructs that inhibit expression of the ubiquitin ligase gene (transcription factor inhibitors, antisense and ribozyme molecules, or gene or regulatory sequence replacement constructs), or promote
5 expression of the ubiquitin ligase (*e.g.*, expression constructs in which ubiquitin ligase coding sequences are operatively associated with expression control elements such as promoters, promoter/enhancers, *etc.*). The invention also relates to host cells and animals genetically engineered to express the human (or mutants thereof) or to inhibit or "knock-out" expression of the animal's endogenous ubiquitin ligase.

10 Finally, the ubiquitin ligase protein products and fusion protein products, (*i.e.*, fusions of the proteins or a domain of the protein, *e.g.*, F-box motif), antibodies and anti-idiotypic antibodies (including Fab fragments), antagonists or agonists (including compounds that modulate the ubiquitization pathway can be used for therapy of proliferative or differentiative diseases. Thus, the invention also encompasses
15 pharmaceutical formulations and methods for treating cancer and tumorigenesis.

Various aspects of the invention are described in greater detail in the subsections below.

5.1 FBP GENES

20 The invention provides nucleic acid molecules comprising seven novel nucleotide sequences, and fragments thereof, FBP1, FBP2, FBP3a, FBP4, FBP5, FBP6, and FBP7, nucleic acids which are novel genes identified by the interaction of their gene products with Skp1, a component of the ubiquitin ligase complex. The invention further provides fourteen novel nucleic acid molecules comprising the nucleotide sequences of
25 FBP1, FBP2, FBP3a, FBP3b, FBP4, FBP5, FBP6, FBP7, FBP8, FBP11, FBP12, FBP13, FBP14, FBP15, FBP17, FBP18, FBP20, FBP21, FBP22, FBP23, FBP24, and FBP25, which Nucleic acid sequences of the identified FBP genes are described herein.

As used herein, "an FBP gene" refers to:

(a) a nucleic acid molecule containing the DNA sequences of FBP1, shown
30 in Figure 3 (SEQ ID NO:1), the DNA sequences of FBP2, shown in Figure 4 (SEQ ID NO:3), the DNA sequences of FBP3a, shown in Figure 5 (SEQ ID NO:5), the DNA sequences of FBP3b, shown in Figure 6 (SEQ ID NO:23), the DNA sequences of FBP4, shown in Figure 7 (SEQ ID NO:7), the DNA sequences of FBP5, shown in Figure 8 (SEQ ID NO:9), the DNA sequences of FBP6, shown in Figure 9 (SEQ ID NO:11), the DNA
35 sequences of FBP7, shown in Figure 10 (SEQ ID NO:13), the DNA sequences of FBP8,

- shown in Figure 11 (SEQ ID NO:25), the DNA sequences of FBP9, shown in Figure 12 (SEQ ID NO:27), the DNA sequences of FBP10, shown in Figure 13 (SEQ ID NO:29), the DNA sequences of FBP11, shown in Figure 14 (SEQ ID NO:31), the DNA sequences of FBP12, shown in Figure 15 (SEQ ID NO:33), the DNA sequences of FBP13, shown in
- 5 Figure 16 (SEQ ID NO:35), the DNA sequences of FBP14, shown in Figure 17 (SEQ ID NO:37), the DNA sequences of FBP15, shown in Figure 18 (SEQ ID NO:39), the DNA sequences of FBP16, shown in Figure 19 (SEQ ID NO:41), the DNA sequences of FBP17, shown in Figure 20 (SEQ ID NO:43), the DNA sequences of FBP18, shown in Figure 21 (SEQ ID NO:45), the DNA sequences of FBP19, shown in Figure 22 (SEQ ID NO:47), the
- 10 DNA sequences of FBP20, shown in Figure 23 (SEQ ID NO:49), the DNA sequences of FBP21, shown in Figure 24 (SEQ ID NO:51), the DNA sequences of FBP22, shown in Figure 25 (SEQ ID NO:53), the DNA sequences of FBP23, shown in Figure 26 (SEQ ID NO:55), the DNA sequences of FBP24, shown in Figure 27 (SEQ ID NO:57), the DNA sequences of FBP25, shown in Figure 28 (SEQ ID NO:59).
- 15 (b) any DNA sequence that encodes a polypeptide containing: the amino acid sequence of FBP1 shown in Figure 3A (SEQ ID NO:2), the amino acid sequence of FBP2, shown in Figure 4A (SEQ ID NO:4), the amino acid sequence of FBP3a shown in Figure 5A (SEQ ID NO:6), the amino acid sequence of FBP3b shown in Figure 6A (SEQ ID NO:24), the amino acid sequence of FBP4 shown in Figure 7A (SEQ ID NO:8), the amino
- 20 acid sequence of FBP5 shown in Figure 8A (SEQ ID NO:10), or the amino acid sequence of FBP6 shown in Figure 9A (SEQ ID NO:12), the amino acid sequences of FBP7, shown in Figure 10 (SEQ ID NO:14), the amino acid sequences of FBP8, shown in Figure 11 (SEQ ID NO:26), the amino acid sequences of FBP9, shown in Figure 12 (SEQ ID NO:28), the amino acid sequences of FBP10, shown in Figure 13 (SEQ ID NO:30), the amino acid
- 25 sequences of FBP11, shown in Figure 14 (SEQ ID NO:32), the amino acid sequences of FBP12, shown in Figure 15 (SEQ ID NO:34), the amino acid sequences of FBP13, shown in Figure 16 (SEQ ID NO:36), the amino acid sequences of FBP14, shown in Figure 17 (SEQ ID NO:38), the amino acid sequences of FBP15, shown in Figure 18 (SEQ ID NO:40), the amino acid sequences of FBP16, shown in Figure 19 (SEQ ID NO:42), the
- 30 amino acid sequences of FBP17, shown in Figure 20 (SEQ ID NO:44), the amino acid sequences of FBP18, shown in Figure 21 (SEQ ID NO:46), the amino acid sequences of FBP19, shown in Figure 22 (SEQ ID NO:48), the amino acid sequences of FBP20, shown in Figure 23 (SEQ ID NO:50), the amino acid sequences of FBP21, shown in Figure 24 (SEQ ID NO:52), the amino acid sequences of FBP22, shown in Figure 25 (SEQ ID
- 35 NO:54), the amino acid sequences of FBP23, shown in Figure 26 (SEQ ID NO:56), the

amino acid sequences of FBP24, shown in Figure 27 (SEQ ID NO:58), the amino acid sequences of FBP25, shown in Figure 28 (SEQ ID NO:60).

(c) any DNA sequence that hybridizes to the complement of the DNA sequences that encode any of the amino acid sequences of (SEQ ID NO: 2, 4, 6, 8, 10, 12 or 14) or Figure 15 under highly stringent conditions, *e.g.*, hybridization to filter-bound DNA in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65 C, and washing in 0.1xSSC/0.1% SDS at 68 C (Ausubel F.M. et al., eds., 1989, Current Protocols in Molecular Biology, Vol. I, Green Publishing Associates, Inc., and John Wiley & sons, Inc., New York, at p. 2.10.3); and/or

(d) any DNA sequence that hybridizes to the complement of the DNA sequences that encode any of the amino acid sequences in (SEQ ID NO: 2, 4, 6, 8, 10, 12 or 14) or Figure 15, under less stringent conditions, such as moderately stringent conditions, *e.g.*, washing in 0.2xSSC/0.1% SDS at 42 C (Ausubel et al., 1989, *supra*), and encodes a gene product functionally equivalent to an FBP gene product.

It is understood that the FBP gene sequences of the present invention do not encompass the previously described genes encoding other mammalian F-box proteins, Skp2, Elongin A, Cyclin F, mouse Md6, (see Pagano, 1997, *supra*; Zhang et al., 1995, *supra*; Bai et al., 1996, *supra*; Skowyra et al., 1997, *supra*). It is further understood that the nucleic acid molecules of the invention do not include nucleic acid molecules that consist solely of the nucleotide sequence in GenBank Accession Nos. AC002428, AI457595, AI105408, H66467, T47217, H38755, THC274684, AI750732, AA976979, AI571815, T57296, Z44228, Z45230, N42405, AA018063, AI751015, AI400663, T74432, AA402415, AI826000, AI590138, AF174602, Z45775, AF174599, THC288870, AI017603, AF174598, THC260994, AI475671, AA768343, AF174595, THC240016, N70417, T10511, AF174603, EST04915, AA147429, AI192344, AF174594, AI147207, AI279712, AA593015, AA644633, AA335703, N26196, AF174604, AF053356, AF174606, AA836036, AA853045, AI479142, AA772788, AA039454, AA397652, AA463756, AA007384, AA749085, AI640599, THC253263, AB020647, THC295423, AA434109, AA370939, AA215393, THC271423, AF052097, THC288182, AL049953, CAB37981, AL022395, AL031178, THC197682, and THC205131.

FBP sequences of the present invention are derived from a eukaryotic genome, preferably a mammalian genome, and more preferably a human or murine genome. Thus, the nucleotide sequences of the present invention do not encompass those derived from yeast genomes. In a specific embodiment, the nucleotides of the present invention encompass any DNA sequence derived from a mammalian genome which hybridizes under

highly stringent conditions to SEQ ID NO: 1, 3, 5, 7, 9, 11 or 13, or to DNA sequence shown in Figure 14, encodes a gene product which contains an F-box motif and binds to Skp1. In a specific embodiment, the nucleotides of the present invention encompass any DNA sequence derived from a mammalian genome which hybridize under highly stringent
5 conditions to SEQ ID NO: 1, 3, 5, 7, 9, 11 or 13 encodes a gene product which contains an F-box motif and another domain selected from the group comprising WD-40, leucine rich region, leucine zipper motif, or other protein-protein interaction domain, and binds to Skp-1 and is at least 300 or 400 nucleotides in length.

FBP sequences can include, for example, either eukaryotic genomic DNA
10 (cDNA) or cDNA sequences. When referring to a nucleic acid which encodes a given amino acid sequence, therefore, it is to be understood that the nucleic acid need not only be a cDNA molecule, but can also, for example, refer to a cDNA sequence from which an mRNA species is transcribed that is processed to encode the given amino acid sequence.

As used herein, an FBP gene may also refer to degenerate variants of DNA
15 sequences (a) through (d).

The invention also includes nucleic acid molecules derived from mammalian nucleic acids, preferably DNA molecules, that hybridize to, and are therefore the complements of, the DNA sequences (a) through (d), in the preceding paragraph. Such hybridization conditions may be highly stringent or less highly stringent, as described
20 above. In instances wherein the nucleic acid molecules are deoxyoligonucleotides ("oligos"), highly stringent conditions may refer, *e.g.*, to washing in 6xSSC/0.05% sodium pyrophosphate at 37 C (for 14-base oligos), 48 C (for 17-base oligos), 55 C (for 20-base oligos), and 60 C (for 23-base oligos). These nucleic acid molecules may encode or act as FBP gene antisense molecules, useful, for example, in FBP gene regulation (for and/or as
25 antisense primers in amplification reactions of FBP gene nucleic acid sequences). With respect to FBP gene regulation, such techniques can be used to regulate, for example, an FBP-regulated pathway, in order to block cell proliferation associated with cancer. Further, such sequences may be used as part of ribozyme and/or triple helix sequences, also useful for FBP gene regulation. Still further, such molecules may be used as components of
30 diagnostic methods whereby, for example, the presence of a particular FBP allele responsible for causing an FBP-related disorder, *e.g.*, proliferative or differentiative disorders such as tumorigenesis or cancer, may be detected.

The invention also encompasses:

(a) DNA vectors that contain any of the foregoing FBP coding sequences
35 and/or their complements (*i.e.*, antisense);

(b) DNA expression vectors that contain any of the foregoing FBP coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences; and

(c) genetically engineered host cells that contain any of the foregoing FBP coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences in the host cell.

As used herein, regulatory elements include but are not limited to inducible and non-inducible promoters, enhancers, operators and other elements known to those skilled in the art that drive and regulate expression. Such regulatory elements include but are not limited to the cytomegalovirus hCMV immediate early gene, the early or late promoters of SV40 adenovirus, the lac system, the trp system, the TAC system, the TRC system, the major operator and promoter regions of phage A, the control regions of fd coat protein, the promoter for 3-phosphoglycerate kinase, the promoters of acid phosphatase, and the promoters of the yeast-mating factors.

The invention further includes fragments of any of the DNA sequences disclosed herein.

In one embodiment, the FBP gene sequences of the invention are mammalian gene sequences, with human sequences being preferred.

In yet another embodiment, the FBP gene sequences of the invention are gene sequences encoding FBP gene products containing polypeptide portions corresponding to (that is, polypeptide portions exhibiting amino acid sequence similarity to) the amino acid sequence depicted in Figures 2, 4-9 or 15, wherein the corresponding portion exhibits greater than about 50% amino acid identity with the depicted sequence, averaged across the FBP gene product's entire length.

In specific embodiments, F-box encoding nucleic acids comprise the cDNA sequences of SEQ ID NOs: 1, 3, 5, 23, 7, 9, 11, 13, 15, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, or 59, nucleotide sequence of Figures 3B, 4B, 5B, 6B, 7B, 8B, 9B, 10B, 11B, 12B, 13B, 14B, 15B, 16B, 17B, 18B, 19B, 20B, 21B, 22B, 23B, 24B, 25B, 26B, 27B, or 28B, respectively, or the coding regions thereof, or nucleic acids encoding an F-box protein (e.g., a protein having the sequence of SEQ ID NOs: 2, 4, 6, 24, 8, 10, 12, 14, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 68, or 60, or as shown in Figures 3A, 4A, 5A, 6A, 7A, 8A, 9A, 10A, 11A, 12A, 13A, 14A, 15A, 16A, 17A, 18A, 19A, 20A, 21A, 22A, 23A, 24A, 25A, 26A, 27A, or 28A, respectively).

The invention further provides nucleotide fragments of nucleotide sequences encoding FBP1, FBP2, FBP3a, FBP4, FBP5, FBP6, or FBP7 (SEQ ID NOs: 1, 3, 5, 7, 9, 11

and 13, respectively) of the invention. Such fragments consist of at least 8 nucleotides (*i.e.*, a hybridizable portion) of an FBP gene sequence; in other embodiments, the nucleic acids consist of at least 25 (continuous) nucleotides, 50 nucleotides, 100 nucleotides, 150 nucleotides, or 200 nucleotides of an F-box sequence, or a full-length F-box coding
5 sequence. In another embodiment, the nucleic acids are smaller than 35, 200 or 500 nucleotides in length. Nucleic acids can be single or double stranded. The invention also relates to nucleic acids hybridizable to or complementary to the foregoing sequences. In specific aspects, nucleic acids are provided which comprise a sequence complementary to at least 10, 25, 50, 100, or 200 nucleotides or the entire coding region of an F-box gene.

10 The invention further relates to the human genomic nucleotide sequences of nucleic acids. In specific embodiments, F-box encoding nucleic acids comprise the genomic sequences of SEQ ID NOs: 1, 3, 5, 7, 9, 11 or 13 or the coding regions thereof, or nucleic acids encoding an FBP protein (*e.g.*, a protein having the sequence of SEQ ID Nos: 2, 4, 6, 8, 10, 12 or 14). The invention provides purified nucleic acids consisting of at least
15 8 nucleotides (*i.e.*, a hybridizable portion) of an FBP gene sequence; in other embodiments, the nucleic acids consist of at least 25 (continuous) nucleotides, 50 nucleotides, 100 nucleotides, 150 nucleotides, or 200 nucleotides of an FBP gene sequence or a full-length FBP gene coding sequence. In another embodiment, the nucleic acids are smaller than 35, 200 or 500 nucleotides in length. Nucleic acids can be single or double stranded. The
20 invention also relates to nucleic acids hybridizable to or complementary to the foregoing sequences. In specific aspects, nucleic acids are provided which comprise a sequence complementary to at least 10, 25, 50, 100, or 200 nucleotides or the entire coding region of an FBP gene sequence.

In addition to the human FBP nucleotide sequences disclosed herein, other
25 FBP gene sequences can be identified and readily isolated, without undue experimentation, by molecular biological techniques well known in the art, used in conjunction with the FBP gene sequences disclosed herein. For example, additional human FBP gene sequences at the same or at different genetic loci as those disclosed in SEQ ID Nos: 1, 3, 5, 7, 9, 11 or 13 can be isolated readily. There can exist, for example, genes at other genetic or physical loci
30 within the human genome that encode proteins that have extensive homology to one or more domains of the FBP gene products and that encode gene products functionally equivalent to an FBP gene product. Further, homologous FBP gene sequences present in other species can be identified and isolated readily.

The FBP nucleotide sequences of the invention further include nucleotide
35 sequences that encode polypeptides having at least 30%, 35%, 40%, 45%, 50%, 55%, 60%,

65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, or higher amino acid sequence identity to the polypeptides encoded by the FBP nucleotide sequences of SEQ ID No. 1, 3, 5, 7, 9, 11 or 13.

To determine the percent identity of two amino acid sequences or of two nucleic acids, the sequences are aligned for optimal comparison purposes (*e.g.*, gaps can be introduced in the sequence of a first amino acid or nucleic acid sequence for optimal alignment with a second amino or nucleic acid sequence). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences (*i.e.*, % identity = # of identical overlapping positions/total # of overlapping positions x 100%). In one embodiment, the two sequences are the same length.

The determination of percent identity between two sequences can also be accomplished using a mathematical algorithm. A preferred, non-limiting example of a mathematical algorithm utilized for the comparison of two sequences is the algorithm of Karlin and Altschul, 1990, Proc. Natl. Acad. Sci. USA 87:2264-2268, modified as in Karlin and Altschul, 1993, Proc. Natl. Acad. Sci. USA 90:5873-5877. Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul, et al., 1990, J. Mol. Biol. 215:403-410. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to a nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to a protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al., 1997, Nucleic Acids Res. 25:3389-3402. Alternatively, PSI-Blast can be used to perform an iterated search which detects distant relationships between molecules (Altschul et al., 1997, *supra*). When utilizing BLAST, Gapped BLAST, and PSI-Blast programs, the default parameters of the respective programs (*e.g.*, XBLAST and NBLAST) can be used (see <http://www.ncbi.nlm.nih.gov>). Another preferred, non-limiting example of a mathematical algorithm utilized for the comparison of two sequences is the algorithm of Karlin and Altschul, 1990, Proc. Natl. Acad. Sci. USA 87:2264-2268, modified as in Karlin and Altschul, 1993, Proc. Natl. Acad. Sci. USA 90:5873-5877. Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul, et al., 1990, J. Mol.

Biol. 215:403-410. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to a nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences
5 homologous to a protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al., 1997, Nucleic Acids Res. 25:3389-3402. Alternatively, PSI-Blast can be used to perform an iterated search which detects distant relationships between molecules (Altschul et al., 1997, *supra*). When utilizing BLAST, Gapped BLAST, and PSI-Blast programs, the default
10 parameters of the respective programs (*e.g.*, XBLAST and NBLAST) can be used (see <http://www.ncbi.nlm.nih.gov>). Another preferred, non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Myers and Miller, 1988, *CABIOS* 4:11-17. Such an algorithm is incorporated into the ALIGN program (version 2.0) which is part of the GCG sequence alignment software package. When
15 utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used.

The percent identity between two sequences can be determined using techniques similar to those described above, with or without allowing gaps. In calculating percent identity, typically only exact matches are counted.

20 With respect to identification and isolation of FBP gene sequences present at the same genetic or physical locus as those sequences disclosed herein, such sequences can, for example, be obtained readily by utilizing standard sequencing and bacterial artificial chromosome (BAC) technologies.

With respect to the cloning of an FBP gene homologue in human or other
25 species (*e.g.*, mouse), the isolated FBP gene sequences disclosed herein may be labeled and used to screen a cDNA library constructed from mRNA obtained from appropriate cells or tissues (*e.g.*, brain tissues) derived from the organism (*e.g.*, mouse) of interest. The hybridization conditions used should be of a lower stringency when the cDNA library is derived from an organism different from the type of organism from which the labeled
30 sequence was derived.

Alternatively, the labeled fragment may be used to screen a genomic library derived from the organism of interest, again, using appropriately stringent conditions. Low stringency conditions are well known to those of skill in the art, and will vary predictably depending on the specific organisms from which the library and the labeled sequences are
35 derived. For guidance regarding such conditions see, for example, Sambrook, et al., 1989,

Molecular Cloning, A Laboratory Manual, Second Edition, Cold Spring Harbor Press, N.Y.; and Ausubel, et al., *supra*. Further, an FBP gene homologue may be isolated from, for example, human nucleic acid, by performing PCR using two degenerate oligonucleotide primer pools designed on the basis of amino acid sequences within any FBP gene product disclosed herein.

The PCR product may be subcloned and sequenced to ensure that the amplified sequences represent the sequences of an FBP gene nucleic acid sequence. The PCR fragment may then be used to isolate a full length cDNA clone by a variety of methods. For example, the amplified fragment may be labeled and used to screen a bacteriophage cDNA library. Alternatively, the labeled fragment may be used to isolate genomic clones via the screening of a genomic library.

PCR technology may also be utilized to isolate full length cDNA sequences. For example, RNA may be isolated, following standard procedures, from an appropriate cellular or tissue source (*i.e.*, one known, or suspected, to express the FBP gene, such as, for example, blood samples or brain tissue samples obtained through biopsy or post-mortem). A reverse transcription reaction may be performed on the RNA using an oligonucleotide primer specific for the most 5' end of the amplified fragment for the priming of first strand synthesis. The resulting RNA/DNA hybrid may then be "tailed" with guanines using a standard terminal transferase reaction, the hybrid may be digested with RNAase H, and second strand synthesis may then be primed with a poly-C primer. Thus, cDNA sequences upstream of the amplified fragment may easily be isolated. For a review of cloning strategies that may be used, see *e.g.*, Sambrook et al., *supra*.

FBP gene sequences may additionally be used to identify mutant FBP gene alleles. Such mutant alleles may be isolated from individuals either known or proposed to have a genotype that contributes to the symptoms of an FBP gene disorder, such as proliferative or differentiative disorders involved in tumorigenesis or causing cancer, for example. Mutant alleles and mutant allele products may then be utilized in the therapeutic, diagnostic and prognostic systems described below. Additionally, such FBP gene sequences can be used to detect FBP gene regulatory (*e.g.*, promoter) defects which can be associated with an FBP disorder, such as proliferative or differentiative disorders involved in tumorigenesis or causing cancer, for example.

FBP alleles may be identified by single strand conformational polymorphism (SSCP) mutation detection techniques, Southern blot, and/or PCR amplification techniques. Primers can routinely be designed to amplify overlapping regions of the whole FBP sequence including the promoter region. In one embodiment, primers are designed to cover

the exon-intron boundaries such that, first, coding regions can be scanned for mutations. Genomic DNA isolated from lymphocytes of normal and affected individuals is used as PCR template. PCR products from normal and affected individuals are compared, either by single strand conformational polymorphism (SSCP) mutation detection techniques and/or by sequencing. SSCP analysis can be performed as follows: 100 ng of genomic DNA is amplified in a 10 μ l reaction, adding 10 pmols of each primer, 0.5 U of Taq DNA polymerase (Promega), 1 μ Ci of α -[32P]dCTP (NEN; specific activity, 3000 Ci/mmol), in 2.5 μ M dNTPs (Pharmacia), 10 mM Tris-HCl (pH 8.8), 50 mM KCl, 1 mM MgCl₂, 0.01% gelatin, final concentration. Thirty cycles of denaturation (94°C), annealing (56°C to 64°C, depending on primer melting temperature), and extension (72°C) is carried out in a thermal-cycler (MJ Research, Boston, MA, USA), followed by a 7 min final extension at 72°C. Two microliters of the reaction mixture is diluted in 0.1% SDS, 10 mM EDTA and then mixed 1: 1 with a sequencing stop solution containing 20 mM NaOH. Samples are heated at 95 C for 5 min, chilled on ice for 3 min and then 3 μ l will be loaded onto a 6% acrylamide/TBE gel containing 5% (v/v) glycerol. Gels are run at 8 W for 12-15 h at room temperature. Autoradiography is performed by exposure to film at -70 C with intensifying screens for different periods of time. The mutations responsible for the loss or alteration of function of the mutant FBP gene product can then be ascertained.

Alternatively, a cDNA of a mutant FBP gene may be isolated, for example, using PCR. In this case, the first cDNA strand may be synthesized by hybridizing an oligo-dT oligonucleotide to mRNA isolated from tissue known or suspected to be expressed in an individual putatively carrying the mutant FBP allele, and by extending the new strand with reverse transcriptase. The second strand of the cDNA is then synthesized using an oligonucleotide that hybridizes specifically to the 5' end of the normal gene. Using these two primers, the product is then amplified via PCR, cloned into a suitable vector, and subjected to DNA sequence analysis through methods well known to those of skill in the art. By comparing the DNA sequence of the mutant FBP allele to that of the normal FBP allele, the mutation(s) responsible for the loss or alteration of function of the mutant FBP gene product can be ascertained.

Alternatively, a genomic library can be constructed using DNA obtained from an individual suspected of or known to carry a mutant FBP allele, or a cDNA library can be constructed using RNA from a tissue known, or suspected, to express a mutant FBP allele. An unimpaired FBP gene or any suitable fragment thereof may then be labeled and used as a probe to identify the corresponding mutant FBP allele in such libraries. Clones

35

containing the mutant FBP gene sequences may then be purified and subjected to sequence analysis according to methods well known to those of skill in the art.

Additionally, an expression library can be constructed utilizing cDNA synthesized from, for example, RNA isolated from a tissue known, or suspected, to express a mutant FBP allele in an individual suspected of or known to carry such a mutant allele. In this manner, gene products made by the putatively mutant tissue may be expressed and screened using standard antibody screening techniques in conjunction with antibodies raised against the normal FBP gene product, as described, below, in Section 5.3. (For screening techniques, see, for example, Harlow and Lane, eds., 1988, "Antibodies: A Laboratory Manual", Cold Spring Harbor Press, Cold Spring Harbor.)

Nucleic acids encoding derivatives and analogs of FBP proteins, and FBP antisense nucleic acids can be isolated by the methods recited above. As used herein, a "nucleic acid encoding a fragment or portion of an F-box protein" shall be construed as referring to a nucleic acid encoding only the recited fragment or portion of the FBP and not the other contiguous portions of the FBP protein as a continuous sequence.

Fragments of FBP gene nucleic acids comprising regions conserved between (*i.e.*, with homology to) other FBP gene nucleic acids, of the same or different species, are also provided. Nucleic acids encoding one or more FBP domains can be isolated by the methods recited above.

In cases where an FBP mutation results in an expressed gene product with altered function (*e.g.*, as a result of a missense or a frameshift mutation), a polyclonal set of anti-FBP gene product antibodies are likely to cross-react with the mutant FBP gene product. Library clones detected via their reaction with such labeled antibodies can be purified and subjected to sequence analysis according to methods well known to those of skill in the art.

5.2 PROTEINS AND POLYPEPTIDES OF FBP GENES

The amino acid sequences depicted in Figures 1, 2, and parts B of Figures 3 to 28 represent FBP gene products. The FBP1 gene product, sometimes referred to herein as a "FBP1 protein", includes those gene products encoded by the FBP1 gene sequences described in Section 5.1, above. Likewise, the FBP2, FBP3a, FBP3b, FBP4, FBP5, FBP6, FBP7, FBP8, FBP9, FBP10, FBP11, FBP12, FBP13, FBP14, FBP15, FBP16, FBP17, FBP18, FBP19, FBP20, FBP21, FBP22, FBP23, FBP24, and FBP25 gene products, referred to herein as an FBP2, FBP3a, FBP3b, FBP4, FBP5, FBP6, FBP7, FBP8, FBP9, FBP10, FBP11, FBP12, FBP13, FBP14, FBP15, FBP16, FBP17, FBP18, FBP19, FBP20, FBP21,

FBP22, FBP23, FBP24, and FBP25 proteins, include those gene products encoded by the FBP2, FBP3, FBP4, FBP5, FBP6, FBP7, FBP8, FBP9, FBP10, FBP11, FBP12, FBP13, FBP14, FBP15, FBP16, FBP17, FBP18, FBP19, FBP20, FBP21, FBP22, FBP23, FBP24, and FBP25 genes. In accordance with the present invention, the nucleic acid sequences
5 encoding the FBP gene products are derived from eukaryotic genomes, including mammalian genomes. In a preferred embodiment the nucleic acid sequences encoding the FBP gene products are derived from human or murine genomes.

FBP gene products, or peptide fragments thereof, can be prepared for a variety of uses. For example, such gene products, or peptide fragments thereof, can be used
10 for the generation of antibodies, in diagnostic and prognostic assays, or for the identification of other cellular or extracellular gene products involved in the ubiquitination pathway and thereby implicated in the regulation of cell cycle and proliferative disorders.

In addition, FBP gene products of the present invention may include proteins that represent functionally equivalent (see Section 5.1 for a definition) gene products. FBP
15 gene products of the invention do not encompass the previously identified mammalian F-box proteins Skp2, Cyclin F, Elongin A, or mouse Md6 (see Pagano, 1997, *supra*; Zhang et al., 1995 *supra*; Bai et al., 1996 *supra*; Skowyra et al., 1997, *supra*).

Functionally equivalent FBP gene products may contain deletions, including internal deletions, additions, including additions yielding fusion proteins, or substitutions of
20 amino acid residues within and/or adjacent to the amino acid sequence encoded by the FBP gene sequences described, above, in Section 5.1, but that result in a "silent" change, in that the change produces a functionally equivalent FBP gene product. Amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues involved. For example,
25 nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine; polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine; positively charged (basic) amino acids include arginine, lysine, and histidine; and negatively charged (acidic) amino acids include aspartic acid and glutamic acid.

30 Alternatively, where alteration of function is desired, deletion or non-conservative alterations can be engineered to produce altered FBP gene products. Such alterations can, for example, alter one or more of the biological functions of the FBP gene product. Further, such alterations can be selected so as to generate FBP gene products that are better suited for expression, scale up, *etc.* in the host cells chosen. For example,

35

cysteine residues can be deleted or substituted with another amino acid residue in order to eliminate disulfide bridges.

The FBP gene products, peptide fragments thereof and fusion proteins thereof, may be produced by recombinant DNA technology using techniques well known in the art. Thus, methods for preparing the FBP gene polypeptides, peptides, fusion peptide and fusion polypeptides of the invention by expressing nucleic acid containing FBP gene sequences are described herein. Methods that are well known to those skilled in the art can be used to construct expression vectors containing FBP gene product coding sequences and appropriate transcriptional and translational control signals. These methods include, for example, in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. See, for example, the techniques described in Sambrook, et al., *supra*, and Ausubel, et al., *supra*. Alternatively, RNA capable of encoding FBP gene product sequences may be chemically synthesized using, for example, synthesizers. See, for example, the techniques described in "Oligonucleotide Synthesis", 1984, Gait, ed., IRL Press, Oxford.

A variety of host-expression vector systems may be utilized to express the FBP gene coding sequences of the invention. Such host-expression systems represent vehicles by which the coding sequences of interest may be produced and subsequently purified, but also represent cells that may, when transformed or transfected with the appropriate nucleotide coding sequences, exhibit the FBP gene product of the invention in situ. These include but are not limited to microorganisms such as bacteria (*e.g.*, *E. coli*, *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing FBP gene product coding sequences; yeast (*e.g.*, *Saccharomyces*, *Pichia*) transformed with recombinant yeast expression vectors containing the FBP gene product coding sequences; insect cell systems infected with recombinant virus expression vectors (*e.g.*, baculovirus) containing the FBP gene product coding sequences; plant cell systems infected with recombinant virus expression vectors (*e.g.*, cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (*e.g.*, Ti plasmid) containing FBP gene product coding sequences; or mammalian cell systems (*e.g.*, COS, CHO, BHK, 293, 3T3) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (*e.g.*, metallothionein promoter) or from mammalian viruses (*e.g.*, the adenovirus late promoter; the vaccinia virus 7.5K promoter).

In bacterial systems, a number of expression vectors may be advantageously selected depending upon the use intended for the FBP gene product being expressed. For

example, when a large quantity of such a protein is to be produced, for the generation of pharmaceutical compositions of FBP protein or for raising antibodies to FBP protein, for example, vectors that direct the expression of high levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not limited, to the E. coli
5 expression vector pUR278 (Ruther et al., 1983, EMBO J. 2, 1791), in which the FBP gene product coding sequence may be ligated individually into the vector in frame with the lac Z coding region so that a fusion protein is produced; pIN vectors (Inouye and Inouye, 1985, Nucleic Acids Res. 13, 3101-3109; Van Heeke and Schuster, 1989, J. Biol. Chem. 264, 5503-5509); and the like. pGEX vectors may also be used to express foreign polypeptides
10 as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. The pGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene product can be released from the GST moiety.

15 In an insect system, Autographa californica, nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in Spodoptera frugiperda cells. The FBP gene coding sequence may be cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter). Successful insertion of FBP
20 gene coding sequence will result in inactivation of the polyhedrin gene and production of non-occluded recombinant virus (*i.e.*, virus lacking the proteinaceous coat coded for by the polyhedrin gene). These recombinant viruses are then used to infect Spodoptera frugiperda cells in which the inserted gene is expressed (*e.g.*, see Smith et al., 1983, J. Virol. 46: 584; Smith, U.S. Patent No. 4,215,051).

25 In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, the FBP gene coding sequence of interest may be ligated to an adenovirus transcription/translation control complex, *e.g.*, the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by in vitro or in vivo recombination. Insertion in
30 a non-essential region of the viral genome (*e.g.*, region E1 or E3) will result in a recombinant virus that is viable and capable of expressing FBP gene product in infected hosts. (*e.g.*, See Logan and Shenk, 1984, Proc. Natl. Acad. Sci. USA 81, 3655-3659). Specific initiation signals may also be required for efficient translation of inserted FBP gene product coding sequences. These signals include the ATG initiation codon and adjacent
35 sequences. In cases where an entire FBP gene, including its own initiation codon and

adjacent sequences, is inserted into the appropriate expression vector, no additional translational control signals may be needed. However, in cases where only a portion of the FBP gene coding sequence is inserted, exogenous translational control signals, including, perhaps, the ATG initiation codon, must be provided. Furthermore, the initiation codon
5 must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, *etc.* (see Bittner, et al., 1987, Methods in Enzymol. 153, 516-544).

10 In addition, a host cell strain may be chosen that modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Such modifications (*e.g.*, glycosylation) and processing (*e.g.*, cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and
15 modification of proteins and gene products. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end, eukaryotic host cells that possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such mammalian host cells include but are not limited to CHO, VERO, BHK, HeLa,
20 COS, MDCK, 293, 3T3, and WI38.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines that stably express the FBP gene product may be engineered. Rather than using expression vectors that contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression
25 control elements (*e.g.*, promoter, enhancer, sequences, transcription terminators, polyadenylation sites, *etc.*), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid
30 into their chromosomes and grow to form foci that in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines that express the FBP gene product. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that affect the endogenous activity of the *FBP* gene product.

A number of selection systems may be used, including but not limited to the
35 herpes simplex virus thymidine kinase (Wigler, et al., 1977, Cell 11:223), hypoxanthine-

guanine phosphoribosyltransferase (Szybalska and Szybalski, 1962, Proc. Natl. Acad. Sci. USA 48, 2026), and adenine phosphoribosyltransferase (Lowy, et al., 1980, Cell 22, 817) genes can be employed in tk-, hgp^rt- or ap^rt- cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: dhfr, which confers resistance to methotrexate (Wigler, et al., 1980, Natl. Acad. Sci. USA 77, 3567; O'Hare, et al., 1981, Proc. Natl. Acad. Sci. USA 78, 1527); gpt, which confers resistance to mycophenolic acid (Mulligan and Berg, 1981, Proc. Natl. Acad. Sci. USA 78, 2072); neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin, et al., 1981, J. Mol. Biol. 150, 1); and hyg^r, which confers resistance to hygromycin (Santerre, et al., 1984, Gene 30, 147).

Alternatively, any fusion protein may be readily purified by utilizing an antibody specific for the fusion protein being expressed. For example, a system described by Janknecht, et al. allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht, et al., 1991, Proc. Natl. Acad. Sci. USA 88, 8972-8976). In this system, the gene of interest is subcloned into a vaccinia recombination plasmid such that the gene's open reading frame is translationally fused to an amino-terminal tag consisting of six histidine residues. Extracts from cells infected with recombinant vaccinia virus are loaded onto Ni²⁺ nitriloacetic acid-agarose columns and histidine-tagged proteins are selectively eluted with imidazole-containing buffers.

The FBP gene products can also be expressed in transgenic animals. Animals of any species, including, but not limited to, mice, rats, rabbits, guinea pigs, pigs, micro-pigs, goats, sheep, and non-human primates, *e.g.*, baboons, monkeys, and chimpanzees may be used to generate FBP transgenic animals. The term "transgenic," as used herein, refers to animals expressing FBP gene sequences from a different species (*e.g.*, mice expressing human FBP sequences), as well as animals that have been genetically engineered to overexpress endogenous (*i.e.*, same species) FBP sequences or animals that have been genetically engineered to no longer express endogenous FBP gene sequences (*i.e.*, "knock-out" animals), and their progeny.

In particular, the present invention relates to FBP1 knockout mice. The present invention also relates to transgenic mice which express human wild-type FBP1 and Skp2 gene sequences in addition to mice engineered to express human mutant FBP1 and Skp2 gene sequences deleted of their F-box domains. Any technique known in the art may be used to introduce an FBP gene transgene into animals to produce the founder lines of transgenic animals. Such techniques include, but are not limited to pronuclear microinjection (Hoppe and Wagner, 1989, U.S. Pat. No. 4,873,191); retrovirus mediated

gene transfer into germ lines (Van der Putten, et al., 1985, Proc. Natl. Acad. Sci., USA 82, 6148-6152); gene targeting in embryonic stem cells (Thompson, et al., 1989, Cell 56, 313-321); electroporation of embryos (Lo, 1983, Mol. Cell. Biol. 3, 1803-1814); and sperm-mediated gene transfer (Lavitrano et al., 1989, Cell 57, 717-723) (For a review of such techniques, see Gordon, 1989, Transgenic Animals, Intl. Rev. Cytol. 115, 171-229)

Any technique known in the art may be used to produce transgenic animal clones containing an FBP transgene, for example, nuclear transfer into enucleated oocytes of nuclei from cultured embryonic, fetal or adult cells induced to quiescence (Campbell, et al., 1996, Nature 380, 64-66; Wilmut, et al., Nature 385, 810-813).

The present invention provides for transgenic animals that carry an FBP transgene in all their cells, as well as animals that carry the transgene in some, but not all their cells, *i.e.*, mosaic animals. The transgene may be integrated as a single transgene or in concatamers, *e.g.*, head-to-head tandems or head-to-tail tandems. The transgene may also be selectively introduced into and activated in a particular cell type by following, for example, the teaching of Lasko et al. (Lasko, et al., 1992, Proc. Natl. Acad. Sci. USA 89, 6232-6236). The regulatory sequences required for such a cell-type specific activation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art. Examples of regulatory sequences that can be used to direct tissue-specific expression of an FBP transgene include, but are not limited to, the elastase I gene control region which is active in pancreatic acinar cells (Swift *et al.*, 1984, Cell 38:639-646; Ornitz *et al.*, 1986, Cold Spring Harbor Symp. Quant. Biol. 50:399-409; MacDonald, 1987, Hepatology 7:42S-51S); the insulin gene control region which is active in pancreatic beta cells (Hanahan, 1985, Nature 315:115-122); immunoglobulin gene control region which is active in lymphoid cells (Grosschedl *et al.*, 1984, Cell 38:647-658; Adams *et al.*, 1985, Nature 318:533-538; Alexander *et al.*, 1987, Mol. Cell. Biol. 7:1436-1444); albumin gene control region which is active in liver (Pinkert *et al.*, 1987, Genes and Devel. 1:268-276) alpha-fetoprotein gene control region which is active in liver (Krumlauf *et al.*, 1985, Mol. Cell. Biol. 5:1639-1648; Hammer *et al.*, 1987, Science 235:53-58); alpha-1-antitrypsin gene control region which is active in liver (Kelsey *et al.*, 1987, Genes and Devel. 1:161-171); beta-globin gene control region which is active in myeloid cells (Magram *et al.*, 1985, Nature 315:338-340; Kollias *et al.*, 1986, Cell 46:89-94); myelin basic protein gene control region which is active in oligodendrocyte cells in the brain (Readhead *et al.*, 1987, Cell 48:703-712); myosin light chain-2 gene control region which is active in skeletal muscle (Shani, 1985, Nature 314:283-286); and gonadotropic releasing hormone gene control region which is active in the hypothalamus (Mason *et al.*, 1986, Science 234:1372-1378).

Promoters isolated from the genome of viruses that grow in mammalian cells, (e.g., vaccinia virus 7.5K, SV40, HSV, adenoviruses MLP, MMTV, LTR and CMV promoters) may be used, as well as promoters produced by recombinant DNA or synthetic techniques.

When it is desired that the FBP gene transgene be integrated into the chromosomal site of the endogenous FBP gene, gene targeting is preferred. Briefly, when such a technique is to be utilized, vectors containing some nucleotide sequences homologous to the endogenous FBP gene are designed for the purpose of integrating, via homologous recombination with chromosomal sequences, into and disrupting the function of the nucleotide sequence of the endogenous FBP gene. The transgene may also be selectively introduced into a particular cell type, thus inactivating the endogenous FBP gene in only that cell type, by following, for example, the teaching of Gu, et al. (Gu, et al., 1994, Science 265, 103-106). The regulatory sequences required for such a cell-type specific inactivation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art.

Once transgenic animals have been generated, the expression of the recombinant FBP gene may be assayed utilizing standard techniques. Initial screening may be accomplished by Southern blot analysis or PCR techniques to analyze animal tissues to assay whether integration of the transgene has taken place. The level of mRNA expression of the transgene in the tissues of the transgenic animals may also be assessed using techniques that include but are not limited to Northern blot analysis of tissue samples obtained from the animal, in situ hybridization analysis, and RT-PCR (reverse transcriptase PCR). Samples of FBP gene-expressing tissue, may also be evaluated immunocytochemically using antibodies specific for the FBP transgene product.

Transgenic mice harboring tissue-directed transgenes can be used to test the effects of FBP gene expression the intact animal. In one embodiment, transgenic mice harboring a human FBP1 transgene in the mammary gland can be used to assess the role of FBPs in mouse mammary development and tumorigenesis. In another embodiment, transgenic mice can be generated that overexpress the human FBP1 dominant negative mutant form (F-box deleted) in the mammary gland. In a specific embodiment, for example, the MMTV LTR promoter (mouse mammary tumor virus long terminal repeat) can be used to direct integration of the transgene in the mammary gland. An MMTV/FBP1 fusion gene can be constructed by fusing sequences of the MMTV LTR promoter to nucleotide sequences upstream of the first ATG of FBP1 gene. An SV40 polyadenylation region can also be fused to sequences downstream of the FBP1 coding region. Transgenic mice are generated by methods well known in the art (Gordon, 1989, Transgenic Animals,

Intl. Rev. Cytol. 115, 171-229). Briefly, immature B6D2F1 female mice are superovulated and mated to CD-1 males. The following morning the females are examined for the presence of vaginal plugs, and fertilized ova are recovered and microinjected with a plasmid vector. Approximately 2000 copies of the material are microinjected into each pronucleus.

- 5 Screening of founder animals is performed by extraction of DNA from spleen and Southern hybridization using the MMTV/FBP1 as a probe. Screening of offspring is performed by PCR of tail DNA. Once transgenic pedigrees are established, the expression pattern of the transgene is determined by Northern blot and RT-PCR analysis in different organs in order to correlate it with subsequent pathological changes.

- 10 The resulting transgenic animals can then be examined for the role of FBP genes in tumorigenesis. In one embodiment, for example, FBP transgenes can be constructed for use as a breast cancer model. Overexpression of FBP1 genes in such mice is expected to increase β -catenin ubiquitination and degradation, resulting in a tumor suppressor phenotype. Conversely, overexpression of the FBP1 deletion mutant is expected
15 to result in stabilization of β -catenin and induce proliferation of mammary gland epithelium. These phenotypes can be tested in both female and male transgenic mice, by assays such as those described in Sections 5.4, 5.5 and 7.

- In another specific embodiment, transgenic mice are generated that express FBP1 transgenes in T-lymphocytes. In this embodiment, a CD2/FBP1 fusion gene is
20 constructed by fusion of the CD2 promoter, which drives expression in both CD4 positive and negative T-cells, to sequences located upstream of the first ATG of an FBP gene, *e.g.*, the wild-type and mutant FBP1 genes. The construct can also contain an SV40 polyadenylation region downstream of the FBP gene. After generation and testing of transgenic mice, as described above, the expression of the FBP transgene is examined. The
25 transgene is expressed in thymus and spleen. Overexpression of wild-type FBP1 is expected to result in a phenotype. For example, possible expected phenotypes of FBP1 transgenic mice include increased degradation of IKB α , increased activation of NF κ B, or increased cell proliferation. Conversely, overexpression of the dominant negative mutant, FBP1, lacking the F-box domain, can be expected to have the opposite effect, for example,
30 increased stability of IKB α , decreased activation of NF κ B, or decreased cell proliferation. Such transgenic phenotypes can be tested by assays such as those used in Section 5.4 and 5.5.

- In another specific embodiment, the SKP2 gene is expressed in T-lymphocytes of transgenic mice. Conversely, the F-box deletion form acts as dominant
35 negative, stabilizing p27 and inhibiting T-cell activation. Construction of the CD2/SKP2

fusion genes and production of transgenic mice are as described above for CD2/ FBP fusion genes, using wild-type and mutant SKP2 cDNA, instead of FBP1 cDNA, controlled by the CD2 promoter. Founders and their progeny are analyzed for the presence and expression of the SKP2 transgene and the mutant SKP2 transgene. Expression of the transgene in spleen and thymus is analyzed by Northern blot and RT-PCR

5 In another specific embodiment, transgenic mice are constructed by inactivation of the FBP1 locus in mice. Inactivation of the FBP1 locus in mice by homologous recombination involves four stages: 1) the construction of the targeting vector for FBP1; 2) the generation of ES +/- cells; 3) the production of knock-out mice; and 4) 10 the characterization of the phenotype. A 129 SV mouse genomic phage library is used to identify and isolate the mouse FBP1 gene. Bacteriophages are plated at an appropriate density and an imprint of the pattern of plaques can be obtained by gently layering a nylon membrane onto the surface of agarose dishes. Bacteriophage particles and DNA are transferred to the filter by capillary action in an exact replica of the pattern of plaques. 15 After denaturation, the DNA is bound to the filter by baking and then hybridized with ³²P-labeled-FBP1 cDNA. Excess probe is washed away and the filters were then exposed for autoradiography. Hybridizing plaques, identified by aligning the film with the original agar plate, were picked for a secondary and a tertiary screening to obtain a pure plaque preparation. Using this method, positive phage which span the region of interest, for 20 example, the region encoding the F-box, are isolated. Using PCR, Southern hybridization, restriction mapping, subcloning and DNA sequencing the partial structure of the wild-type FBP1 gene can be determined.

To inactivate the Fbp1 locus by homologous recombination, a gene targeting vector in which exon 3 in the Fbp1 locus is replaced by a selectable marker, for example, 25 the neoR gene, in an antisense orientation can be constructed. Exon 3 encodes the F-box motif which is known to be critical for Fbp1 interaction with Skp1. The targeting construct possesses a short and a long arm of homology flanking a selectable marker gene. One of the vector arms is relatively short (2 kb) to ensure efficient amplification since homologous recombinant ES clones will be screened by PCR. The other arm is >6 kb to maximize the 30 frequency of homologous recombination. A thymidine kinase (tk) gene, included at the end of the long homology arm of the vector provides an additional negative selection marker (using gancyclovir) against ES clones which randomly integrate the targeting vector. Since homologous recombination occurs frequently using linear DNA, the targeting vector is linearized prior to transfection of ES cells. Following electroporation and double drug 35 selection of embryonic stem cell clones, PCR and Southern analysis is used to determine

whether homologous recombination has occurred at the FBP1 locus. Screening by PCR is advantageous because a larger number of colonies can be analyzed with this method than with Southern analysis. In addition, PCR screening allows rapid elimination of negative clones thus to avoid feeding and subsequently freezing all the clones while recombinants are identified. This PCR strategy for detection of homologous recombinants is based on the use of a primer pair chosen such that one primer anneals to a sequence specific to the targeting construct, *e.g.*, sequences of the neomycin gene or other selectable marker, and not in the endogenous locus, and the other primer anneals to a region outside the construct, but within the endogenous locus. Southern analysis is used to confirm that a homologous recombination event has occurred (both at the short arm of homology and at the long arm of homology) and that no gene duplication events have occurred during the recombination.

Such FBP1 knockout mice can be used to test the role of Fbp1 in cellular regulation and control of proliferation. In one embodiment, phenotype of such mice lacking Fbp1 is cellular hyperplasia and increased tumor formation. In another embodiment, FBP1 null mice phenotypes include, but are not limited to, increased β -catenin activity, stabilization of β -catenin, increased cellular proliferation, accumulation of IK-Ba, decreased NF-KB activity, deficient immune response, inflammation, or increased cell death or apoptotic activity. Alternatively, a deletion of the of the FBP1 gene can result in an embryonic lethality. In this case, heterozygous mice at the FBP1 allele can be tested using the above assays, and embryos of null FBP mice can be tested using the assays described above.

Transgenic mice bearing FBP transgenes can also be used to screen for compounds capable of modulating the expression of the FBP gene and/or the synthesis or activity of the FBP1 gene or gene product. Such compounds and methods for screening are described.

5.3 GENERATION OF ANTIBODIES TO F-BOX PROTEINS AND THEIR DERIVATIVES

According to the invention, F-box motif, its fragments or other derivatives, or analogs thereof, may be used as an immunogen to generate antibodies which immunospecifically bind such an immunogen. Such antibodies include but are not limited to polyclonal, monoclonal, chimeric, single chain, Fab fragments, and an Fab expression library. In a specific embodiment, antibodies to a human FBP protein are produced. In

another embodiment, antibodies to a domain (*e.g.*, the F-box domain or the substrate-binding domain) of an FBP are produced.

Various procedures known in the art may be used for the production of polyclonal antibodies to an FBP or derivative or analog. In a particular embodiment, rabbit polyclonal antibodies to an epitope of an FBP encoded by a sequence of FBP1, FBP2, FBP3a, FBP3b, FBP4, FBP5, FBP6, FBP7, FBP8, FBP9, FBP10, FBP11, FBP12, FBP13, FBP14, FBP15, FBP16, FBP17, FBP18, FBP19, FBP20, FBP21, FBP22, FBP23, FBP24, and FBP25, or a subsequence thereof, can be obtained (Pagano, M., 1995, "From peptide to purified antibody", in *Cell Cycle: Materials and Methods*. M. Pagano, ed. Springer-Verlag. 217-281). For the production of antibody, various host animals can be immunized by injection with the native FBP, or a synthetic version, or derivative (*e.g.*, fragment) thereof, including but not limited to rabbits, mice, rats, *etc.* Various adjuvants may be used to increase the immunological response, depending on the host species, and including but not limited to Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanins, dinitrophenol, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and corynebacterium parvum.

For preparation of monoclonal antibodies directed toward an FBP sequence or analog thereof, any technique which provides for the production of antibody molecules by continuous cell lines in culture may be used. For example, the hybridoma technique originally developed by Kohler and Milstein (1975, *Nature* 256:495-497), as well as the trioma technique, the human B-cell hybridoma technique (Kozbor et al., 1983, *Immunology Today* 4:72), and the EBV-hybridoma technique to produce human monoclonal antibodies (Cole et al., 1985, in *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96). In an additional embodiment of the invention, monoclonal antibodies can be produced in germ-free animals utilizing recent technology (PCT/US90/02545). According to the invention, human antibodies may be used and can be obtained by using human hybridomas (Cote et al., 1983, *Proc. Natl. Acad. Sci. U.S.A.* 80:2026-2030) or by transforming human B cells with EBV virus in vitro (Cole et al., 1985, in *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, pp. 77-96). In fact, according to the invention, techniques developed for the production of "chimeric antibodies" (Morrison et al., 1984, *Proc. Natl. Acad. Sci. U.S.A.* 81:6851-6855; Neuberger et al., 1984, *Nature* 312:604-608; Takeda et al., 1985, *Nature* 314:452-454) by splicing the genes from a mouse antibody molecule specific for FBP together with genes from a human antibody molecule

35

of appropriate biological activity can be used; such antibodies are within the scope of this invention.

According to the invention, techniques described for the production of single chain antibodies (U.S. Patent No. 4,946,778) can be adapted to produce FBP-specific single chain antibodies. An additional embodiment of the invention utilizes the techniques described for the construction of Fab expression libraries (Huse et al., 1989, Science 246:1275-1281) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity for FBPs, derivatives, or analogs.

Antibody fragments which contain the idiotype of the molecule can be generated by known techniques. For example, such fragments include but are not limited to: the F(ab')₂ fragment which can be produced by pepsin digestion of the antibody molecule; the Fab' fragments which can be generated by reducing the disulfide bridges of the F(ab')₂ fragment, the Fab fragments which can be generated by treating the antibody molecule with papain and a reducing agent, and Fv fragments.

In the production of antibodies, screening for the desired antibody can be accomplished by techniques known in the art, e.g. ELISA (enzyme-linked immunosorbent assay). For example, to select antibodies which recognize a specific domain of an FBP, one may assay generated hybridomas for a product which binds to an FBP fragment containing such domain. For selection of an antibody that specifically binds a first FBP homolog but which does not specifically bind a different FBP homolog, one can select on the basis of positive binding to the first FBP homolog and a lack of binding to the second FBP homolog.

Antibodies specific to a domain of an FBP are also provided, such as an F-box motif.

The foregoing antibodies can be used in methods known in the art relating to the localization and activity of the FBP sequences of the invention, e.g., for imaging these proteins, measuring levels thereof in appropriate physiological samples, in diagnostic methods, etc.

In another embodiment of the invention (see *infra*), anti-FBP antibodies and fragments thereof containing the binding domain are used as therapeutics.

5.4 SCREENING ASSAYS FOR THE IDENTIFICATION OF AGENTS THAT INTERACT WITH F-BOX PROTEINS AND/OR INTERFERE WITH THEIR ENZYMATIC ACTIVITIES

Novel components of the ubiquitin ligase complex, including FBP1, FBP2, FBP3a, FBP3b, FBP4, FBP5, FBP6, FBP7, FBP8, FBP9, FBP10, FBP11, FBP12, FBP13, FBP14, FBP15, FBP16, FBP17, FBP18, FBP19, FBP20, FBP21, FBP22, FBP23, FBP24,

and FBP25, interact with cellular proteins to regulate cellular proliferation. One aspect of the present invention provides methods for assaying and screening fragments, derivatives and analogs of the novel components to identify polypeptides or peptides or other compounds that interact with the novel ubiquitin ligases such as potential substrates of ubiquitin ligase activity. The present invention also provides screening assays to identify compounds that modulate or inhibit the interaction of the novel FBPs with other subunits or numbers of the ubiquitin ligase complex, such as Skp1, or ubiquitinating enzymes with which the novel FBPs interact.

In yet another embodiment, the assays of the present invention may be used to identify polypeptides or peptides or other compounds which inhibit or modulate the interaction between the novel ubiquitin ligases or known (*e.g.*, Skp1) components of the ubiquitin ligase complex with novel or known substrates. By way of example, but not by limitation, the screening assays described herein may be used to identify peptides or proteins that interfere with the interaction between known ubiquitin ligase component, Skp2, and its novel substrate, p27. In another example, compounds that interfere with the interaction between FBP1 and its novel substrate, β -catenin, are identified using the screening assay. In another example, compounds that interfere with the interaction between Skp2 and another putative substrate, E2F, are identified using the screening assay. In yet another example, compounds that interfere with the interaction between FBP1 and another putative substrate, IKB α , are identified using the screening assay.

In yet another embodiment, the assays of the present invention may be used to identify polypeptides or peptides which inhibit or activate the enzymatic activators of the novel FBPs.

5.4.1 ASSAYS FOR PROTEIN-PROTEIN INTERACTIONS

Derivatives, analogs and fragments of proteins that interact with the novel components of the ubiquitin ligase complex of the present invention can be identified by means of a yeast two hybrid assay system (Fields and Song, 1989, Nature 340:245-246 and U.S. Patent No. 5,283,173). Because the interactions are screened for in yeast, the intermolecular protein interactions detected in this system occur under physiological conditions that mimic the conditions in mammalian cells (Chien et al., 1991, Proc. Natl. Acad. Sci. U.S.A. 88:9578-9581).

Identification of interacting proteins by the improved yeast two hybrid system is based upon the detection of expression of a reporter gene, the transcription of which is dependent upon the reconstitution of a transcriptional regulator by the interaction of two proteins, each fused to one half of the transcriptional regulator. The "bait" (*i.e.*, the novel components of the ubiquitin ligase complex of the present invention or derivatives or

analogous thereof) and "prey" (proteins to be tested for ability to interact with the bait) proteins are expressed as fusion proteins to a DNA binding domain, and to a transcriptional regulatory domain, respectively, or vice versa. In various specific embodiments, the prey has a complexity of at least about 50, about 100, about 500, about 1,000, about 5,000, about 10,000, or about 50,000; or has a complexity in the range of about 25 to about 100,000, about 100 to about 100,000, about 50,000 to about 100,000, or about 100,000 to about 500,000. For example, the prey population can be one or more nucleic acids encoding mutants of a protein (e.g., as generated by site-directed mutagenesis or another method of making mutations in a nucleotide sequence). Preferably, the prey populations are proteins encoded by DNA, e.g., cDNA or genomic DNA or synthetically-generated DNA. For example, the populations can be expressed from chimeric genes comprising cDNA sequences from an un-characterized sample of a population of cDNA from mRNA.

In a specific embodiment, recombinant biological libraries expressing random peptides can be used as the source of prey nucleic acids.

In general, proteins of the bait and prey populations are provided as fusion (chimeric) proteins (preferably by recombinant expression of a chimeric coding sequence) comprising each protein contiguous to a pre-selected sequence. For one population, the pre-selected sequence is a DNA binding domain. The DNA binding domain can be any DNA binding domain, as long as it specifically recognizes a DNA sequence within a promoter. For example, the DNA binding domain is of a transcriptional activator or inhibitor. For the other population, the pre-selected sequence is an activator or inhibitor domain of a transcriptional activator or inhibitor, respectively. The regulatory domain alone (not as a fusion to a protein sequence) and the DNA-binding domain alone (not as a fusion to a protein sequence) preferably do not detectably interact (so as to avoid false positives in the assay). The assay system further includes a reporter gene operably linked to a promoter that contains a binding site for the DNA binding domain of the transcriptional activator (or inhibitor). Accordingly, in the present method of the present invention, binding of a ubiquitin ligase fusion protein to a prey fusion protein leads to reconstitution of a transcriptional activator (or inhibitor) which activates (or inhibits) expression of the reporter gene. The activation (or inhibition) of transcription of the reporter gene occurs intracellularly, e.g., in prokaryotic or eukaryotic cells, preferably in cell culture.

The promoter that is operably linked to the reporter gene nucleotide sequence can be a native or non-native promoter of the nucleotide sequence, and the DNA binding site(s) that are recognized by the DNA binding domain portion of the fusion protein can be native to the promoter (if the promoter normally contains such binding site(s)) or non-native to the promoter.

Alternatively, the transcriptional activation binding site of the desired gene(s) can be deleted and replaced with GAL4 binding sites (Bartel et al., 1993, BioTechniques 14:920-924, Chasman et al., 1989, Mol. Cell. Biol. 9:4746-4749). The

reporter gene preferably contains the sequence encoding a detectable or selectable marker, the expression of which is regulated by the transcriptional activator, such that the marker is either turned on or off in the cell in response to the presence of a specific interaction. Preferably, the assay is carried out in the absence of background levels of the transcriptional activator (e.g., in a cell that is mutant or otherwise lacking in the transcriptional activator).

5 The activation domain and DNA binding domain used in the assay can be from a wide variety of transcriptional activator proteins, as long as these transcriptional activators have separable binding and transcriptional activation domains. For example, the GAL4 protein of *S. cerevisiae* (Ma et al., 1987, Cell 48:847-853), the GCN4 protein of *S. cerevisiae* (Hope & Struhl, 1986, Cell 46:885-894), the ARD1 protein of *S. cerevisiae* (Thukral et al., 1989, Mol. Cell. Biol. 9:2360-2369), and the human estrogen receptor (Kumar et al., 1987, Cell 51:941-951), have separable DNA binding and activation domains. The DNA binding domain and activation domain that are employed in the fusion proteins need not be from the same transcriptional activator. In a specific embodiment, a GAL4 or LEXA DNA binding domain is employed. In another specific embodiment, a GAL4 or herpes simplex virus VP16 (Triezenberg et al., 1988, Genes Dev. 2:730-742) activation domain is employed. In a specific embodiment, amino acids 1-147 of GAL4 (Ma et al., 1987, Cell 48:847-853; Ptashne et al., 1990, Nature 346:329-331) is the DNA binding domain, and amino acids 411-455 of VP16 (Triezenberg et al., 1988, Genes Dev. 2:730-742; Cress et al., 1991, Science 251:87-90) comprise the activation domain.

In a preferred embodiment, the yeast transcription factor GAL4 is reconstituted by protein-protein interaction and the host strain is mutant for GAL4. In another embodiment, the DNA-binding domain is Ace1N and/or the activation domain is Ace1, the DNA binding and activation domains of the Ace1 protein, respectively. Ace1 is a yeast protein that activates transcription from the CUP1 operon in the presence of divalent copper. CUP1 encodes metallothionein, which chelates copper, and the expression of CUP1 protein allows growth in the presence of copper, which is otherwise toxic to the host cells. The reporter gene can also be a CUP1-lacZ fusion that expresses the enzyme beta-galactosidase (detectable by routine chromogenic assay) upon binding of a reconstituted Ace1N transcriptional activator (see Chaudhuri et al., 1995, FEBS Letters 357:221-226). In another specific embodiment, the DNA binding domain of the human estrogen receptor is used, with a reporter gene driven by one or three estrogen receptor response elements (Le Douarin et al., 1995, Nucl. Acids. Res. 23:876-878). The DNA binding domain and the transcriptional activator/inhibitor domain each preferably has a nuclear localization signal (see Ylikomi et al., 1992, EMBO J. 11:3681-3694, Dingwall and Laskey, 1991, TIBS 16:479-481) functional in the cell in which the fusion proteins are to be expressed.

To facilitate isolation of the encoded proteins, the fusion constructs can further contain sequences encoding affinity tags such as glutathione-S-transferase or maltose-binding protein or an epitope of an available antibody, for affinity purification (e.g.,

binding to glutathione, maltose, or a particular antibody specific for the epitope, respectively) (Allen et al., 1995, TIBS 20:511-516). In another embodiment, the fusion constructs further comprise bacterial promoter sequences for recombinant production of the fusion protein in bacterial cells.

5 The host cell in which the interaction assay occurs can be any cell, prokaryotic or eukaryotic, in which transcription of the reporter gene can occur and be detected, including, but not limited to, mammalian (*e.g.*, monkey, mouse, rat, human, bovine), chicken, bacterial, or insect cells, and is preferably a yeast cell. Expression constructs encoding and capable of expressing the binding domain fusion proteins, the transcriptional activation domain fusion proteins, and the reporter gene product(s) are
10 provided within the host cell, by mating of cells containing the expression constructs, or by cell fusion, transformation, electroporation, microinjection, *etc.*

Various vectors and host strains for expression of the two fusion protein populations in yeast are known and can be used (see *e.g.*, U.S. Patent No. 5,1468,614; Bartel et al., 1993, "Using the two-hybrid system to detect protein-protein interactions" In:
15 Cellular Interactions in Development, Hartley, ed., Practical Approach Series xviii, IRL Press at Oxford University Press, New York, NY, pp. 153-179; Fields and Sternglanz, 1994, Trends In Genetics 10:286-292).

If not already lacking in endogenous reporter gene activity, cells mutant in the reporter gene may be selected by known methods, or the cells can be made mutant in the target reporter gene by known gene-disruption methods prior to introducing the reporter
20 gene (Rothstein, 1983, Meth. Enzymol. 101:202-211).

In a specific embodiment, plasmids encoding the different fusion protein populations can be introduced simultaneously into a single host cell (*e.g.*, a haploid yeast cell) containing one or more reporter genes, by co-transformation, to conduct the assay for protein-protein interactions. Or, preferably, the two fusion protein populations are
25 introduced into a single cell either by mating (*e.g.*, for yeast cells) or cell fusions (*e.g.*, of mammalian cells). In a mating type assay, conjugation of haploid yeast cells of opposite mating type that have been transformed with a binding domain fusion expression construct (preferably a plasmid) and an activation (or inhibitor) domain fusion expression construct (preferably a plasmid), respectively, will deliver both constructs into the same diploid cell. The mating type of a yeast strain may be manipulated by transformation with the HO gene
30 (Herskowitz and Jensen, 1991, Meth. Enzymol. 194:132-146).

In a preferred embodiment, a yeast interaction mating assay is employed using two different types of host cells, strain-type α and a of the yeast *Saccharomyces cerevisiae*. The host cell preferably contains at least two reporter genes, each with one or more binding sites for the DNA-binding domain (*e.g.*, of a transcriptional activator). The activator domain and DNA binding domain are each parts of chimeric proteins formed from
35 the two respective populations of proteins. One strain of host cells, for example the a strain,

contains fusions of the library of nucleotide sequences with the DNA-binding domain of a transcriptional activator, such as GAL4. The hybrid proteins expressed in this set of host cells are capable of recognizing the DNA-binding site in the promoter or enhancer region in the reporter gene construct. The second set of yeast host cells, for example, the alpha strain, contains nucleotide sequences encoding fusions of a library of DNA sequences fused to the activation domain of a transcriptional activator.

In another embodiment, the fusion constructs are introduced directly into the yeast chromosome via homologous recombination. The homologous recombination for these purposes is mediated through yeast sequences that are not essential for vegetative growth of yeast, *e.g.*, the MER2, MER1, ZIPI, REC102, or ME14 gene.

Bacteriophage vectors can also be used to express the DNA binding domain and/or activation domain fusion proteins. Libraries can generally be prepared faster and more easily from bacteriophage vectors than from plasmid vectors.

In a specific embodiment, the present invention provides a method of detecting one or more protein-protein interactions comprising (a) recombinantly expressing a novel ubiquitin ligase component of the present invention or a derivative or analog thereof in a first population of yeast cells being of a first mating type and comprising a first fusion protein containing the sequence of a novel ubiquitin ligase component of the present invention and a DNA binding domain, wherein said first population of yeast cells contains a first nucleotide sequence operably linked to a promoter driven by one or more DNA binding sites recognized by said DNA binding domain such that an interaction of said first fusion protein with a second fusion protein, said second fusion protein comprising a transcriptional activation domain, results in increased transcription of said first nucleotide sequence; (b) negatively selecting to eliminate those yeast cells in said first population in which said increased transcription of said first nucleotide sequence occurs in the absence of said second fusion protein; (c) recombinantly expressing in a second population of yeast cells of a second mating type different from said first mating type, a plurality of said second fusion proteins, each second fusion protein comprising a sequence of a fragment, derivative or analog of a protein and an activation domain of a transcriptional activator, in which the activation domain is the same in each said second fusion protein; (d) mating said first population of yeast cells with said second population of yeast cells to form a third population of diploid yeast cells, wherein said third population of diploid yeast cells contains a second nucleotide sequence operably linked to a promoter driven by a DNA binding site recognized by said DNA binding domain such that an interaction of a first fusion protein with a second fusion protein results in increased transcription of said second nucleotide sequence, in which the first and second nucleotide sequences can be the same or different; and (e) detecting said increased transcription of said first and/or second nucleotide sequence, thereby detecting an interaction between a first fusion protein and a second fusion protein.

5.4.2 ASSAYS TO IDENTIFY F-BOX PROTEIN INTERACTIONS WITH KNOWN PROTEINS INCLUDING POTENTIAL SUBSTRATES

The cellular abundance of cell-cycle regulatory proteins, such as members of the cyclin family or the Cki inhibitory proteins, is regulated by the ubiquitin pathway. The enzymes responsible for the ubiquitination of mammalian cell cycle regulation are not known. In yeast, SCF complexes represent the ubiquitin ligases for cell cycle regulators. The F-box component of the ubiquitin ligase complexes, such as the novel F-box proteins of the invention, determines the specificity of the target of the ubiquitin ligase complex. The invention therefore provides assays to screen known molecules for specific binding to F-box protein nucleic acids, proteins, or derivatives under conditions conducive to binding, and then molecules that specifically bind to the FBP protein are identified.

In a specific embodiment, the invention provides a method for studying the interaction between the F-box protein FBP1 and the Cul1/Skp1 complex, and its role in regulating the stability of β -catenin. Protein-protein interactions can be probed in vivo and in vitro using antibodies specific to these proteins, as described in detail in the experiments in Section 7.

In another specific embodiment, methods for detecting the interaction between Skp2 and p27, a cell cycle regulated cyclin-dependent kinase (Cdk) inhibitor, are provided, as described in Section 8. The interaction between Skp2 and p27 may be targeted to identify modulators of Skp2 activity, including its interaction with cell cycle regulators, such as p27. The ubiquitination of Skp2-specific substrates, such as p27 may be used as a means of measuring the ability of a test compound to modulate Skp2 activity. In another embodiment of the screening assays of the present invention, immunodepletion assays, as described in Section 8, can be used to identify modulators of the Skp2/p27 interaction. In particular, Section 8 describes a method for detection of ubiquitination activity in vitro using p27 as a substrate, which can also be used to identify modulators of the Skp2-dependent ubiquitination of p27. In another embodiment of the screening assays of the present invention, antisense oligonucleotides, as described in Section 5.7.1, can be used as inhibitors of the Skp2 activity. Such identified modulators of p27 ubiquitination/degradation and of the Skp2/p27 interaction can be useful in anti-cancer therapies.

In another specific embodiment, methods for detecting the interaction between Skp2 and Cks1 and Skp2, Cks1, and p27 are provided. The interaction between Skp2 and Cks1, and Skp2, Cks1 and p27 may be targeted to identify modulators of Skp2 activity, including its interaction with molecules involved in the cell cycle, such as Cks1

and p27. The ubiquitination of Skp2-specific substrates, such as p27 may be used as a means of measuring the ability of a test compound to modulate Skp2 activity in the presence or absence of Cks1. Section 9 describes another embodiment of the screening assays of the present invention for detection of ubiquitination activity by Skp2 with or without Cks1 in vitro using p27 or a phospho-peptide corresponding to the carboxy terminus of p27 with or without a phosphothreonine at position 187 as a substrate, which can also be used to identify modulators of the Skp2-dependent ubiquitination of p27. In another embodiment of the screening assays of the present invention, antisense oligonucleotides, as described in Section 5.7.1, can be used as inhibitors of the Skp2 activity. Such identified modulators of p27 ubiquitination/degradation and of the Skp2/Cks1/p27 interaction can be useful in anti-cancer therapies.

In another specific embodiment, the invention provides for a method for detecting the interaction between the F-box protein Skp2 and E2F-1, a transcription factor involved in cell cycle progression. Insect cells can be infected with baculoviruses co-expressing Skp2 and E2F-1, and cell extracts can be prepared and analyzed for protein-protein interactions. As described in detail in Section 10, this assay has been used successfully to identify potential targets, such as E2F, for known F-box proteins, such as Skp2. This assay can be used to identify other Skp2 targets, as well as targets for novel F-box proteins.

The invention further provides methods for screening ubiquitin ligase complexes having novel F-box proteins (or fragments thereof) as one of their components for ubiquitin ligase activity using known cell-cycle regulatory molecules as potential substrates for ubiquitination. For example, cells engineered to express FBP nucleic acids can be used to recombinantly produce FBP proteins either wild-type or dominant negative mutants in cells that also express a putative ubiquitin-ligase substrate molecule. Such candidates for substrates of the novel FBP of the present invention include, but are not limited to, such potential substrates as IKB α , β -catenin, myc, E2F-1, p27, p21, cyclin A, cyclin B, cycD1, cyclin E and p53. Then the extracts can be used to test the association of F-box proteins with their substrates, (by Western blot immunoassays) and whether the presence of the FBP increases or decreases the level of the potential substrates.

5.5 ASSAYS FOR THE IDENTIFICATION OF COMPOUNDS THAT MODULATE THE ACTIVITY OF F-BOX PROTEINS

The present invention relates to in vitro and in vivo assay systems described in the subsections below, which can be used to identify compounds or compositions that

modulate the interaction of known FBPs with novel substrates and novel components of the ubiquitin ligase complex. The screening assays of the present invention may also be used to identify compounds or compositions that modulate the interaction of novel FBPs with their identified substrates and components of the ubiquitin ligase complex.

5 Methods to screen potential agents for their ability to disrupt or moderate FBP expression and activity can be designed based on the Applicants' discovery of novel FBPs and their interaction with other components of the ubiquitin ligase complex as well as its known and potential substrates. For example, candidate compounds can be screened for their ability to modulate the interaction of an FBP and Skp1, or the specific interactions of
10 Skp2 with E2F-1, Skp2 with Cks1, Skp2 with Cks1 and p27, or the FBP1/Cul1/Skp1 complex with β -catenin. In principle, many methods known to those of skill in the art, can be readily adapted in designed the assays of the present invention.

 The screening assays of the present invention also encompass high-throughput screens and assays to identify modulators of FBP expression and activity. In
15 accordance with this embodiment, the systems described below may be formulated into kits. To this end, cells expressing FBP and components of the ubiquitination ligase complex and the ubiquitination pathway, or cell lysates, thereof can be packaged in a variety of containers, *e.g.*, vials, tubes, microtitre well plates, bottles, and the like. Other reagents can be included in separate containers and provided with the kit; *e.g.*, positive control samples,
20 negative control samples, buffers, cell culture media, *etc.*

 The invention provides screening methodologies useful in the identification of proteins and other compounds which bind to, or otherwise directly interact with, the FBP genes and their gene products. Screening methodologies are well known in the art (see *e.g.*, PCT International Publication No. WO 96/34099, published October 31, 1996, which is
25 incorporated by reference herein in its entirety). The proteins and compounds include endogenous cellular components which interact with the identified genes and proteins in vivo and which, therefore, may provide new targets for pharmaceutical and therapeutic interventions, as well as recombinant, synthetic, and otherwise exogenous compounds which may have binding capacity and, therefore, may be candidates for pharmaceutical
30 agents. Thus, in one series of embodiments, cell lysates or tissue homogenates may be screened for proteins or other compounds which bind to one of the normal or mutant FBP genes and FBP proteins.

 Alternatively, any of a variety of exogenous compounds, both naturally occurring and/or synthetic (*e.g.*, libraries of small molecules or peptides), may be screened
35 for binding capacity. All of these methods comprise the step of mixing an FBP protein or

fragment with test compounds, allowing time for any binding to occur, and assaying for any bound complexes. All such methods are enabled by the present disclosure of substantially pure FBP proteins, substantially pure functional domain fragments, fusion proteins, antibodies, and methods of making and using the same.

5

5.5.1 ASSAYS FOR F-BOX PROTEIN AGONISTS AND ANTAGONISTS

FBP nucleic acids, F-box proteins, and derivatives can be used in screening assays to detect molecules that specifically bind to FBP nucleic acids, proteins, or derivatives and thus have potential use as agonists or antagonists of FBPs, in particular, molecules that thus affect cell proliferation. In a preferred embodiment, such assays are performed to screen for molecules with potential utility as anti-cancer drugs or lead compounds for drug development. The invention thus provides assays to detect molecules that specifically bind to FBP nucleic acids, proteins, or derivatives. For example, recombinant cells expressing FBP nucleic acids can be used to recombinantly produce FBP proteins in these assays, to screen for molecules that bind to an FBP protein. Similar methods can be used to screen for molecules that bind to FBP derivatives or nucleic acids. Methods that can be used to carry out the foregoing are commonly known in the art. The assays of the present invention may be first optimized on a small scale (*i.e.*, in test tubes), and then scaled up for high-throughput assays. The screening assays of the present may be performed *in vitro*, *i.e.* in test tubes, using purified components or cell lysates. The screening assays of the present invention may also be carried out in intact cells in culture and in animal models. In accordance with the present invention, test compounds which are shown to modulate the activity of the FBP as described herein *in vitro*, will further be assayed *in vivo*, including cultured cells and animal models to determine if the test compound has the similar effects *in vivo* and to determine the effects of the test compound on cell cycle progression, the accumulation or degradation of positive and negative regulators, cellular proliferation *etc.*

In accordance with the present invention, screening assays may be designed to detect molecules which act as agonists or antagonists of the activity of the novel F-box proteins. In accordance with this aspect of the invention, the test compound may be added to an assay system to measure its effect on the activity of the novel FBP, *i.e.*, ubiquitination of its substrates, interaction with other components of the ubiquitin ligase complex, *etc.* These assays should be conducted both in the presence and absence of the test compound.

In accordance with the present invention, ubiquitination activity of a novel FBP in the presence or absence of a test compound can be measured *in vitro* using purified

components of the ubiquitination pathway or may be measured using crude cellular extracts obtained from tissue culture cells or tissue samples. In another embodiment of the aspect of the present invention the screening may be performed by adding the test agent to in vitro translation systems such as a rabbit reticulocyte lysate (RRL) system and then proceeding with the established analysis. As another alternative, purified or partially purified components which have been determined to interact with one another by the methods described above can be placed under conditions in which the interaction between them would normally occur, with and without the addition of the test agent, and the procedures previously established to analyze the interaction can be used to assess the impact of the test agent. In this approach, the purified or partially purified components may be prepared by fractionation of extracts of cells expressing the components of the ubiquitin ligase complex and pathway, or they may be obtained by expression of cloned genes or cDNAs or fragments thereof, optionally followed by purification of the expressed material.

Within the broad category of in vitro selection methods, several types of method are likely to be particularly convenient and/or useful for screening test agents. These include but are not limited to methods which measure a binding interaction between two or more components of the ubiquitin ligase complex or interaction with the target substrate, methods which measure the activity of an enzyme which is one of the interacting components, and methods which measure the activity or expression of "reporter" protein, that is, an enzyme or other detectable or selectable protein, which has been placed under the control of one of the components.

Binding interactions between two or more components can be measured in a variety of ways. One approach is to label one of the components with an easily detectable label, place it together with the other component(s) in conditions under which they would normally interact, perform a separation step which separates bound labeled component from unbound labeled component, and then measure the amount of bound component. The effect of a test agent included in the binding reaction can be determined by comparing the amount of labeled component which binds in the presence of this agent to the amount which binds in its absence.

In another embodiment, screening can be carried out by contacting the library members with an FBP protein (or nucleic acid or derivative) immobilized on a solid phase and harvesting those library members that bind to the protein (or nucleic acid or derivative). Examples of such screening methods, termed "panning" techniques are described by way of example in Parmley & Smith, 1988, Gene 73:305-318; Fowlkes et al., 1992, BioTechniques 13:422-427; PCT Publication No. WO 94/18318; and in references cited hereinabove.

In another embodiment, the two-hybrid system for selecting interacting proteins or peptides in yeast (Fields & Song, 1989, Nature 340:245-246; Chien et al., 1991, Proc. Natl. Acad. Sci. USA 88:9578-9582) can be used to identify molecules that specifically bind to an FBP protein or derivative.

5 Alternatively, test methods may rely on measurements of enzyme activity, such as ubiquitination of the target substrate. Once a substrate of a novel FBP is identified or a novel putative substrate of a known FBP is identified, such as the novel substrates of Skp2, E2F and p27, these components may be used in assays to determine the effect of a test compound on the ubiquitin ligase activity of the ubiquitin ligase complex.

10 In one embodiment, the screening assays may be conducted with a purified system in the presence and absence of test compound. Purified substrate is incubated together with purified ubiquitin ligase complex, ubiquitin conjugating enzymes, ubiquitin activating enzymes and ubiquitin in the presence or in the absence of test compound. Ubiquitination of the substrate is analyzed by immunoassay (see Pagano et al., 1995, 15 Science 269:682-685). Briefly, ubiquitination of the substrate can be performed in vitro in reactions containing 50-200ng of proteins in 50mM Tris pH 7.5, 5mM MgCl₂, 2mM ATPγ-S, 0.1 mM DTT and 5μM of biotinylated ubiquitin. Total reactions (30μl) can be incubated at 25°C for up to 3 hours in the presence or absence of test compound and then loaded on an 8% SDS gel or a 4-20% gradient gel for analysis. The gels are run and 20 proteins are electrophoretically transferred to nitrocellulose. Ubiquitination of the substrate can be detected by immunoblotting. Ubiquitinated substrates can be visualized using Extravidin-HRP (Sigma), or by using a substrate-specific antibody, and the ECL detection system (NEN).

In another embodiment, ubiquitination of the substrate may be assayed in 25 intact cells in culture or in animal models in the presence and absence of the test compound. For example, the test compound may be administered directly to an animal model or to crude extracts obtained from animal tissue samples to measure ubiquitination of the substrate in the presence and absence of the test compounds. For these assays, host cells to which the test compound is added may be genetically engineered to express the FBP 30 components of the ubiquitin ligase pathway and the target substrate, the expression of which may be transient, induced or constitutive, or stable. For the purposes of the screening methods of the present invention, a wide variety of host cells may be used including, but not limited to, tissue culture cells, mammalian cells, yeast cells, and bacteria. Each cell type has its own set of advantages and drawbacks. Mammalian cells such as primary cultures of 35 human tissue cells may be a preferred cell type in which to carry out the assays of the

present invention, however these cell types are sometimes difficult to cultivate. Bacteria and yeast are relatively easy to cultivate but process proteins differently than mammalian cells. This ubiquitination assay may be conducted as follows: first, the extracts are prepared from human or animal tissue. To prepare animal tissue samples preserving ubiquitinating enzymes, 1 g of tissue can be sectioned and homogenized at 15,000 r.p.m. with a
5 Brinkmann Polytron homogenizer (PT 3000, Westbury, NY) in 1 ml of ice-cold double-distilled water. The sample is frozen and thawed 3 times. The lysate is spun down at 15,000 r.p.m. in a Beckman JA-20.1 rotor (Beckman Instruments, Palo Alto, CA) for 45 min at 4°C. The supernatant is retrieved and frozen at -80°C. This method of preparation of
10 total extract preserves ubiquitinating enzymes (Loda et al. 1997, Nature Medicine 3:231-234, incorporated by reference herein in its entirety).

Purified recombinant substrate is added to the assay system and incubated at 37°C for different times in 30 µl of ubiquitination mix containing 100 µg of protein tissue homogenates, 50 mM Tris-HCl (pH 8.0), 5 mM MgCl₂, and 1 mM DTT, 2 mM ATP, 10
15 mM creatine phosphokinase, 10 mM creatine phosphate and 5 µM biotinylated ubiquitin. The substrate is then re-purified with antibodies or affinity chromatography. Ubiquitination of the substrate is measured by immunoassays with either antibodies specific to the substrates or with Extravidin-HRP.

In addition, *Drosophila* can be used as a model system in order to detect
20 genes that phenotypically interact with FBP. For example, overexpression of FBP in *Drosophila* eye leads to a smaller and rougher eye. Mutagenesis of the fly genome can be performed, followed by selecting flies in which the mutagenesis has resulted in suppression or enhancement of the small rough eye phenotype; the mutated genes in such flies are likely to encode proteins that interact/bind with FBP. Active compounds identified with methods
25 described above will be tested in cultured cells and/or animal models to test the effect of blocking in vivo FBP activity (*e.g.* effects on cell proliferation, accumulation of substrates, *etc.*).

In various other embodiments, screening the can be accomplished by one of many commonly known methods. See, *e.g.*, the following references, which disclose
30 screening of peptide libraries: Parmley & Smith, 1989, Adv. Exp. Med. Biol. 251:215-218; Scott & Smith, 1990, Science 249:386-390; Fowlkes et al., 1992; BioTechniques 13:422-427; Oldenburg et al., 1992, Proc. Natl. Acad. Sci. USA 89:5393-5397; Yu et al., 1994, Cell 76:933-945; Staudt et al., 1988, Science 241:577-580; Bock et al., 1992, Nature 355:564-566; Tuerk et al., 1992, Proc. Natl. Acad. Sci. USA 89:6988-6992; Ellington et al., 1992,
35 Nature 355:850-852; U.S. Patent No. 5,096,815, U.S. Patent No. 5,223,409, and U.S.

Patent No. 5,198,346, all to Ladner et al.; Rebar & Pabo, 1993, Science 263:671-673; and PCT Publication No. WO 94/18318.

Compounds, peptides, and small molecules can be used in screening assays to identify candidate agonists and antagonists. In one embodiment, peptide libraries may be
5 used to screen for agonists or antagonists of the FBP of the present invention diversity libraries, such as random or combinatorial peptide or non-peptide libraries can be screened for molecules that specifically bind to FBP. Many libraries are known in the art that can be used, *e.g.*, chemically synthesized libraries, recombinant (*e.g.*, phage display libraries), and in vitro translation-based libraries.

10 Examples of chemically synthesized libraries are described in Fodor et al., 1991, Science 251:767-773; Houghten et al., 1991, Nature 354:84-86; Lam et al., 1991, Nature 354:82-84; Medynski, 1994, Bio/Technology 12:709-710; Gallop et al., 1994, J. Medicinal Chemistry 37(9):1233-1251; Ohlmeyer et al., 1993, Proc. Natl. Acad. Sci. USA 90:10922-10926; Erb et al., 1994, Proc. Natl. Acad. Sci. USA 91:11422-11426; Houghten
15 et al., 1992, Biotechniques 13:412; Jayawickreme et al., 1994, Proc. Natl. Acad. Sci. USA 91:1614-1618; Salmon et al., 1993, Proc. Natl. Acad. Sci. USA 90:11708-11712; PCT Publication No. WO 93/20242; and Brenner and Lerner, 1992, Proc. Natl. Acad. Sci. USA 89:5381-5383.

Examples of phage display libraries are described in Scott & Smith, 1990,
20 Science 249:386-390; Devlin et al., 1990, Science, 249:404-406; Christian, et al., 1992, J. Mol. Biol. 227:711-718; Lenstra, 1992, J. Immunol. Meth. 152:149-157; Kay et al., 1993, Gene 128:59-65; and PCT Publication No. WO 94/18318 dated August 18, 1994.

In vitro translation-based libraries include but are not limited to those described in PCT Publication No. WO 91/05058 dated April 18, 1991; and Mattheakis et
25 al., 1994, Proc. Natl. Acad. Sci. USA 91:9022-9026.

By way of examples of non-peptide libraries, a benzodiazepine library (see *e.g.*, Bunin et al., 1994, Proc. Natl. Acad. Sci. USA 91:4708-4712) can be adapted for use. Peptoid libraries (Simon et al., 1992, Proc. Natl. Acad. Sci. USA 89:9367-9371) can also be used. Another example of a library that can be used, in which the amide functionalities in
30 peptides have been permethylated to generate a chemically transformed combinatorial library, is described by Ostresh et al. (1994, Proc. Natl. Acad. Sci. USA 91:11138-11142).

5.5.2 ASSAYS FOR THE IDENTIFICATION OF COMPOUNDS THAT MODULATE THE INTERACTION OF F-box PROTEINS WITH OTHER PROTEINS

Once a substrate or interacting protein is identified, as described in detail in Section 5.4, then one can assay for modulators of the F-box protein interaction with such a protein. The present invention provides for methods of detecting agonists and antagonists of such interactions.

In one embodiment, the invention encompasses methods to identify modulators, such as inhibitors or agonists, of the interaction between the F-box protein Skp2 and E2F-1, identified in Section 7 and Figure 10. Such methods comprise both in vivo and in vitro assays for modulator activity. For example, in an in vivo assay, insect cells can be co-infected with baculoviruses co-expressing Skp2 and E2F-1 as well as potential modulators of the Skp2/E2F-1 interaction. The screening methods of the present invention encompass in vitro assays which measure the ability of a test compound to inhibit the enzymatic activity of Skp2 as described above in Section 5.5.1. Cell extracts can be prepared and analyzed for protein-protein interactions by gel electrophoresis and detected by immunoblotting, as described in detail in Section 7 and presented in Figure 10.

Alternatively, an in vitro protein-protein interaction assay can be used. Recombinant purified Skp2, E2F-1, and putative agonist or antagonist molecules can be incubated together, under conditions that allow binding to occur, such as 37 C for 30 minutes. Protein-protein complex formation can be detected by gel analysis, such as those described herein in Section 7. This assay can be used to identify modulators of interactions of known FBP, such as Skp2 with novel substrates.

In another embodiment, the invention provides for a method for identification of modulators of F-box protein/Skp1 interaction. Such agonist and antagonists can be identified in vivo or in vitro. For example, in an in vitro assay to identify modulators of F-box protein/Skp1 interactions, purified Skp1 and the novel FBP can be incubated together, under conditions that allow binding occur, such as 37C for 30 minutes. In a parallel reaction, a potential agonist or antagonist, as described above in Section 5.5.1, is added either before or during the box protein/Skp1 incubation. Protein-protein interactions can be detected by gel analysis, such as those described herein in Section 7. Modulators of FBP activities and interactions with other proteins can be used as therapeutics using the methods described herein, in Section 5.7.

These assays may be carried out utilizing any of the screening methods described herein, including the following in vitro assay. The screening can be performed by adding the test agent to intact cells which express components of the ubiquitin pathway, and

then examining the component of interest by whatever procedure has been established. Alternatively, the screening can be performed by adding the test agent to in vitro translation reactions and then proceeding with the established analysis. As another alternative, purified or partially purified components which have been determined to interact with one another
5 by the methods described above can be placed under conditions in which the interaction between them would normally occur, with and without the addition of the test agent, and the procedures previously established to analyze the interaction can be used to assess the impact of the test agent. In this approach, the purified or partially purified components may be prepared by fractionation of extracts of cells expressing the components of the ubiquitin
10 ligase complex and pathway, or they may be obtained by expression of cloned genes or cDNAs or fragments thereof, optionally followed by purification of the expressed material.

Within the broad category of in vitro selection methods, several types of method are likely to be particularly convenient and/or useful for screening test agents. These include but are not limited to methods which measure a binding interaction between
15 two or more components of the ubiquitin ligase complex or interaction with the target substrate, methods which measure the activity of an enzyme which is one of the interacting components, and methods which measure the activity or expression of "reporter" protein, that is, an enzyme or other detectable or selectable protein, which has been placed under the control of one of the components.

20 Binding interactions between two or more components can be measured in a variety of ways. One approach is to label one of the components with an easily detectable label, place it together with the other component(s) in conditions under which they would normally interact, perform a separation step which separates bound labeled component from unbound labeled component, and then measure the amount of bound component. The effect
25 of a test agent included in the binding reaction can be determined by comparing the amount of labeled component which binds in the presence of this agent to the amount which binds in its absence.

The separation step in this type of procedure can be accomplished in various ways. In one approach, (one of) the binding partner(s) for the labeled component can be
30 immobilized on a solid phase prior to the binding reaction, and unbound labeled component can be removed after the binding reaction by washing the solid phase. Attachment of the binding partner to the solid phase can be accomplished in various ways known to those skilled in the art, including but not limited to chemical cross-linking, non-specific adhesion to a plastic surface, interaction with an antibody attached to the solid phase, interaction

35

between a ligand attached to the binding partner (such as biotin) and a ligand-binding protein (such as avidin or streptavidin) attached to the solid phase, and so on.

Alternatively, the separation step can be accomplished after the labeled component had been allowed to interact with its binding partner(s) in solution. If the size differences between the labeled component and its binding partner(s) permit such a separation, the separation can be achieved by passing the products of the binding reaction through an ultrafilter whose pores allow passage of unbound labeled component but not of its binding partner(s) or of labeled component bound to its partner(s). Separation can also be achieved using any reagent capable of capturing a binding partner of the labeled component from solution, such as an antibody against the binding partner, a ligand-binding protein which can interact with a ligand previously attached to the binding partner, and so on.

5.6 METHODS AND COMPOSITIONS FOR DIAGNOSTIC USE OF F-BOX PROTEINS, DERIVATIVES, AND MODULATORS

Cell cycle regulators are the products of oncogenes (cyclins, β -catenin, *etc.*), or tumor suppressor genes (ckis, p53, *etc.*) The FBPs, part of ubiquitin ligase complexes, might therefore be products of oncogenes or tumor suppressor genes, depending on which cell cycle regulatory proteins for which they regulate cellular abundance.

FBP proteins, analogues, derivatives, and subsequences thereof, FBP nucleic acids (and sequences complementary thereto), anti-FBP antibodies, have uses in diagnostics. The FBP and FBP nucleic acids can be used in assays to detect, prognose, or diagnose proliferative or differentiative disorders, including tumorigenesis, carcinomas, adenomas *etc.* The novel FBP nucleic acids of the present invention are located at chromosome sites associated with karyotypic abnormalities and loss of heterozygosity. The FBP1 nucleic acid of the present invention is mapped and localized to chromosome position 10q24, the loss of which has been demonstrated in 10 % of human prostate tumors and small cell lung carcinomas (SCLC), suggesting the presence of a tumor suppressor gene at this location. In addition, up to 7% of childhood acute T-cell leukemia is accompanied by a translocation involving 10q24 as a breakpoint, either t(10;14)(q24;q11) or t(7;10)(q35;q24). 9q34 region (where FBP2 is located) has been shown to be a site of loss of heterozygosity (LOH) in human ovarian and bladder cancers. The FBP2 nucleic acid of the present invention is mapped and localized to chromosome position 9q34 which has been shown to be a site of loss of heterozygosity (LOH) in human ovarian and bladder cancers. The FBP3 nucleic acid of the present invention is mapped and localized to chromosome position

13q22, a region known to contain a putative tumor suppressor gene with loss of heterozygosity in approx. 75 % of human SCLC. The FBP4 nucleic acid of the present invention is mapped and localized to chromosome position 5p12, a region shown to be a site of karyotypic abnormalities in a variety of tumors, including human breast cancer and nasopharyngeal carcinomas. The FBP5 nucleic acid of the present invention is mapped and localized to chromosome position 6q25-26, a region shown to be a site of loss of heterozygosity in human ovarian, breast and gastric cancers hepatocarcinomas, Burkitt's lymphomas, gliomas, and parathyroid adenomas. The FBP7 nucleic acid of the present invention is mapped and localized to chromosome position 15q15 a region which contains a tumor suppressor gene associated with progression to a metastatic stage in breast and colon cancers and a loss of heterozygosity in parathyroid adenomas.

The molecules of the present invention can be used in assays, such as immunoassays, to detect, prognose, diagnose, or monitor various conditions, diseases, and disorders affecting FBP expression, or monitor the treatment thereof. In particular, such an immunoassay is carried out by a method comprising contacting a sample derived from a patient with an anti-FBP antibody under conditions such that immunospecific binding can occur, and detecting or measuring the amount of any immunospecific binding by the antibody. In a specific aspect, such binding of antibody, in tissue sections, can be used to detect aberrant FBP localization or aberrant (*e.g.*, low or absent) levels of FBP. In a specific embodiment, antibody to FBP can be used to assay a patient tissue or serum sample for the presence of FBP where an aberrant level of FBP is an indication of a diseased condition. By "aberrant levels," is meant increased or decreased levels relative to that present, or a standard level representing that present, in an analogous sample from a portion of the body or from a subject not having the disorder.

The immunoassays which can be used include but are not limited to competitive and non-competitive assay systems using techniques such as western blots, immunohisto-chemistry radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoprecipitation assays, precipitin reactions, gel diffusion precipitin reactions, immunodiffusion assays, agglutination assays, complement-fixation assays, immunoradiometric assays, fluorescent immunoassays, protein A immunoassays, to name but a few.

FBP genes and related nucleic acid sequences and subsequences, including complementary sequences, can also be used in hybridization assays. FBP nucleic acid sequences, or subsequences thereof comprising about at least 8 nucleotides, can be used as hybridization probes. Hybridization assays can be used to detect, prognose, diagnose, or

monitor conditions, disorders, or disease states associated with aberrant changes in FBP expression and/or activity as described supra. In particular, such a hybridization assay is carried out by a method comprising contacting a sample containing nucleic acid with a nucleic acid probe capable of hybridizing to FBP DNA or RNA, under conditions such that

5 hybridization can occur, and detecting or measuring any resulting hybridization.

In specific embodiments, diseases and disorders involving overproliferation of cells can be diagnosed, or their suspected presence can be screened for, or a predisposition to develop such disorders can be detected, by detecting decreased levels of FBP protein, FBP RNA, or FBP functional activity (*e.g.*, ubiquitin ligase target binding

10 activity, F-box domain binding activity, ubiquitin ligase activity *etc.*), or by detecting mutations in FBP RNA, DNA or FBP protein (*e.g.*, translocations in FBP nucleic acids, truncations in the FBP gene or protein, changes in nucleotide or amino acid sequence relative to wild-type FBP) that cause decreased expression or activity of FBP. Such diseases and disorders include but are not limited to those described in Section 5.7.3. By

15 way of example, levels of FBP protein can be detected by immunoassay, levels of FBP RNA can be detected by hybridization assays (*e.g.*, Northern blots, in situ-hybridization), FBP activity can be assayed by measuring ubiquitin ligase activity in E3 ubiquitin ligase complexes formed in vivo or in vitro, F-box domain binding activity can be assayed by measuring binding to Skp1 protein by binding assays commonly known in the art,

20 translocations, deletions and point mutations in FBP nucleic acids can be detected by Southern blotting, FISH, RFLP analysis, SSCP, PCR using primers that preferably generate a fragment spanning at least most of the FBP gene, sequencing of FBP genomic DNA or cDNA obtained from the patient, *etc.*

In a preferred embodiment, levels of FBP mRNA or protein in a patient

25 sample are detected or measured, in which decreased levels indicate that the subject has, or has a predisposition to developing, a malignancy or hyperproliferative disorder; in which the decreased levels are relative to the levels present in an analogous sample from a portion of the body or from a subject not having the malignancy or hyperproliferative disorder, as the case may be.

30 In another specific embodiment, diseases and disorders involving a deficiency in cell proliferation or in which cell proliferation is desirable for treatment, are diagnosed, or their suspected presence can be screened for, or a predisposition to develop such disorders can be detected, by detecting increased levels of FBP protein, FBP RNA, or FBP functional activity (*e.g.*, ubiquitin ligase activity, Skp1 binding activity, *etc.*), or by

35 detecting mutations in FBP RNA, DNA or protein (*e.g.*, translocations in FBP nucleic acids,

truncations in the gene or protein, changes in nucleotide or amino acid sequence relative to wild-type FBP) that cause increased expression or activity of FBP. Such diseases and disorders include but are not limited to those described in Section 5.7.3. By way of example, levels of FBP protein, levels of FBP RNA, ubiquitin ligase activity, FBP binding activity, and the presence of translocations or point mutations can be determined as described above.

In a specific embodiment, levels of FBP mRNA or protein in a patient sample are detected or measured, in which increased levels indicate that the subject has, or has a predisposition to developing, a growth deficiency or degenerative or hypoproliferative disorder; in which the increased levels are relative to the levels present in an analogous sample from a portion of the body or from a subject not having the growth deficiency, degenerative, or hypoproliferative disorder, as the case may be.

Kits for diagnostic use are also provided, that comprise in one or more containers an anti-FBP antibody, and, optionally, a labeled binding partner to the antibody. Alternatively, the anti-FBP antibody can be labeled (with a detectable marker, *e.g.*, a chemiluminescent, enzymatic, fluorescent, or radioactive moiety). A kit is also provided that comprises in one or more containers a nucleic acid probe capable of hybridizing to FBP RNA. In a specific embodiment, a kit can comprise in one or more containers a pair of primers (*e.g.*, each in the size range of 6-30 nucleotides) that are capable of priming amplification [*e.g.*, by polymerase chain reaction (see *e.g.*, Innis et al., 1990, PCR Protocols, Academic Press, Inc., San Diego, CA), ligase chain reaction (see EP 320,308) use of Q replicase, cyclic probe reaction, or other methods known in the art] under appropriate reaction conditions of at least a portion of a FBP nucleic acid. A kit can optionally further comprise in a container a predetermined amount of a purified FBP protein or nucleic acid, *e.g.*, for use as a standard or control.

5.7 METHODS AND COMPOSITIONS FOR THERAPEUTIC USE OF F-box PROTEINS, DERIVATIVES, AND MODULATORS

Described below are methods and compositions for the use of F-box proteins in the treatment of proliferative disorders and oncogenic disease symptoms may be ameliorated by compounds that activate or enhance FBP activity, and whereby proliferative disorders and cancer may be ameliorated.

In certain instances, compounds and methods that increase or enhance the activity of an FBP can be used to treat proliferative and oncogenic disease symptoms. Such a case may involve, for example, a proliferative disorder that is brought about, at least in

part, by a reduced level of FBP gene expression, or an aberrant level of an FBP gene product's activity. For example, decreased activity or under-expression of an FBP component of a ubiquitin ligase complex whose substrate is a positive cell-cycle regulator, such as a member of the Cyclin family, will result in increased cell proliferation. As such, an increase in the level of gene expression and/or the activity of such FBP gene products would bring about the amelioration of proliferative disease symptoms.

In another instance, compounds that increase or enhance the activity of an FBP can be used to treat proliferative and oncogenic disease symptoms resulting from defects in the expression or activity of other genes and gene products involved in cell cycle control, such as FBP substrate molecules. For example, an increase in the expression or activity of a positive cell-cycle positive molecule, such as a member of the Cyclin family, may result in its over-activity and thereby lead to increased cell proliferation. Compounds that increase the expression or activity of the FBP component of a ubiquitin ligase complex whose substrate is such a cell-cycle positive regulator will lead to ubiquitination of the defective molecule, and thereby result in an increase in its degradation. Disease symptoms resulting from such a defect may be ameliorated by compounds that compensate the disorder by increased FBP activity. Techniques for increasing FBP gene expression levels or gene product activity levels are discussed in Section 5.7, below.

Alternatively, compounds and methods that reduce or inactivate FBP activity may be used therapeutically to ameliorate proliferative and oncogenic disease symptoms. For example, a proliferative disorder may be caused, at least in part, by a defective FBP gene or gene product that leads to its overactivity. Where such a defective gene product is a component of a ubiquitin ligase complex whose target is a cell-cycle inhibitor molecule, such as a Cki, an overactive FBP will lead to a decrease in the level of cell-cycle molecule and therefore an increase in cell proliferation. In such an instance, compounds and methods that reduce or inactivate FBP function may be used to treat the disease symptoms.

In another instance, compounds and methods that reduce the activity of an FBP can be used to treat disorders resulting from defects in the expression or activity of other genes and gene products involved in cell cycle control, such as FBP substrate molecules. For example, a defect in the expression or activity of a cell-cycle negative regulatory molecule, such as a Cki, may lead to its under-activity and thereby result in increased cell proliferation. Reduction in the level and/or activity of an FBP component whose substrate was such molecule would decrease the ubiquitination and thereby increase the level of such a defective molecule. Therefore, compounds and methods aimed at

35

reducing the expression and/or activity of such FBP molecules could thereby be used in the treatment of disease symptoms by compensating for the defective gene or gene product.

Techniques for the reduction of target gene expression levels or target gene product activity levels are discussed in Section 5.7 below.

5

5.7.1 THERAPEUTIC USE OF INHIBITORY ANTISENSE, RIBOZYME AND TRIPLE HELIX MOLECULES AND IDENTIFIED AGONISTS AND ANTAGONISTS

In another embodiment, symptoms of certain FBP disorders, such as such as
10 proliferative or differentiative disorders causing tumorigenesis or cancer, may be ameliorated by decreasing the level of FBP gene expression and/or FBP gene product activity by using FBP gene sequences in conjunction with well-known antisense, gene "knock-out" ribozyme and/or triple helix methods to decrease the level of FBP gene expression. Among the compounds that may exhibit the ability to modulate the activity,
15 expression or synthesis of the FBP gene, including the ability to ameliorate the symptoms of an FBP disorder, such as cancer, are antisense, ribozyme, and triple helix molecules. Such molecules may be designed to reduce or inhibit either unimpaired, or if appropriate, mutant target gene activity. Techniques for the production and use of such molecules are well known to those of skill in the art. For example, antisense targeting SKP2 mRNA stabilize
20 the Skp2-substrate p27, as described in Section X (Figure X).

Antisense RNA and DNA molecules act to directly block the translation of mRNA by hybridizing to targeted mRNA and preventing protein translation. Antisense approaches involve the design of oligonucleotides that are complementary to a target gene mRNA. The antisense oligonucleotides will bind to the complementary target gene mRNA
25 transcripts and prevent translation. Absolute complementarity, although preferred, is not required.

A sequence "complementary" to a portion of an RNA, as referred to herein, means a sequence having sufficient complementarity to be able to hybridize with the RNA, forming a stable duplex; in the case of double-stranded antisense nucleic acids, a single
30 strand of the duplex DNA may thus be tested, or triplex formation may be assayed. The ability to hybridize will depend on both the degree of complementarity and the length of the antisense nucleic acid. Generally, the longer the hybridizing nucleic acid, the more base mismatches with an RNA it may contain and still form a stable duplex (or triplex, as the case may be). One skilled in the art can ascertain a tolerable degree of mismatch by use of
35 standard procedures to determine the melting point of the hybridized complex.

In one embodiment, oligonucleotides complementary to non-coding regions of the FBP gene could be used in an antisense approach to inhibit translation of endogenous FBP mRNA. Antisense nucleic acids should be at least six nucleotides in length, and are preferably oligonucleotides ranging from 6 to about 50 nucleotides in length. In specific
5 aspects the oligonucleotide is at least 10 nucleotides, at least 17 nucleotides, at least 25 nucleotides or at least 50 nucleotides.

In an embodiment of the present invention, oligonucleotides complementary to the nucleic acids encoding the F-box motif as indicated in Figures 2 and 4-9.

Regardless of the choice of target sequence, it is preferred that in vitro
10 studies are first performed to quantitate the ability of the antisense oligonucleotide to inhibit gene expression. It is preferred that these studies utilize controls that distinguish between antisense gene inhibition and nonspecific biological effects of oligonucleotides. It is also preferred that these studies compare levels of the target RNA or protein with that of an internal control RNA or protein. Additionally, it is envisioned that results obtained using
15 the antisense oligonucleotide are compared with those obtained using a control oligonucleotide. It is preferred that the control oligonucleotide is of approximately the same length as the test oligonucleotide and that the nucleotide sequence of the oligonucleotide differs from the antisense sequence no more than is necessary to prevent specific hybridization to the target sequence.

20 The oligonucleotides can be DNA or RNA or chimeric mixtures or derivatives or modified versions thereof, single-stranded or double-stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone, for example, to improve stability of the molecule, hybridization, *etc.* The oligonucleotide may include other appended groups such as peptides (*e.g.*, for targeting host cell receptors
25 in vivo), or agents facilitating transport across the cell membrane (see, *e.g.*, Letsinger, et al., 1989, Proc. Natl. Acad. Sci. U.S.A. 86, 6553-6556; Lemaitre, et al., 1987, Proc. Natl. Acad. Sci. 84, 648-652; PCT Publication No. WO88/09810, published December 15, 1988) or the blood-brain barrier (see, *e.g.*, PCT Publication No. WO89/10134, published April 25, 1988), hybridization-triggered cleavage agents (see, *e.g.*, Krol et al., 1988, BioTechniques 6,
30 958-976) or intercalating agents (see, *e.g.*, Zon, 1988, Pharm. Res. 5, 539-549). To this end, the oligonucleotide may be conjugated to another molecule, *e.g.*, a peptide, hybridization triggered cross-linking agent, transport agent, hybridization-triggered cleavage agent, *etc.*

The antisense oligonucleotide may comprise at least one modified base moiety which is selected from the group including but not limited to 5-fluorouracil, 5-
35 bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-

(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine.

The antisense oligonucleotide may also comprise at least one modified sugar moiety selected from the group including but not limited to arabinose, 2-fluoroarabinose, xylulose, and hexose.

In yet another embodiment, the antisense oligonucleotide comprises at least one modified phosphate backbone selected from the group consisting of a phosphorothioate (S-ODNs), a phosphorodithioate, a phosphoramidothioate, a phosphoramidate, a phosphordiamidate, a methylphosphonate, an alkyl phosphotriester, and a formacetal or analog thereof.

In yet another embodiment, the antisense oligonucleotide is an α -anomeric oligonucleotide. An α -anomeric oligonucleotide forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual β -units, the strands run parallel to each other (Gautier, et al., 1987, Nucl. Acids Res. 15, 6625-6641). The oligonucleotide is a 2'-O-methylribonucleotide (Inoue, et al., 1987, Nucl. Acids Res. 15, 6131-6148), or a chimeric RNA-DNA analogue (Inoue, et al., 1987, FEBS Lett. 215, 327-330).

Oligonucleotides of the invention may be synthesized by standard methods known in the art, *e.g.* by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, *etc.*). As examples, phosphorothioate oligonucleotides may be synthesized by the method of Stein, et al. (1988, Nucl. Acids Res. 16, 3209), methylphosphonate oligonucleotides can be prepared by use of controlled pore glass polymer supports (Sarin, et al., 1988, Proc. Natl. Acad. Sci. U.S.A. 85, 7448-7451), *etc.*

While antisense nucleotides complementary to the target gene coding region sequence could be used, those complementary to the transcribed, untranslated region are most preferred.

In one embodiment of the present invention, gene expression downregulation is achieved because specific target mRNAs are digested by RNase H after they have hybridized with the antisense phosphorothioate oligonucleotides (S-ODNs). Since no rules exist to predict which antisense S-ODNs will be more successful, the best strategy is

5 completely empirical and consists of trying several antisense S-ODNs. Antisense phosphorothioate oligonucleotides (S-ODNs) will be designed to target specific regions of mRNAs of interest. Control S-ODNs consisting of scrambled sequences of the antisense S-ODNs will also be designed to assure identical nucleotide content and minimize differences potentially attributable to nucleic acid content. All S-ODNs will be synthesized by Oligos

10 *Etc.* (Wilsonville, OR). In order to test the effectiveness of the antisense molecules when applied to cells in culture, such as assays for research purposes or ex vivo gene therapy protocols, cells will be grown to 60-80% confluence on 100 mm tissue culture plates, rinsed with PBS and overlaid with lipofection mix consisting of 8 ml Opti-MEM, 52.8 μ l Lipofectin, and a final concentration of 200 nM S-ODNs. Lipofections will be carried out

15 using Lipofectin Reagent and Opti-MEM (Gibco BRL). Cells will be incubated in the presence of the lipofection mix for 5 hours. Following incubation the medium will be replaced with complete DMEM. Cells will be harvested at different time points post-lipofection and protein levels will be analyzed by Western blot.

Antisense molecules should be targeted to cells that express the target gene,

20 either directly to the subject in vivo or to cells in culture, such as in ex vivo gene therapy protocols. A number of methods have been developed for delivering antisense DNA or RNA to cells; *e.g.*, antisense molecules can be injected directly into the tissue site, or modified antisense molecules, designed to target the desired cells (*e.g.*, antisense linked to peptides or antibodies that specifically bind receptors or antigens expressed on the target

25 cell surface) can be administered systemically.

However, it is often difficult to achieve intracellular concentrations of the antisense sufficient to suppress translation of endogenous mRNAs. Therefore a preferred approach utilizes a recombinant DNA construct in which the antisense oligonucleotide is placed under the control of a strong pol III or pol II promoter. The use of such a construct

30 to transfect target cells in the patient will result in the transcription of sufficient amounts of single stranded RNAs that will form complementary base pairs with the endogenous target gene transcripts and thereby prevent translation of the target gene mRNA. For example, a vector can be introduced *e.g.*, such that it is taken up by a cell and directs the transcription of an antisense RNA. Such a vector can remain episomal or become chromosomally

35 integrated, as long as it can be transcribed to produce the desired antisense RNA. Such

vectors can be constructed by recombinant DNA technology methods standard in the art. Vectors can be plasmid, viral, or others known in the art, used for replication and expression in mammalian cells. Expression of the sequence encoding the antisense RNA can be by any promoter known in the art to act in mammalian, preferably human cells. Such promoters
5 can be inducible or constitutive. Such promoters include but are not limited to: the SV40 early promoter region (Bernoist and Chambon, 1981, Nature 290, 304-310), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto, et al., 1980, Cell 22, 787-797), the herpes thymidine kinase promoter (Wagner, et al., 1981, Proc. Natl. Acad. Sci. U.S.A. 78, 1441-1445), the regulatory sequences of the metallothionein gene
10 (Brinster, et al., 1982, Nature 296, 39-42), *etc.* Any type of plasmid, cosmid, YAC or viral vector can be used to prepare the recombinant DNA construct which can be introduced directly into the tissue site. Alternatively, viral vectors can be used that selectively infect the desired tissue, in which case administration may be accomplished by another route (*e.g.*, systemically).

15 Ribozyne molecules designed to catalytically cleave target gene mRNA transcripts can also be used to prevent translation of target gene mRNA and, therefore, expression of target gene product (see, *e.g.*, PCT International Publication WO90/11364, published October 4, 1990; Sarver, et al., 1990, Science 247, 1222-1225). In an embodiment of the present invention, oligonucleotides which hybridize to the FBP gene are
20 designed to be complementary to the nucleic acids encoding the F-box motif as indicated in Figures 2 and 4-9.

Ribozymes are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA. (For a review, see Rossi, 1994, Current Biology 4, 469-471). The mechanism of ribozyme action involves sequence specific hybridization of the ribozyme
25 molecule to complementary target RNA, followed by an endonucleolytic cleavage event. The composition of ribozyme molecules must include one or more sequences complementary to the target gene mRNA, and must include the well known catalytic sequence responsible for mRNA cleavage. For this sequence, see, *e.g.*, U.S. Patent No. 5,093,246, which is incorporated herein by reference in its entirety.

30 While ribozymes that cleave mRNA at site specific recognition sequences can be used to destroy target gene mRNAs, the use of hammerhead ribozymes is preferred. Hammerhead ribozymes cleave mRNAs at locations dictated by flanking regions that form complementary base pairs with the target mRNA. The sole requirement is that the target mRNA have the following sequence of two bases: 5'-UG-3'. The construction and
35 production of hammerhead ribozymes is well known in the art and is described more fully

in Myers, 1995, Molecular Biology and Biotechnology: A Comprehensive Desk Reference, VCH Publishers, New York, (see especially Figure 4, page 833) and in Haseloff & Gerlach, 1988, Nature, 334, 585-591, which is incorporated herein by reference in its entirety.

Preferably the ribozyme is engineered so that the cleavage recognition site is
5 located near the 5' end of the target gene mRNA, *i.e.*, to increase efficiency and minimize the intracellular accumulation of non-functional mRNA transcripts.

The ribozymes of the present invention also include RNA endoribonucleases (hereinafter "Cech-type ribozymes") such as the one that occurs naturally in *Tetrahymena thermophila* (known as the IVS, or L-19 IVS RNA) and that has been extensively described
10 by Thomas Cech and collaborators (Zaug, et al., 1984, Science, 224, 574-578; Zaug and Cech, 1986, Science, 231, 470-475; Zaug, et al., 1986, Nature, 324, 429-433; published International patent application No. WO 88/04300 by University Patents Inc.; Been & Cech, 1986, Cell, 47, 207-216). The Cech-type ribozymes have an eight base pair active site which hybridizes to a target RNA sequence whereafter cleavage of the target RNA takes
15 place. The invention encompasses those Cech-type ribozymes which target eight base-pair active site sequences that are present in the target gene.

As in the antisense approach, the ribozymes can be composed of modified oligonucleotides (*e.g.*, for improved stability, targeting, *etc.*) and should be delivered to cells that express the target gene in vivo. A preferred method of delivery involves using a
20 DNA construct "encoding" the ribozyme under the control of a strong constitutive pol III or pol II promoter, so that transfected cells will produce sufficient quantities of the ribozyme to destroy endogenous target gene messages and inhibit translation. Because ribozymes unlike antisense molecules, are catalytic, a lower intracellular concentration is required for efficiency.

25 Endogenous target gene expression can also be reduced by inactivating or "knocking out" the target gene or its promoter using targeted homologous recombination (*e.g.*, see Smithies, et al., 1985, Nature 317, 230-234; Thomas & Capecchi, 1987, Cell 51, 503-512; Thompson, et al., 1989, Cell 5, 313-321; each of which is incorporated by reference herein in its entirety). For example, a mutant, non-functional target gene (or a
30 completely unrelated DNA sequence) flanked by DNA homologous to the endogenous target gene (either the coding regions or regulatory regions of the target gene) can be used, with or without a selectable marker and/or a negative selectable marker, to transfect cells that express the target gene in vivo. Insertion of the DNA construct, via targeted homologous recombination, results in inactivation of the target gene. Such approaches are
35 particularly suited modifications to ES (embryonic stem) cells can be used to generate

animal offspring with an inactive target gene (*e.g.*, see Thomas & Capecchi, 1987 and Thompson, 1989, *supra*). However this approach can be adapted for use in humans provided the recombinant DNA constructs are directly administered or targeted to the required site *in vivo* using appropriate viral vectors.

- 5 Alternatively, endogenous target gene expression can be reduced by targeting deoxyribonucleotide sequences complementary to the regulatory region of the target gene (*i.e.*, the target gene promoter and/or enhancers) to form triple helical structures that prevent transcription of the target gene in target cells in the body. (See generally, Helene, 1991, *Anticancer Drug Des.*, 6(6), 569-584; Helene, et al., 1992, *Ann. N.Y. Acad. Sci.*, 660, 27-10 36; and Maher, 1992, *Bioassays* 14(12), 807-815).

- Nucleic acid molecules to be used in triple helix formation for the inhibition of transcription should be single stranded and composed of deoxynucleotides. The base composition of these oligonucleotides must be designed to promote triple helix formation via Hoogsteen base pairing rules, which generally require sizeable stretches of either purines 15 or pyrimidines to be present on one strand of a duplex. Nucleotide sequences may be pyrimidine-based, which will result in TAT and CGC+ triplets across the three associated strands of the resulting triple helix. The pyrimidine-rich molecules provide base complementarity to a purine-rich region of a single strand of the duplex in a parallel orientation to that strand. In addition, nucleic acid molecules may be chosen that are 20 purine-rich, for example, contain a stretch of G residues. These molecules will form a triple helix with a DNA duplex that is rich in GC pairs, in which the majority of the purine residues are located on a single strand of the targeted duplex, resulting in GGC triplets across the three strands in the triplex.

- Alternatively, the potential sequences that can be targeted for triple helix 25 formation may be increased by creating a so called "switchback" nucleic acid molecule. Switchback molecules are synthesized in an alternating 5'-3', 3'-5' manner, such that they base pair with first one strand of a duplex and then the other, eliminating the necessity for a sizeable stretch of either purines or pyrimidines to be present on one strand of a duplex.

- In instances wherein the antisense, ribozyme, and/or triple helix molecules 30 described herein are utilized to inhibit mutant gene expression, it is possible that the technique may so efficiently reduce or inhibit the transcription (triple helix) and/or translation (antisense, ribozyme) of mRNA produced by normal target gene alleles that the possibility may arise wherein the concentration of normal target gene product present may be lower than is necessary for a normal phenotype. In such cases, to ensure that 35 substantially normal levels of target gene activity are maintained, therefore, nucleic acid

molecules that encode and express target gene polypeptides exhibiting normal target gene activity may, be introduced into cells via gene therapy methods such as those described, below, in Section 5.7.2 that do not contain sequences susceptible to whatever antisense, ribozyme, or triple helix treatments are being utilized. Alternatively, in instances whereby
5 the target gene encodes an extracellular protein, it may be preferable to co-administer normal target gene protein in order to maintain the requisite level of target gene activity.

Anti-sense RNA and DNA, ribozyme, and triple helix molecules of the invention may be prepared by any method known in the art for the synthesis of DNA and RNA molecules, as discussed above. These include techniques for chemically synthesizing
10 oligodeoxyribonucleotides and oligoribonucleotides well known in the art such as for example solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding the antisense RNA molecule. Such DNA sequences may be incorporated into a wide variety of vectors that incorporate suitable RNA polymerase promoters such as the T7 or SP6
15 polymerase promoters. Alternatively, antisense cDNA constructs that synthesize antisense RNA constitutively or inducibly, depending on the promoter used, can be introduced stably into cell lines.

5.7.2 GENE REPLACEMENT THERAPY

20 With respect to an increase in the level of normal FBP gene expression and/or FBP gene product activity, FBP gene nucleic acid sequences, described, above, in Section 5.1 can, for example, be utilized for the treatment of proliferative disorders such as cancer. Such treatment can be administered, for example, in the form of gene replacement
25 therapy. Specifically, one or more copies of a normal FBP gene or a portion of the FBP gene that directs the production of an FBP gene product exhibiting normal FBP gene function, may be inserted into the appropriate cells within a patient, using vectors that include, but are not limited to adenovirus, adeno-associated virus, and retrovirus vectors, in addition to other particles that introduce DNA into cells, such as liposomes.

For FBP genes that are expressed in all tissues or are preferentially
30 expressed, such as FBP1 gene is expressed preferably in the brain, such gene replacement therapy techniques should be capable delivering FBP gene sequences to these cell types within patients. Thus, in one embodiment, techniques that are well known to those of skill in the art (see, e.g., PCT Publication No. WO89/10134, published April 25, 1988) can be used to enable FBP gene sequences to cross the blood-brain barrier readily and to deliver
35 the sequences to cells in the brain. With respect to delivery that is capable of crossing the

blood-brain barrier, viral vectors such as, for example, those described above, are preferable.

In another embodiment, techniques for delivery involve direct administration of such FBP gene sequences to the site of the cells in which the FBP gene sequences are to be expressed.

Additional methods that may be utilized to increase the overall level of FBP gene expression and/or FBP gene product activity include the introduction of appropriate FBP-expressing cells, preferably autologous cells, into a patient at positions and in numbers that are sufficient to ameliorate the symptoms of an FBP disorder. Such cells may be either recombinant or non-recombinant.

Among the cells that can be administered to increase the overall level of FBP gene expression in a patient are cells that normally express the FBP gene.

Alternatively, cells, preferably autologous cells, can be engineered to express FBP gene sequences, and may then be introduced into a patient in positions appropriate for the amelioration of the symptoms of an FBP disorder or a proliferative or differentiative disorders, *e.g.*, cancer and tumorigenesis. Alternately, cells that express an unimpaired FBP gene and that are from a MHC matched individual can be utilized, and may include, for example, brain cells. The expression of the FBP gene sequences is controlled by the appropriate gene regulatory sequences to allow such expression in the necessary cell types. Such gene regulatory sequences are well known to the skilled artisan. Such cell-based gene therapy techniques are well known to those skilled in the art, see, *e.g.*, Anderson, U.S. Patent No. 5,399,349.

When the cells to be administered are non-autologous cells, they can be administered using well known techniques that prevent a host immune response against the introduced cells from developing. For example, the cells may be introduced in an encapsulated form which, while allowing for an exchange of components with the immediate extracellular environment, does not allow the introduced cells to be recognized by the host immune system.

Additionally, compounds, such as those identified via techniques such as those described, above, in Section 5.5, that are capable of modulating FBP gene product activity can be administered using standard techniques that are well known to those of skill in the art. In instances in which the compounds to be administered are to involve an interaction with brain cells, the administration techniques should include well known ones that allow for a crossing of the blood-brain barrier.

5.7.3 TARGET PROLIFERATIVE CELL DISORDERS

With respect to specific proliferative and oncogenic disease associated with ubiquitin ligase activity, the diseases that can be treated or prevented by the methods of the present invention include but are not limited to: human sarcomas and carcinomas, *e.g.*,
5 fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland
10 carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilms' tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma,
15 pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, melanoma, neuroblastoma, retinoblastoma; leukemias, *e.g.*, acute lymphocytic leukemia and acute myelocytic leukemia (myeloblastic, promyelocytic, myelomonocytic, monocytic and erythroleukemia); chronic leukemia (chronic myelocytic (granulocytic) leukemia and chronic lymphocytic leukemia); and polycythemia vera, lymphoma (Hodgkin's disease and
20 non-Hodgkin's disease), multiple myeloma, Waldenström's macroglobulinemia, and heavy chain disease.

Diseases and disorders involving a deficiency in cell proliferation or in which cell proliferation is desired for treatment or prevention, and that can be treated or prevented by inhibiting FBP function, include but are not limited to degenerative disorders,
25 growth deficiencies, hypoproliferative disorders, physical trauma, lesions, and wounds; for example, to promote wound healing, or to promote regeneration in degenerated, lesioned or injured tissues, *etc.* In a specific embodiment, nervous system disorders are treated. In another specific embodiment, a disorder that is not of the nervous system is treated.

30 5.8 PHARMACEUTICAL PREPARATIONS AND METHODS OF ADMINISTRATION

The compounds that are determined to affect FBP gene expression or gene product activity can be administered to a patient at therapeutically effective doses to treat or ameliorate a cell proliferative disorder. A therapeutically effective dose refers to that
35 amount of the compound sufficient to result in amelioration of symptoms of such a disorder.

5.8.1 EFFECTIVE DOSE

Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, for determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD50/ED50. Compounds that exhibit large therapeutic indices are preferred. While compounds that exhibit toxic side effects may be used, care should be taken to design a delivery system that targets such compounds to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range that includes the IC50 (*i.e.*, the concentration of the test compound that achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

5.8.2 FORMULATIONS AND USE

Pharmaceutical compositions for use in accordance with the present invention may be formulated in conventional manner using one or more physiologically acceptable carriers or excipients.

Thus, the compounds and their physiologically acceptable salts and solvates may be formulated for administration by inhalation or insufflation (either through the mouth or the nose) or oral, buccal, parenteral or rectal administration.

For oral administration, the pharmaceutical compositions may take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (*e.g.*, pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (*e.g.*, lactose, microcrystalline cellulose or calcium hydrogen phosphate); lubricants (*e.g.*, magnesium stearate, talc or silica); disintegrants (*e.g.*, potato starch or sodium starch glycolate); or

wetting agents (*e.g.*, sodium lauryl sulphate). The tablets may be coated by methods well known in the art. Liquid preparations for oral administration may take the form of, for example, solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations may
5 be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (*e.g.*, sorbitol syrup, cellulose derivatives or hydrogenated edible fats); emulsifying agents (*e.g.*, lecithin or acacia); non-aqueous vehicles (*e.g.*, almond oil, oily esters, ethyl alcohol or fractionated vegetable oils); and preservatives (*e.g.*, methyl or propyl-p-hydroxybenzoates or sorbic acid). The preparations may also contain buffer salts,
10 flavoring, coloring and sweetening agents as appropriate.

Preparations for oral administration may be suitably formulated to give controlled release of the active compound.

For buccal administration the compositions may take the form of tablets or lozenges formulated in conventional manner.

15 For administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebuliser, with the use of a suitable propellant, *e.g.*, dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit may be determined
20 by providing a valve to deliver a metered amount. Capsules and cartridges of *e.g.*, gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

The compounds may be formulated for parenteral administration by injection, *e.g.*, by bolus injection or continuous infusion. Formulations for injection may be
25 presented in unit dosage form, *e.g.*, in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, *e.g.*, sterile pyrogen-free water, before use.

30 The compounds may also be formulated in rectal compositions such as suppositories or retention enemas, *e.g.*, containing conventional suppository bases such as cocoa butter or other glycerides.

In addition to the formulations described previously, the compounds may also be formulated as a depot preparation. Such long acting formulations may be
35 administered by implantation (for example subcutaneously or intramuscularly) or by

intramuscular injection. Thus, for example, the compounds may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

5 The compositions may, if desired, be presented in a pack or dispenser device that may contain one or more unit dosage forms containing the active ingredient. The pack may for example comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration.

10

6. EXAMPLE: IDENTIFICATION AND CHARACTERIZATION OF NOVEL UBIQUITIN LIGASE F-BOX PROTEINS AND GENES

The following studies were carried out to identify novel F-box proteins which may act to recruit novel specific substrates to the ubiquitination pathways. Studies
15 involving several organisms have shown that some FBPs play a crucial role in the controlled degradation of important cellular regulatory proteins (*e.g.*, cyclins, cdk-inhibitors, β -catenin, IKBa, *etc.*). These FBPs are subunits of ubiquitin protein SCF ligases formed by three basic subunits: a cullin subunit (called Cdc53 in *S. cerevisiae* and Cul1 in humans); Skp1; and one of many FBPs. SCF ligases target ubiquitin conjugating enzymes (either
20 Ubc3 or Ubc4) to specific substrates which are recruited by different FBPs. Schematically, the Ubc is bound to the ligase through the cullin subunit while the substrate interacts with the FBP subunit. Although FBPs can bind the cullin subunit directly, the presence of fourth subunit, Skp1, which simultaneously can bind the cullin –terminus and the F-box of the FBP, stabilizes the complex. Thus, the substrate specificity of the ubiquitin ligase complex
25 is provided by the F-box subunit.

6.1 MATERIALS AND METHODS USED FOR THE IDENTIFICATION AND CHARACTERIZATION OF NOVEL F-BOX GENES

Yeast Two-Hybrid Screening In order to clone the human genes encoding F-box proteins,
30 proteins associated with Skp1 were identified using a modified yeast 2-hybrid system (Vidal et al., 1996, Proc. Nat. Acad. Sci., 93:10315-20; Vidal et al., 1996, Proc. Nat. Acad. Sci., 93:10321-26). This modified system takes advantage of using three reporter genes expressed from three different Gal4 binding site promoters, thereby decreasing the number of false positive interactions. This multiple reporter gene assay facilitates identification of
35 true interactors.

Human Skp1 was used as a bait to search for proteins that interact with Skp1, such as novel F-box proteins and the putative human homolog of Cdc4. The plasmids pPC97-CYH2 and pPC86 plasmids, encoding the DNA binding domain (DB, aa 1 - 147) and the transcriptional activation domain (AD, aa 768 - 881) of yeast GAL4, and containing
5 LEU2 and TRP1 as selectable markers, respectively, were used (Chevray and Nathans, 1992, Proc. Nat. Acad. Sci., 89:5789-93; Vidal et al., *supra*).

An in-frame fusion between Skp1 and DB was obtained by homologous recombination of the PCR product described below. The following 2 oligonucleotides were designed and obtained as purified primers from Gene Link Inc.: 5'-AGT-AGT-AAC-
10 AAA-GGT-CAA-AGA-CAG-TTG-ACT-GTA-TCG-TCG-AGG-ATG-CCT-TCA-ATT-AAG-TT (SEQ ID NO: 80); 3'-GCG-GTT-ACT-TAC-TTA-GAG-CTC-GAC-GTC-TTA-CTT-ACT-TAG-CTC-ACT-TCT-CTT-CAC-ACC-A (SEQ ID NO: 81). The 5' primer corresponds to a sequence located in the DB of the pPC97-CYH2 plasmid (underlined) flanked by the 5' sequence of the *skp1* gene. The 3' primer corresponds to a sequence
15 located by polylinker of the pPC97-CYH2 plasmid (underlined) flanked by the 3' sequence of the *skp1* gene. These primers were used in a PCR reaction containing the following components: 100 ng DNA template (*skp1* pET plasmid), 1 μ M of each primer, 0.2 mM dNTP, 2 mM MgCl₂, 10 mM KCl, 20 mM TrisCl pH 8.0, 0.1% Triton X-100, 6 mM (NH₄)₂SO₄, 10 μ g/ml nuclease-free BSA, 1 unit of Pfu DNA polymerase (4' at 94°C, 1' at 50°C, 10'
20 at 72°C for 28 cycles). Approximately 100 ng of PCR product were transformed into yeast cells (MaV103 strain; Vidal et al., 1996, Proc. Natl. Acad. Sci. U.S.A. 93:10315-10320; Vidal et al., 1996, Proc. Natl. Acad. Sci. U.S.A. 93:10321-10326) in the presence or in the absence of 100 ng of pPC97-CYH2 plasmid previously digested with BglII and Sall. As a result of the homologous recombination, only yeast cells containing the pPC97-CYH2
25 plasmid homologously recombined with *skp1* cDNA, grew in the absence of leucine. Six colonies were isolated and analyzed by immunoblotting for the expression of Skp1, as described (Vidal et al., *supra*). All 6 colonies, but not control colonies, expressed a Mr 36.000 fusion-protein that was recognized by our affinity purified anti-Skp1 antibody.

The AD fusions were generated by cloning cDNA fragments in the frame
30 downstream of the AD domains and constructs were confirmed by sequencing, immunoblot, and interaction with Skp1. The pPC86-Skp2s (pPC86) include: pPC86-Skp2, and pPC86-Skp2-CT (aa 181-435 of Skp2). The first fusion represents our positive control since Skp2 is a known interactor of Skp1 (Zhang, et al, 1995, Cell, 82: 915-25); the latter fusion was used as a negative control since it lacked the F-box required for the interaction with Skp1.

35

MaV103 strain harboring the DB-skp1 fusions was transformed with an activated T-cell cDNA library (Alala 2; Hu, et al., Genes & Dev. 11: 2701-14) in pPC86 using the standard lithium acetate method. Transformants were first plated onto synthetic complete (SC)-Leu-Trp plates, followed by replica plating onto (SC)-Leu-Trp-His plates containing 20 mM 3-aminotriazole (3-AT) after 2 days. Yeast colonies grown out after additional 3-4 days of incubation were picked as primary positives and further tested in three reporter assays: i) growth on SC-Leu-Trp-His plates supplemented with 20 mM 3-AT; ii) -galactosidase activity; and iii) URA3 activation on SC-Leu-Trp plates containing 0.2% 5-fluoroorotic acid, as a counterselection method. Of the 3×10^6 yeast transformants screened AD plasmids were rescued from the fifteen selected positive colonies after all three. MaV103 cells were re-transformed with either rescued AD plasmids and the DBskp1 fusion or rescued AD plasmid and the pPC97-CYH2vector without a cDNA insert as control. Eleven AD plasmids from colonies that repeatedly tested positive in all three reporter assays (very strong interactors) and four additional AD plasmids from clones that were positive on some but not all three reporter assays (strong interactors) were recovered and sequenced with the automated ABI 373 DNA sequencing system.

Cloning of full length FBPs Two of the clones encoding FBP4 and FBP5 appeared to be full-length, while full length clones of 4 other cDNAs encoding FBP1, FBP2, FBP3 and FBP7 were obtained with RACE using Marathon-Ready cDNA libraries (Clontech, cat. # 7406, 7445, 7402) according to the manufacturer's instructions. A full-length clone encoding FBP6 was not obtained. Criteria for full length clones included at least two of the following: i) the identification of an ORF yielding a sequence related to known F-box proteins; ii) the presence of a consensus Kozak translation initiation sequence at a putative initiator methionine codon; iii) the identification of a stop codon in the same reading frame but upstream of the putative initiation codon; iv) the inability to further increase the size of the clone by RACE using three different cDNA libraries.

Analysis by Immunoblotting of Protein from Yeast Extracts Yeast cells were grown to mid-logarithmic phase, harvested, washed and resuspended in buffer (50 mM Tris pH 8.0, 20% glycerol, 1 mM EDTA, 0.1% Triton X-100, 5 mM MgCl₂, 10 mM β -mercaptoethanol, 1 mM PMSF, 1 mg/ml Leupeptin, 1 mg/ml Pepstatin) at a cell density of about 109 cells/ml. Cells were disrupted by vortexing in the presence of glass beads for 10 min at 40C. Debris was pelleted by centrifugation at 12,000 RPM for 15 min at 40C. Approximately 50 g of

proteins were subjected to immunoblot analysis as described (Vidal et al., 1996a, *supra*; Vidal et al., 1996b, *supra*).

DNA database searches and analysis of protein motifs ESTs (expressed sequence tags)

- 5 with homology to FBP genes were identified using BLAST, PSI-BLAST (<http://www.ncbi.nlm.nih.gov/BLAST/>) and TGI Sequence Search (http://www.tigr.org/cgi-bin/BlastSearch/blast_tgi.cgi). ESTs that overlapped more than 95 % in at least 100 bps were assembled into novel contiguous ORFs using Sequencher 3.0. Protein domains were identified with ProfileScan Server
- 10 (http://www.isrec.isb-sib.ch/software/PFSCAN_form.html), BLOCKS Sercher (http://www.blocks.fhcrc.org/blocks_search.html) and IMB Jena (<http://genome.imb-jena.de/cgi-bin/GDEWWW/menu.cgi>).

Construction of F-box mutants Delta-F-box mutants [(ΔF)FBP1, residues 32-179;

- 15 (ΔF)FBP2, residues 60-101; (ΔF)FBP3a, residues 40-76; (ΔF)FBP4, residues 55-98] were obtained by deletion with the appropriate restriction enzymes with conservation of the reading frame. (ΔF)Skp2 mutant was obtained by removing a DNA fragment (nucleotides 338-997) with BspEI and XbaI restriction enzymes, and replacing it with a PCR fragment containing nucleotides 457 to 997. The final construct encoded a protein lacking residues
- 20 113-152. The leucine 51-to-alanine FBP3a mutant [FBP3a(L51A)] and the tryptophan 76-to-alanine FBP3a mutant [FBP3a(W76A)] were generated by oligonucleotide-directed mutagenesis using the polymerase chain reaction of the QuikChange site-directed mutagenesis kit (Stratagene). All mutants were sequenced in their entirety.

- 25 Recombinant proteins cDNA fragments encoding the following human proteins: Flag-tagged FBP1, Flag-tagged (ΔF)FBP1, Flag-tagged FBP3a, Skp2, HA-tagged Cul1, HA-tagged Cul2, (β-catenin, His-tagged cyclin D1, Skp1, His-tagged Skp1, His-tagged Elongin C were inserted into the baculovirus expression vector pBacpak-8 (Clontech) and cotransfected into Sf9 cells with linearized baculovirus DNA using the BaculoGold
- 30 transfection kit (Pharmingen). Recombinant viruses were used to infect 5B cells and assayed for expression of their encoded protein by immunoblotting as described above. His-proteins were purified with Nickel-agarose (Invitrogen) according to the manufacturer's instructions.

Antibodies. Anti-Cul1 antibodies was generated by injecting rabbits and mice with the following amino acid peptide: (C)DGEKDTYSYLA (SEQ ID NO: 82). This peptide corresponds to the carboxy-terminus of human Cul1 and is not conserved in other cullins. Anti-Cul2 antibodies was generated by injecting rabbits with the following amino acid peptide: (C)ESSFSLNMFSSKRTKFKITTSMQ (SEQ ID NO: 83). This peptide is located 87 amino acids from the carboxy-terminus of human Cul2 and is not conserved in other cullins. The anti-Skp1 antibody was generated by injecting rabbits with the peptide (C)EEAQVRKENQW (SEQ ID NO: 84), corresponding to the carboxy-terminus of human Skp1. The cysteine residues (C) were added in order to couple the peptides to keyhole limpet hemocyanin (KLH). All of the antibodies were generated, affinity-purified (AP) and characterized as described (Pagano, M., ed., 1995, "From Peptide to Purified Antibody", in Cell Cycle: Materials and Methods, Springer-Verlag, 217-281). Briefly, peptides whose sequence showed high antigenic index (high hydrophilicity, good surface probability, good flexibility, and good secondary structure) were chosen. Rabbits and mice were injected with peptide-KLH mixed with complete Freund's adjuvant. Subsequently they were injected with the peptide in incomplete Freund's adjuvant, every 2 weeks, until a significant immunoreactivity was detected by immunoprecipitation of 35S-methionine labeled HeLa extract. These antisera recognized bands at the predicted size in both human extracts and extracts containing recombinant proteins.

Monoclonal antibody (Mab) to Ubc3 was generated and characterized in collaboration with Zymed Inc. Mab to cyclin B (cat # sc-245) was from Santa Cruz; Mabs to p21 (cat # C24420) and p27 (cat # K25020) from Transduction lab. (Mabs) cyclin E, (Faha, 1993, J. of Virology 67: 2456); AP rabbit antibodies to human p27, Skp2, Cdk2 (Pagano, 1992, EMBO J. 11: 761), and cyclin A (Pagano, 1992, EMBO J. 11: 761), and phospho-site p27 specific antibody, were obtained or generated by standard methods. Where indicated, an AP goat antibody to an N-terminal Skp2 peptide (Santa Cruz, cat # sc-1567) was used. Rat anti-HA antibody was from Boehringer Mannheim (cat. #1867423), rabbit anti-HA antibody was from Santa Cruz (cat. # sc-805), mouse anti-Flag antibody was from Kodak (cat. # IB13010), rabbit anti-Flag antibody was from Zymed (cat. #71-5400), anti-Skp1 and anti-(β -catenin mouse antibodies were from Transduction Laboratories (cat. # C19220 and P46020, respectively). The preparation, purification and characterization of a Mab to human cyclin D1 (clone AM29, cat. #33-2500) was performed in collaboration with Zymed Inc. Antiserum to human cyclin D1 was produced as described(Ohtsubo et al., 1995, Mol Cell Biol, 15, 2612-2624).

35

Extract preparation and cell synchronization Protein extraction was performed as previously described (Pagano, 1993, J. Cell Biol. 121: 101) with the only difference that 1 μ M okadaic acid was present in the lysis buffer. Human lung fibroblasts IMR-90 were synchronized in G0/G1 by serum starvation for 48 hours and then restimulated to re-enter the cell cycle by serum readdition. HeLa cells were synchronized by mitotic shake-off as described (Pagano, 1992, EMBO J. 11: 761). Synchronization was monitored by flow cytometry. For in vitro ubiquitination and degradation assays, G1 HeLa cells were obtained with a 48-hour lovastatin treatment and protein extraction performed as described below.

- 10 Immunoprecipitation and Immunoblotting. Cell extracts were prepared by addition of 3-5 volumes of standard lysis buffers (Pagano et al., 1992, Science 255, 1144-1147), and conditions for immunoprecipitation were as described (Jenkins and Xiong, 1995; Pagano et al., 1992a Science 255-1144-1147). Proteins were transferred from gel to a nitrocellulose membrane (Novex) by wet blotting as described (Tam et al., 1994 Oncogene 9, 2663).
- 15 Filters were subjected to immunoblotting using a chemiluminescence (DuPont-NEN) detection system according to the manufacturer's instructions

- Protein extraction for in vitro ubiquitination assay Logarithmically growing, HeLa-S3 cells were collected at a density of 6×10^5 cells/ml. Approx. 4 ml of HeLa S3 cell pellet were
- 20 suspended in 6 ml of ice-cold buffer consisting of 20 mM Tris-HCl (pH 7.2), 2 mM DTT, 0.25 mM EDTA, 10 μ g/ml leupeptin, and 10 μ g/ml pepstatin. The suspension was transferred to a cell nitrogen-disruption bomb (Parr, Moline, IL, cat #4639) that had been rinsed thoroughly and chilled on ice before use. The bomb chamber was connected to a nitrogen tank and the pressure was brought slowly to 1000 psi. The chamber was left on ice
- 25 under the same pressure for 30 minutes and then the pressure was released slowly. The material was transferred to an Eppendorf tube and centrifuged in a microcentrifuge at 10,000 g for 10 minutes. The supernatant (S-10) was divided into smaller samples and frozen at -80°C.

- 30 In vitro ubiquitination The ubiquitination assay was performed as described (Lyapina, 1998, Proc Natl Acad Sci U S A, 95: 7451). Briefly, immuno-beads containing Flag-tagged FBPs immunoprecipitated with anti-Flag antibody were added with purified recombinant human E1 and E2 enzymes (Ubc2, Ubc3 or Ubc4) to a reaction mix containing biotinylated-ubiquitin. Samples were then analyzed by blotting with HRP-streptavidin. E1 and E2

35

enzymes and biotinylated-ubiquitin were produced as described (Pagano, 1995, Science 269: 682).

Transient transfections cDNA fragments encoding the following human proteins:

- 5 FBP1, (Δ F)FBP1, FBP2, (Δ F)FBP2, FBP3a, (Δ F)FBP3a, FBP3a(L51A), FBP3a(W76A), FBP4, (Δ F)FBP4, Skp2, (Δ F)Skp2, HA-tagged β -catenin, untagged β -catenin, Skp1, cyclin D1 were inserted into the mammalian expression vector pcDNA3 (Invitrogen) in frame with a Flag-tag at their C-terminus. Cells were transfected with FuGENE transfection reagent (Boehringer, cat. #1-814-443) according to the manufacture's instruction.

10

Immunofluorescence Transfected cell monolayers growing on glass coverslips were rinsed in PBS and fixed with 4% paraformaldehyde in PBS for 10 minutes at 4°C followed by permeabilization for 10 minutes with 0.25% Triton X-100 in PBS. Other fixation protocols gave comparable results. Immunofluorescence stainings were performed using 1 μ g/ml

- 15 rabbit anti-Flag antibody as described (Pagano, 1994, Genes & Dev., 8:1627).

Northern Blot Analysis

Northern blots were performed using human multiple-tissue mRNAs from Clontech Inc. Probes were radiolabeled with [α -32P] dCTP (Amersham Inc.) using a random primer DNA labeling kit (Gibco BRL) (2 x 10⁶ cpm/ml). Washes

20 were performed with 0.2 x SSC, 0.1% SDS, at 55 - 60°C. FBP1 and FBP3a probes were two HindIII restriction fragments (nucleotides 1 - 571 and 1 - 450, respectively), FBP2, FBP4, and FBP1 probes were their respective full-length cDNAs, and β -ACTIN probe was from Clontech Inc.

- 25 Fluorescence in situ hybridization (FISH) Genomic clones were isolated by high-stringency screening (65°C, 0.2 x SSC, 0.1 % SDS wash) of a λ FIX II placenta human genomic library (Stratagene) with cDNA probes obtained from the 2-hybrid screening. Phage clones were confirmed by high-stringency Southern hybridization and partial sequence analysis. Purified whole phage DNA was labeled and FISH was performed as
- 30 described (M. Pagano., ed., 1994, in Cell Cycle: Materials and Methods, 29).

6.2 RESULTS

6.2.1 Characterization of novel F-box Proteins and their activity in vivo

- 35 An improved version of the yeast two-hybrid system was used to search for interactors of human Skp1. The MaV103 yeast strain harboring the Gal4 DB-Skp1 fusion

protein as bait was transformed with an activated T-cell cDNA library expressing Gal4 AD fusion proteins as prey. After initial selection and re-transformation steps, 3 different reporter assays were used to obtain 13 positive clones that specifically interact with human Skp1. After sequence analysis, the 13 rescued cDNAs were found to be derived from 7 different open reading frames all encoding FBPs. These novel FBPs were named as follows: FBP1, shown in Figure 3 (SEQ ID NO:1); FBP2, shown in Figure 4 (SEQ ID NO:3), FBP3a, shown in Figure 5 (SEQ ID NO:5), FBP4, shown in Figure 7 (SEQ ID NO:7), FBP5, shown in Figure 8 (SEQ ID NO:9), FBP6, shown in Figure 9 (SEQ ID NO:11), FBP7, shown in Figure 10 (SEQ ID NO:13). One of the seven FBPs, FBP1 (SEQ ID NO:1) was also identified by others while our screen was in progress (Margottin et al., 1998, Molecular Cell, 1:565-74).

BLAST programs were used to search for predicted human proteins containing an F-box in databases available through the National Center for Biotechnology Information and The Institute for Genomic Research. The alignment of the F-box motifs from these predicted human FBPs is shown in Figure 1. Nineteen previously uncharacterized human FBPs were identified by aligning available sequences (GenBank Accession Nos. AC002428, AI457595, AI105408, H66467, T47217, H38755, THC274684, AI750732, AA976979, AI571815, T57296, Z44228, Z45230, N42405, AA018063, AI751015, AI400663, T74432, AA402415, AI826000, AI590138, AF174602, Z45775, AF174599, THC288870, AI017603, AF174598, THC260994, AI475671, AA768343, AF174595, THC240016, N70417, T10511, AF174603, EST04915, AA147429, AI192344, AF174594, AI147207, AI279712, AA593015, AA644633, AA335703, N26196, AF174604, AF053356, AF174606, AA836036, AA853045, AI479142, AA772788, AA039454, AA397652, AA463756, AA007384, AA749085, AI640599, THC253263, AB020647, THC295423, AA434109, AA370939, AA215393, THC271423, AF052097, THC288182, AL049953, CAB37981, AL022395, AL031178, THC197682, and THC205131), with the nucleotide sequences derived from the F-box proteins disclosed above.

The nineteen previously uncharacterized FBP nucleotide sequences thus identified were named as follows: FBP3b, shown in Figure 6 (SEQ ID NO:23); FBP8, shown in Figure 11 (SEQ ID NO:25); FBP9, shown in Figure 12 (SEQ ID NO:27); FBP10, shown in Figure 13 (SEQ ID NO:29); FBP11, shown in Figure 14 (SEQ ID NO:31); FBP12, shown in Figure 15 (SEQ ID NO:33); FBP13, shown in Figure 16 (SEQ ID NO:35); FBP14, shown in Figure 17 (SEQ ID NO:37); FBP15, shown in Figure 18 (SEQ ID NO:39); FBP16, shown in Figure 19 (SEQ ID NO:41); FBP17, shown in Figure 20 (SEQ ID NO:43); FBP18, shown in Figure 21 (SEQ ID NO:45); FBP19, shown in Figure 22 (SEQ ID NO:47); FBP20,

shown in Figure 23 (SEQ ID NO:49); FBP21, shown in Figure 24 (SEQ ID NO:51); FBP22, shown in Figure 25 (SEQ ID NO:53); FBP23, shown in Figure 26 (SEQ ID NO:55); FBP24, shown in Figure 27 (SEQ ID NO:57); and FBP25, shown in Figure 28 (SEQ ID NO:59).

The alignment of the F-box motifs from these predicted human FBPs is shown in Figure

- 5 1A. Of these sequences, the nucleotide sequences of fourteen identified FBPs, FBP3b (SEQ ID NO:23), FBP8 (SEQ ID NO:25), FBP11 (SEQ ID NO:31), FBP12 (SEQ ID NO:33), FBP13 (SEQ ID NO:35), FBP14 (SEQ ID NO:37), FBP15 (SEQ ID NO:39), FBP17 (SEQ ID NO:43), FBP18 (SEQ ID NO:45), FBP20 (SEQ ID NO:49), FBP21 (SEQ ID NO:51), FBP22 (SEQ ID NO:53), FBP23 (SEQ ID NO:55), and FBP25 (SEQ ID NO:59) were not
10 previously assembled and represent novel nucleic acid molecules. The five remaining sequences, FBP9 (SEQ ID NO:27), FBP10 (SEQ ID NO:29), FBP16 (SEQ ID NO:41), FBP19 (SEQ ID NO:47), and FBP24 (SEQ ID NO:57) were previously assembled and disclosed in the database, but were not previously recognized as F-box proteins.

- Computer analysis of human FBPs revealed several interesting features (see
15 the schematic representation of FBPs in Figure 2. Three FBPs contain WD-40 domains; seven FBPs contain LRRs, and six FBPs contain other potential protein-protein interaction modules not yet identified in FBPs, such as leucine zippers, ring fingers, helix-loop-helix domains, proline rich motifs and SH2 domains.

- As examples of the human FBP family, a more detailed characterization of
20 some FBPs was performed. To confirm the specificity of interaction between the novel FBPs and human Skp1, eight in vitro translated FBPs were tested for binding to His-tagged-Skp1 pre-bound to Nickel-agarose beads. As a control Elongin C was used, the only known human Skp1 homolog. All 7 FBPs were able to bind His-Skp1 beads but not to His-tagged-Elongin C beads (Figure 29). The small amount of FBPs that bound to His-tagged-Elongin
25 C beads very likely represents non-specific binding since it was also present when a non-relevant protein (His-tagged-p27) bound to Nickel-agarose beads was used in pull-down assays (see as an example, Figure 29, lane 12).

- F-box deletion mutants, (Δ F)FBP1, (Δ F)FBP2, (Δ F)FBP3a, and mutants containing single point mutations in conserved amino acid residues of the F-box,
30 FBP3a(L51A) and FBP3a(W76A) were constructed. Mutants lacking the F-box and those with point mutations lost their ability to bind Skp1 (Figure 29), confirming that human FBPs require the integrity of their F-box to specifically bind Skp1.

- In order to determine whether FBP1, FBP2, FBP3a, FBP4 and FBP7 interact with human Skp1 and Cul1 in vivo (as Skp2 is known to do), flag-tagged-FBP1, -
35 Δ (F)FBP1, -FBP2, Δ (F)FBP2, -FBP3a, Δ (F)FBP3a, -FBP4 and -FBP7 were expressed in

HeLa cells from which cell extracts were made and subjected to immunoprecipitation with an anti-Flag antibody. As detected in immunoblots with specific antibodies to Cul1, Cul2 (another human cullin), and Skp1, the anti-Flag antibody co-precipitated Cul1 and Skp1, but not Cul2, exclusively in extracts from cells expressing wild-type FBPs (Figure 29 and data
5 not shown). These data indicate that as in yeast, the human Skp1/cullin complex forms a scaffold for many FBPs.

The binding of FBPs to the Skp1/Cul1 complex is consistent with the possibility that FBPs associate with a ubiquitin ligation activity. To test this possibility, Flag-tagged were expressed in HeLa cells, FBPs together with human Skp1 and Cul1.
10 Extracts were subjected to immunoprecipitation with an anti-Flag antibody and assayed for ubiquitin ligase activity in the presence of the human ubiquitin-activating enzyme (E1) and a human Ubc. All of the wild type FBPs tested, but not FBP mutants, associated with a ubiquitin ligase activity which produced a high molecular weight smear characteristic of ubiquitinated proteins (Figure 30). The ligase activity was N-ethylmaleimide (NEM)
15 sensitive (Figure 30, lane 2) and required the presence of both Ubc4 and E1. Results similar to those with Ubc4 were obtained using human Ubc3, whereas Ubc2 was unable to sustain the ubiquitin ligase activity of these SCFs (Figure 30, lanes 12, 13).

Using indirect immunofluorescence techniques, the subcellular distribution of FBP1, FBP2, FBP3a, FBP4 and FBP7 was studied in human cells. Flag-tagged-versions
20 of these proteins were expressed in HeLa, U2OS, and 293T cells and subjected to immunofluorescent staining with an anti-Flag antibody. FBP1, FBP4 and FBP7 were found to be distributed both in the cytoplasm and in the nucleus, while FBP2 was detected mainly in the cytoplasm and FBP3a mainly in the nucleus. Figure 32 shows, as an example, the subcellular localization of FBP1, FBP2, FBP3a, FBP4 observed in HeLa cells. The
25 localization of (Δ F)FBP1, (Δ F)FBP2, (Δ F)FBP3a mutants was identical to those of the respective wild-type proteins (Figure 32) demonstrating that the F-box and the F-box-dependent binding to Skp1 do not determine the subcellular localization of FBPs. Immunofluorescence stainings were in agreement with the results of biochemical subcellular fractionation.

30

6.2.2 Northern Blot Analysis of Novel Ubiquitin Ligase Gene Transcripts

RNA blot analysis was performed on poly(A)+ mRNA from multiple normal human tissues (heart, brain, placenta, lung, liver, skeletal, muscle, kidney, pancreas, spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocytes, see
35 Figure 33). FBP1 mRNA transcripts (a major band of ~7-kb and two minor bands of ~3.5 -

and ~2.5 kb) were expressed in all of the 16 human tissues tested but were more prevalent in brain and testis. Testis was the only tissue expressing the smaller FBP1 mRNA forms in amounts equal to, if not in excess of, the 7 kb form. FBP2 transcripts (~7.7-kb and ~2.4-kb) were expressed in all tissues tested, yet the ratio of the FBP2 transcripts displayed some tissue differences. An approximately 4 kb FBP3a transcript was present in all tissues tested and two minor FBP3a forms of approximately 3 kb and 2 kb became visible, upon longer exposure, especially in the testis. An approximately 4.8 kb FBP4 transcript was expressed in all normal human tissues tested, but was particularly abundant in heart and pancreas. Finally, the pattern of expression of the new FBPs was compared to that of FBP1 whose mRNA species (a major band ~4 kb and a minor band of ~8.5 kb) were found in all tissues but was particularly abundant in placenta.

6.2.3 Chromosomal Localization Of The Human FBP Genes

Unchecked degradation of cellular regulatory proteins (*e.g.*, p53, p27, β -catenin) has been observed in certain tumors, suggesting the hypothesis that deregulated ubiquitin ligases play a role in this altered degradation (reviewed in A. Ciechanover, 1998, *Embo J*, 17: 7151). A well understood example is that of MDM2, a proto-oncogene encoding a ubiquitin ligase whose overexpression destabilize its substrate, the tumor suppressor p53 (reviewed by Brown and Pagano, 1997, *Biochim Biophys Acta*, 1332: 1, 1998). To map the chromosomal localization of the human FBP genes and to determine if these positions coincided with loci known to be altered in tumors or in inherited disease, fluorescence in situ hybridization (FISH) was used. The FBP1 gene was mapped and localized to 10q24 (Fig. 34A), FBP2 to 9q34 (Figure 34B), FBP3a to 13q22 (Figure 34C), FBP4 to 5p12 (Figure 34D) and FBP5 to 6q25-26 (Figure 34E). FBP genes (particularly FBP1, FBP3a, and FBP5) are localized to chromosomal loci frequently altered in tumors (for references and details see Online Mendelian Inheritance in Man database, <http://www3.ncbi.nlm.nih.gov/omim/>). In particular, loss of 10q24 (where FBP1 is located) has been demonstrated in approx. 10 % of human prostate tumors and small cell lung carcinomas (SCLC), suggesting the presence of a tumor suppressor gene at this location. In addition, up to 7% of childhood acute T-cell leukemia is accompanied by a translocation involving 10q24 as a breakpoint, either t(10;14)(q24;q11) or t(7;10)(q35;q24). Although rarely, the 9q34 region (where FBP2 is located) has been shown to be a site of loss of heterozygosity (LOH) in human ovarian and bladder cancers. LOH is also observed in the region. Finally, 6q25-26 (where FBP5 is located) has been shown to be a site of loss of

heterozygosity in human ovarian, breast and gastric cancers hepatocarcinomas, Burkitt's lymphomas, and parathyroid adenomas.

7. EXAMPLE: FBP1 REGULATES THE STABILITY OF β -CATENIN

5 Deregulation of β -catenin proteolysis is associated with malignant transformation. *Xenopus* Slimb and *Drosophila* FBP1 negatively regulate the Wnt/ β -catenin signaling pathway (Jiang and Struhl, 1998, *supra*; Marikawa and Elinson, 1998). Since ubiquitin ligase complexes physically associate with their substrates, the studies in this Example were designed to determine whether FBP1 can interact with β -catenin. The
10 results show that FBP1 forms a novel ubiquitin ligase complex that regulates the in vivo stability of β -catenin. Thus, the identification of FBP1 as a component of the novel ubiquitin ligase complex that ubiquitinates β -catenin, provides new targets that can be used in screens for agonists, antagonists, ligands, and novel substrates using the methods of the present invention. Molecules identified by these assays are potentially useful drugs as
15 therapeutic agents against cancer and proliferative disorders.

7.1 MATERIALS AND METHODS FOR IDENTIFICATION OF FBP1 FUNCTION

Recombinant proteins, Construction of F-box mutants, Antibodies, Transient transfections,
20 Immunoprecipitation, Immunoblotting, Cell culture and Extract preparation Details of the methods are described in Section 6.1, *supra*.

7.2 RESULTS

25 7.2.1 Human FBP1 Interacts With β -Catenin

Flag-tagged FBP1 and β -catenin viruses were used to co-infect insect cells, and extracts were analyzed by immunoprecipitation followed by immunoblotting. β -catenin was co-immunoprecipitated by an anti-Flag antibody (Figure 35A), indicating that in intact
30 cells β -catenin and FBP1 physically interact. It has been shown that binding of the yeast FBP Cdc4 to its substrate Sic1 is stabilized by the presence of Skp1 (Skowyra et al., 1997, Cell, 91, 209-219). Simultaneous expression of human Skp1 had no effect on the strength of the interaction between FBP1 and β -catenin. To test the specificity of the FBP1/ β -catenin interaction, cells were co-infected with human cyclin D1 and FBP1 viruses. The
35 choice of this cyclin was dictated by the fact that human cyclin D1 can form a complex with the Skp2 ubiquitin ligase complex (Skp1-Cul1-Skp2; Yu et al., 1998, Proc. Natl. Acad. Sci.

U.S.A, 95:11324-9). Under the same conditions used to demonstrate the formation of the FBP1/ β -catenin complex, cyclin D1 could not be co-immunoprecipitated with Flag-tagged FBP1, and anti-cyclin D1 antibodies were unable to co-immunoprecipitate FBP1 (Figure 35B, lanes 1-3). Co-expression of Skp1 (Figure 35B, lanes 4-6) or Cdk4 with FBP1 and cyclin D1 did not stimulate the association of cyclin D1 with FBP1.

Mammalian expression plasmids carrying HA-tagged β -catenin and Flag-tagged FBP1 (wild type or mutant) were then co-transfected in human 293 cells. β -catenin was detected in anti-Flag immunoprecipitates when co-expressed with either wild type or (Δ F)FBP1 mutant (Figure 35C, lanes 4-6), confirming the presence of a complex formed between β -catenin and FBP1 in human cells.

7.2.2 F-box Deleted FBP1 Mutant Stabilizes β -Catenin In Vivo

The association of (Δ F)FBP1 to β -catenin suggested that (Δ F)FBP1 might act as a dominant negative mutant in vivo by being unable to bind Skp1/Cul1 complex, on the one hand, while retaining the ability to bind β -catenin, on the other. HA-tagged β -catenin was co-expressed together with Flag-tagged (Δ F)FBP1 or with another F-box deleted FBP, (Δ F)FBP2. FBP2 was also obtained with our screening for Skp1-interactors; and, like FBP1, contains several WD-40 domains. The presence of (Δ F)FBP1 specifically led to the accumulation of higher quantities of β -catenin (Figure 36A). To determine whether this accumulation was due to an increase in β -catenin stability, we measured the half-life of β -catenin using pulse chase analysis. Human 293 cells were transfected with HA-tagged β -catenin alone or in combination with the wild type or mutant FBP1. While wild type Fbp1 had little effect on the degradation of β -catenin, the F-box deletion mutant prolonged the half life of β -catenin from 1 to 4 hours (Figure 36B).

FBP1 is also involved in CD4 degradation induced by the HIV-1 Vpu protein (Margottin et al., *supra*). It has been shown that Vpu recruits FBP1 to DC4 and (Δ F) FBP1 inhibits Vpu-mediated CD4 regulation. In addition, FBP1-ubiquitin ligase complex also controls the stability of IKB α (Yaron et al., 1998, Nature, 396: 590). Thus, the interactions between FBP1 and β -catenin, Vpu protein, CD4, and IKB α are potential targets that can be used to screen for agonists, antagonists, ligands, and novel substrates using the methods of the present invention.

8. EXAMPLE: METHODS FOR IDENTIFYING p27 AS A SUBSTRATE OF THE FBP Skp2

Degradation of the mammalian G1 cyclin-dependent kinase (Cdk) inhibitor p27 is required for the cellular transition from quiescence to the proliferative state. The ubiquitination and degradation of p27 depend upon its phosphorylation by cyclin/Cdk complexes. Skp2, an F-box protein essential for entry into S phase, specifically recognizes p27 in a phosphorylation-dependent manner. Furthermore, both in vivo and in vitro, Skp2 is a rate-limiting component of the machinery that ubiquitinates and degrades phosphorylated p27. Thus, p27 degradation is subject to dual control by the accumulation of both Skp2 and cyclins following mitogenic stimulation.

This Example discloses novel assays that have been used to identify the interaction of Skp2 and p27 in vitro. First, an in vitro ubiquitination assay performed using p27 as a substrate is described. Second, Skp2 is depleted from cell extracts using anti-Skp2 antibody, and the effect on p27 ubiquitin ligase activity is assayed. Purified Skp2 is added back to such immunodepleted extracts to restore p27 ubiquitination and degradation. Also disclosed is the use of a dominant negative mutant, (Δ F)Skp2, which interferes with p27 ubiquitination and degradation.

The assays described herein can be used to test for compounds that inhibit cell proliferation. The assays can be carried out in the presence or absence of molecules, compounds, peptides, or other agents described in Section 5.5. Agents that either enhance or inhibit the interactions or the ubiquitination activity can be identified by an increase or decrease the formation of a final product are identified. Such agents can be used, for example, to inhibit Skp2-regulated p27 ubiquitination and degradation in vivo. Molecules identified by these assays are potentially useful drugs as therapeutic agents against cancer and proliferative disorders.

Dominant negative mutants, for example the mutant (Δ F)Skp2, and antisense oligos targeting SKP2, mRNA interfere with p27 ubiquitination and degradation, and can be used in gene therapies against cancer. The assays described herein can also be used to identify novel substrates of the novel FBP proteins, as well as modulators of novel ubiquitin ligase complex - substrate interactions and activities.

8.1 MATERIALS AND METHODS FOR IDENTIFICATION OF p27 AS A Skp2 SUBSTRATE

Protein extraction for in vitro ubiquitination assay Approx. 4 ml of HeLa S3 cell pellet were suspended in 6 ml of ice-cold buffer consisting of 20 mM Tris-HCl (pH 7.2), 2 mM

DTT, 0.25 mM EDTA, 10 μ g/ml leupeptin, and 10 μ g/ml pepstatin. The suspension was transferred to a cell nitrogen-disruption bomb (Parr, Moline, IL, cat #4639) that had been rinsed thoroughly and chilled on ice before use. The bomb chamber was connected to a nitrogen tank and the pressure was brought slowly to 1000 psi. The chamber was left on ice
5 under the same pressure for 30 minutes and then the pressure was released slowly. The material was transferred to an Eppendorf tube and centrifuged in a microcentrifuge at 10,000 g for 10 minutes. The supernatant (S-10) was divided into smaller samples and frozen at -80°C. This method of extract preparation based on the use of a cell nitrogen-disruption bomb extract preserves the activity to in vitro ubiquitinate p27 better than the
10 method previously described (Pagano et al., 1995, Science 269:682-685).

Reagents and antibodies Ubiquitin aldehyde (Hershko & Rose, 1987, Proc. Natl. Acad. Sci. USA 84:1829-33), methyl-ubiquitin (Hershko & Heller, 1985, Biochem. Biophys. Res. Commun. 128:1079-86) and p13 beads (Brizuela et al., 1987, EMBO J. 6:3507-3514) were
15 prepared as described. β , γ -imidoadenosine-50-triphosphate (AMP-PNP), staurosporine, hexokinase, and deoxy-glucose were from Sigma; lovastatin obtained from Merck; flavopiridol obtained from Hoechst Marion Roussel. The phospho-site p27 specific antibody was generated in collaboration with Zymed Inc. by injecting rabbits with the phospho-peptide NAGSVEQT*PKKPGLRRRQT (SEQ ID NO: 85), corresponding to the
20 carboxy terminus of the human p27 with a phosphothreonine at position 187 (T*). The antibody was then purified from serum with two rounds of affinity chromatography using both phospho- and nonphospho-peptide chromatography. All the other antibodies are described in Section 6.1.

Immunodepletion Assays For immunodepletion assays, 3 μ l of an Skp2 antiserum was adsorbed to 15 μ l Affi-Prep Protein-A beads (BioRad), at 4°C for 90 min. The beads were washed and then mixed (4°C, 2 hours) with 40 μ l of HeLa extract (approximately 400 μ g of protein). Beads were removed by centrifugation and supernatants were filtered through a 0.45- μ Microspin filter (Millipore). Immunoprecipitations and immunoblots were
25 performed as described (M. Pagano, et al., 1995, *supra*. Rabbit polyclonal antibody against purified GST-Skp2 was generated, affinity-purified (AP) and characterized as described (M. Pagano, in Cell Cycle-Materials and Methods, M. Pagano Ed. (Springer, NY, 1995), chap. 24; E. Harlow and D. Lane, in Using antibodies. A Laboratory Manual (Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, 1998), in collaboration with Zymed Inc. (cat # 51-
30 1900). Monoclonal antibodies (Mabs) to human Cull1, and cyclin E, (Faha et al., 1993, J. of
35 1900).

Virology 67:2456); AP rabbit antibodies to human p27, Skp1 (Latres et al., 1999, Oncogene 18:849), Cdk2 (Pagano, et al., 1992, Science 255:1144) and phospho-site p27 specific antibody. Mab to cyclin B was from Santa Cruz (cat # sc-245); Mabs to p21 (cat # C24420) and p27 (cat # K25020) Transduction lab; anti-Flag rabbit antibody from Zymed (cat # 71-5400). An AP goat antibody to an N-terminal Skp2 peptide (Santa Cruz, cat # sc-1567) was used.

Construction of Skp2 F-box mutant (Δ F)Skp2 mutant was obtained by removing a DNA fragment (nucleotides 338-997) with BspEI and XbaI restriction enzymes, and replacing it with a PCR fragment containing nucleotides 457 to 997. The final construct encoded a protein lacking residues 113-152.

Recombinant proteins cDNA fragments encoding the following human proteins: Flag-tagged FBP1, Flag-tagged (Δ F)FBP1, Flag-tagged FBP3a, Skp2, HA-tagged Cul1, HA-tagged Cul2, β -catenin, His-tagged cyclin D1, Skp1, His-tagged Skp1, His-tagged Elongin C were inserted into the baculovirus expression vector pBacpak-8 (Clontech) and cotransfected into Sf9 cells with linearized baculovirus DNA using the BaculoGold transfection kit (Pharmingen). Baculoviruses expressing human His-tagged cyclin E and HA-tagged Cdk2 were supplied by D. Morgan (Desai, 1992, Molecular Biology of the Cell 3: 571). Recombinant viruses were used to infect 5B cells and assayed for expression of their encoded protein by immunoblotting as described above. His-proteins were purified with Nickel-agarose (Invitrogen) according to the manufacturer's instructions. The different complexes were formed by co-expression of the appropriate baculoviruses and purified by nickel-agarose chromatography, using the His tag at the 5' of Skp1 and cyclin E. Unless otherwise stated, recombinant proteins were added to incubations at the following amounts: cyclin E/Cdk2, ~0.5 pmol; Skp1, ~0.5 pmol; Skp2, ~0.1 pmol; FBP1, ~0.1 pmol; FBP3a, ~0.1 pmol, Cul1, ~0.1 pmol. The molar ratio of Skp1/Skp2, Skp1/FBP1, Skp1/FBP3a, and Skp1/Cul1 in the purified preparations was ~5.

Extract preparation and cell synchronization, Transient transfections, Immunoprecipitation and Immunoblotting Methods were carried out as described in Section 6.1, *supra*.

8.2 RESULTS

8.2.1 p27 in vitro ubiquitination assay

In an exemplary in vitro ubiquitination assay, logarithmically growing, HeLa-S3 cells were collected at a density of 6×10^5 cells/ml. Cells are arrested in G1 by 48-hour treatment with 70 μ M lovastatin as described (O'Connor & Jackman, 1995 in Cell Cycle-Materials and Methods, M. Pagano, ed., Springer, NY, chap. 6). 1 μ l of in vitro translated [35S]p27 is incubated at 30°C for different times (0 - 75 minutes) in 10 μ l of ubiquitination mix containing: 40 mM Tris pH 7.6, 5 mM $MgCl_2$, 1 mM DTT, 10 % glycerol, 1 μ M ubiquitin aldehyde, 1 mg/ml methyl ubiquitin, 10 mM creatine phosphate, 0.1 mg/ml creatine phosphokinase, 0.5 mM ATP, 1 μ M okadaic acid, 20-30 μ g HeLa cell extract. Ubiquitin aldehyde can be added to the ubiquitination reaction to inhibit the isopeptidases that would remove the chains of ubiquitin from p27. Addition of methyl ubiquitin competes with the ubiquitin present in the cellular extracts and terminates p27 ubiquitin chains. Such chains appear as discrete bands instead of a high molecular smear. These shorter polyubiquitin chains have lower affinity for the proteasome and therefore are more stable. Reactions are terminated with Laemmli sample buffer containing β -mercaptoethanol and the products can be analyzed on protein gels under denaturing conditions.

Polyubiquitinated p27 forms are identified by autoradiography. p27 degradation assay is performed in a similar manner, except that (i) Methylated ubiquitin and ubiquitin aldehyde were omitted; (ii) The concentration of HeLa extract is approximately 7 μ g/ μ l; (iii) Extracts are prepared by hypotonic lysis (Pagano et al., 1995, Science 269:682), which preserves proteasome activity better than the nitrogen bomb disruption procedure. In the absence of methyl ubiquitin, p27 degradation activity, instead of p27 ubiquitination activity, can be measured.

The samples are immunoprecipitated with an antibody to p27 followed by a subsequent immunoprecipitation with an anti-ubiquitin antibody and run on an 8% SDS gel. The high molecular species as determined by this assay are ubiquitinated. As a control, a p27 mutant lacking all 13 lysines was used. This mutant form of p27 is not ubiquitinated and runs at higher molecular weight on the 8% SDS gel.

8.2.2 p27-Skp2 interaction assays and p27-Skp2 immunodepletion assay

The recruitment of specific substrates by yeast and human FBPs to Skp1/cullin complexes is phosphorylation-dependent. Accordingly, peptides derived from I κ B α and β -catenin bind to FBP1 specifically and in a phosphorylation-dependent manner

(Yaron, 1998, *Nature* 396: 590; Winston et al., 1999, *Genes Dev.* 13: 270). A p27 phosphopeptide with a phosphothreonine at position 187 was assayed for its ability to bind to human FBPs, including Skp2 and the FBP1, FBP2, FBP3a, FBP4, FBP5, FBP6, and FBP7, isolated by using a 2-hybrid screen using Skp1 as bait, as described in Section 6, above. Four of
5 these FBPs contain potential substrate interaction domains, such as WD-40 domains in FBP1 and FBP2, and leucine-rich repeats in Skp2 and FBP3a. The phospho-p27 peptide was immobilized to Sepharose beads and incubated with these seven in vitro translated FBPs (Figure 37A). Only one FBP, Skp2, was able to bind to the phospho-T187 p27
10 peptide. Then, beads linked to p27 peptides (in either phosphorylated or unphosphorylated forms) or with an unrelated phospho-peptide were incubated with HeLa cell extracts. Proteins stably associated with the beads were examined by immunoblotting. Skp2 and its associated proteins, Skp1 and Cull1, were readily detected as proteins bound to the phospho-p27 peptide but not to control peptides (Figure 37B).

To further study p27 association to Skp2, in vitro translated p27 was
15 incubated with either Skp1/Skp2 complex, cyclin E/Cdk2 complex, or the combination of both complexes under conditions in which p27 is phosphorylated on T187 by cyclin E/Cdk2 (Montagnoli, A., et al., 1999, *Genes & Dev* 13: 1181). Samples were then immunoprecipitated with an anti-Skp2 antibody. p27 was co-immunoprecipitated with Skp2 only in the presence of cyclin E/Cdk2 complex (Fig. 37C). Notably, under the same
20 conditions, a T187-to-alanine p27 mutant, p27(T187A), was not co-immunoprecipitated by the anti-Skp2 antibody. Finally, we tested Skp2 and p27 association in vivo. Extracts from HeLa cells and IMR90 human diploid fibroblasts were subjected to immunoprecipitation with two different antibodies to Skp2 and then immunoblotted. p27 and Cull1, but not cyclin D1 and cyclin B1, were specifically detected in Skp2 immunoprecipitates (Fig. 38).
25 Importantly, using a phospho-T187 site p27 specific antibody we demonstrated that the Skp2-bound p27 was phosphorylated on T187 (Fig. 38, lane 2, bottom panel). Furthermore, an anti-peptide p27 antibody specifically co-immunoprecipitated Skp2. These results indicate that the stable interaction of p27 with Skp2 was highly specific and dependent upon phosphorylation of p27 on T187.

30 A cell-free assay for p27 ubiquitination which faithfully reproduced the cell cycle stage-specific ubiquitination and degradation of p27 has been developed (Montagnoli et al., supra). Using this assay, a p27-ubiquitin ligation activity is higher in extracts from asynchronously growing cells than in those from G1-arrested cells (Figure 39A, lanes 2 and 4). In accordance with previous findings (Montagnoli, A., et al., supra), the addition of
35 cyclin E/Cdk2 stimulated the ubiquitination of p27 in both types of extracts (Figure 39A,

lanes 3 and 5). However, this stimulation was much lower in extracts from G1-arrested cells than in those from growing cells, suggesting that in addition to cyclin E/Cdk2, some other component of the p27-ubiquitin ligation system is rate-limiting in G1. This component could be Skp2 since, in contrast to other SCF subunits, its levels are lower in
5 extracts from G1 cells than in those from asynchronous cells and are inversely correlated with levels of p27 (Figures 39B and 43). Skp2 was thus tested to determine if it is a rate-limiting component of a p27 ubiquitin ligase activity. The addition of recombinant purified Skp1/Skp2 complex alone to G1 extracts did not stimulate p27 ubiquitination significantly (Figure 39A, lane 6). In contrast, the combined addition of Skp1/Skp2 and cyclin E/Cdk2
10 complexes strongly stimulated p27 ubiquitination in G1 extracts (Figure 39A, lane 7). Similarly, the combined addition of Skp1/Skp2 and cyclin E/Cdk2 strongly stimulated p27 proteolysis as measured by a degradation assay (Figure 39A, lanes 13-16). Since the Skp1/Skp2 complex used for these experiments was isolated from insect cells co-expressing baculovirus His-tagged-Skp1 and Skp2 (and co-purified by nickel-agarose
15 chromatography), it was possible that an insect-derived F-box protein co-purified with His-Skp1 and was responsible for the stimulation of p27 ubiquitination in G1 extracts. This possibility was eliminated by showing that the addition of a similar amount of His-tagged-Skp1, expressed in the absence of Skp2 in insect cells and purified by the same procedure, did not stimulate p27 ubiquitination in the presence of cyclin E/Cdk2 (Figure 39A, lane 8).
20 Furthermore, we found that neither FBP1 nor FBP3a could replace Skp2 for the stimulation of p27-ubiquitin ligation in G1 extracts (Figure 39A, lanes 9-12). Stimulation of p27-ubiquitination in G1 extracts by the combined addition of Skp1/Skp2 and cyclin E/Cdk2 could be observed only with wild-type p27, but not with the p27(T187A) mutant (lanes 17-20), indicating that phosphorylation of p27 on T187 is required for the Skp2-mediated
25 ubiquitination of p27. These findings indicated that both cyclin E/Cdk2 and Skp1/Skp2 complexes are rate-limiting for p27 ubiquitination and degradation in the G1 phase.

To further investigate the requirement of Skp2 for p27 ubiquitin ligation, Skp2 was specifically removed from extracts of asynchronously growing cells by immunodepletion with an antibody to Skp2. The immunodepletion procedure efficiently
30 removed most of Skp2 from these extracts and caused a drastic reduction of p27-ubiquitin ligation activity (Figure 40A, lane 4) as well as of p27 degradation activity. This effect was specific as shown by the following observations: (i) Similar treatment with pre-immune serum did not inhibit p27-ubiquitination (Figure 40A, lane 3); (ii) Pre-incubation of anti-Skp2 antibody with recombinant GST-Skp2 (lane 5), but not with a control protein (lane 4),
35 prevented the immunodepletion of p27-ubiquitination activity from extracts; (iii) p27-

ubiquitinating activity could be restored in Skp2-depleted extracts by the addition of His-Skp1/Skp2 complex (Figure 40B, lane 3) but not His-Skp1 (lane 2), His-Skp1/Cul1 complex (lane 4), or His-Skp1/FBP1.

We then immunoprecipitated Skp2 from HeLa extracts and tested whether this immunoprecipitate contained a p27 ubiquitinating activity. The anti-Skp2 beads, but not a immunoprecipitate made with a pre-immune (PI) serum, was able to induce p27 ubiquitination in the presence of cyclin E/Cdk2 (Figure 40C, lanes 2 and 3). The addition of purified recombinant E1 ubiquitin-activating enzyme, and purified recombinant Ubc3 did not greatly increase the ability of the Skp2 immunoprecipitate to sustain p27 ubiquitination, (Figure 40C, lane 5), likely due to the presence of both proteins in the rabbit reticulocyte lysate used for p27 in vitro translation.

8.2.3 F-BOX deleted SKP2mutant stabilizes p27 in vivo

Skp2 also targets p27 for ubiquitin-mediated degradation in vivo. The F-box-deleted FBP1 mutant, (Δ F)FBP1, acts in vivo as a dominant negative mutant, most likely because without the F-box is unable to bind Skp1/Cul1 complex but retains the ability to bind its substrates. Therefore, once expressed in cells, (Δ F)Fb sequesters β -catenin and IKB α and causes their stabilization. An F-box deleted Skp2 mutant, (Δ F)Skp2, was constructed. p27 was expressed in murine cells either alone or in combination with (Δ F)Skp2 or (Δ F)FBP1 (see Figure 41). The presence of (Δ F)Skp2 led to the accumulation of higher quantities of p27. To determine whether this accumulation was due to an increase in p27 stability, the half-life of p27 was measured using pulse chase analysis (for details, see Section 8, above). Indeed, (Δ F)Skp2 prolonged p27 half-life from less than 1 hour to ~3 hours. Since in these experiments the efficiency of transfection was approximately 10%, (Δ F)Skp2 affected only the stability of co-expressed human exogenous p27, but not of murine endogenous p27.

8.2.4 SKP2 ANTISENSE EXPERIMENTS

SKP2 mRNA was targeted with antisense oligonucleotides to determine whether a decrease in Skp2 levels would influence the abundance of endogenous p27. Two different antisense oligos, but not control oligodeoxynucleotides induced a decrease in Skp2 protein levels (Figure 42). Concomitant with the Skp2 decrease, there was a substantial increase in the level of endogenous p27 protein. Similar results were obtained with cells blocked at the G1/S transition with hydroxyurea or aphidicolin treatment (lanes 9-16).

Thus, the effect of the SKP2 antisense oligos on p27 was not a secondary consequence of a possible block in G1 due to the decrease in Skp2 levels.

Antisense experiments were performed as described in (Yu, 1998, Proc. Natl. Acad. Sci. U. S. A. 95: 11324). Briefly, four oligodeoxynucleotides that contain a
5 phosphorothioate backbone and C-5 propyne pyrimidines were synthesized (Keck
Biotechnology Resource Laboratory at Yale University): (1) 5'-
CCTGGGGGATGTTCTCA-3' (SEQ ID NO: 86) (the antisense direction of human Skp2
cDNA nucleotides 180-196); (2) 5'-GGCTTCCGGGCATTAG-3' (SEQ ID NO: 87) [the
scrambled control of (1)]; (3) 5'-CATCTGGCAGATTCCA-3' (SEQ ID NO: 88) (the
10 antisense direction of Skp2 cDNA nucleotides 1137-1153); (4) 5'-
CCGCTCATCGTATGACA-3' (89) [the scrambled control for (3)]. The oligonucleotides
were delivered into HeLa cells using Cytofectin GS (Glen Research) according to the
manufacturers instructions. The cells were then harvested between 16 and 18 hours
posttransfection.

15

9. EXAMPLE: METHOD FOR IDENTIFYING Cks1 AS A MEDIATOR OF THE FBP Skp2/p27 INTERACTION

As stated in Example 8, p27 is recognized by Skp2 in a phosphorylation-
dependent manner for entry into S phase and Skp2 is a rate-limiting component of the
20 machinery that ubiquitinates and degrades phosphorylated p27. This Example discloses
novel assays that have been used to identify the interactions of Cks1 with Skp2 and Cks1
with p27 in vitro and in a purified system. First, extracts of HeLa cells are fractionated and
the activity of the fractions to promote the ligation of p27 is tested. Second, identification
of Cks1 as the factor required for p27-ubiquitin ligation is confirmed with use of
25 recombinant Cks1. Third, identification of Cks1's involvement in the p27-ubiquitin ligation
after p27 is phosphorylated. Fourth, Cks1 increases the binding of Skp2 to p27. Fifth, Cks1
binds to Skp2. Sixth, Cks1 binds to the C-terminus of p27.

The assays described herein can be used to test for compounds that inhibit
cell proliferation. The assays can be carried out in the presence or absence of molecules,
30 compounds, peptides, or other agents described in Section 5.5. Agents that either enhance
or inhibit the interactions or the ubiquitination activity can be identified by an increase or
decrease the formation of a final product are identified. Such agents can be used, for
example, to inhibit Skp2-regulated p27 ubiquitination and degradation in vivo. Molecules
identified by these assays are potentially useful drugs as therapeutic agents against cancer
35 and proliferative disorders.

- Dominant negative mutants and antisense mRNA, oligos targeting the gene for Cks1, interfere with p27 ubiquitination and degradation, and can be used in gene therapies against cancer. The assays described herein can also be used to identify additional novel substrates of the novel FBP proteins, as well as additional modulators of novel
- 5 ubiquitin ligase complex - substrate interactions and activities.

9.1 MATERIALS AND METHODS FOR IDENTIFYING Cks1 AS A MEDIATOR OF THE FBP Skp2/p27 INTERACTION

- Proteins His₆-tagged p27 and Cdc34 were expressed in *E. coli* and purified by nickel-
 10 agarose chromatography. Cks2 and p13^{Suc1} were expressed in bacteria and purified by gel filtration chromatography. His₆-Skp1/Skp2, His₆-Skp1/β-TrCP, His₆-cyclin E/Cdk2, and His₆-Cul-1/ROC1 were produced by co-infection of 5B insect cells with baculoviruses encoding the corresponding proteins and were purified by nickel-agarose chromatography as described previously (Montagnoli, et al., 1999, *Genes & Dev.* 13:1501; Carrano, et al.,
 15 1999, *Nat. Cell Biol.* 1:193). The approximate concentrations of recombinant proteins in these preparations were (in pmole/μl): Skp1, 5; Skp2, 0.5; Cul-1, 4; ROC1, 1; cyclin E, 8; Cdk2, 1.5. Purified recombinant human Nedd8 was the generous gift of C. Pickart, and purified recombinant human Cks1 was the generous gift of S. Reed. Purified GST-IκBα(1-154) and its constitutively active kinase IKKβ^{S177E,S181E} were generously provided by Z.-Q.
 20 Pan. ³⁵S-labeled p27, Skp2 and Cks proteins were prepared by *in vitro* transcription-translation, using the TnT Quick kit (Promega) and ³⁵S-methionine (Amersham).
Purification of Nedd8-conjugating enzymes Purified recombinant human Nedd8 was the generous gift of C. Pickart. A mixture of Nedd8-conjugating enzymes (E1-like APP-BP1-Uba3 heterodimer and E2-like Ubc12: Osaka, et al., 1998, *Genes Dev.* 12:2263; Gong, L.,
 25 Yeh, E.T., 1999, *J. Biol. Chem.* 274:12036) was co-purified from lysates of rabbit reticulocytes by a "covalent affinity" chromatography procedure similar to that used for the purification of E2s (Hershko, et al., 1983, *J. Biol. Chem.* 258:8206), except that unfractionated reticulocyte lysate was applied to a column of GST-Nedd8-Sepharose (5 mg/ml). Following a wash with 1M KCl, all proteins bound to immobilized Nedd8 by
 30 thiolester linkages were co-eluted with a solution containing 20 mM DTT. The DTT eluate was concentrated by ultrafiltration to approx. 1/10 of the original volume of reticulocyte lysate. This preparation had strong activity in the ligation of Nedd8 to Cul-1, without any detectable hydrolase activity that removes Nedd8 from Cul-1.
Purification of the factor required for p27-ubiquitin ligation A frozen pellet from 50g
 35 of HeLa S3 cells (National Cell Culture Center) was disrupted by a nitrogen cell disruption

bomb (Parr, Moline, IL) as described Montagnoli, et al., 1999, Genes & Dev. 13:1181, except that the buffer also contained 10 µg/ml chymostatin and 5 µg/ml aprotinin. The extract was centrifuged at 15,000xg for 20 min and the supernatants were centrifuged again at 100,000xg for 60 min. The supernatant was subjected to fractionation on DEAE-cellulose as described (Hershko, et al., 1983, J. Biol. Chem. 258:8206), except that 2,500 mg of protein was loaded on 250 ml of resin. The fraction not adsorbed to the resin (Fraction 1) was collected and was concentrated by centrifuge ultrafiltration to approx. 10 mg/ml. Fraction 1 (100 mg of protein) was subjected to heat-treatment at 90°C for 10 minutes. The sample was allowed to stay on ice for 30 min, and then the precipitate was removed by centrifugation (10,000xg, 15 min). Approximately 99% of protein was removed by heat-treatment. The supernatant was concentrated by ultrafiltration and then was applied to a MonoS HR 5/5 column (Pharmacia) equilibrated with 50 mM Tris-HCl, 1 mM DTT and 0.1% (w/v) Brij-35 (Boehringer). The column was washed with 15 ml of the above buffer and was then eluted with a gradient of 0-200 mM NaCl. Activity in column fractions was followed by the p27-ubiquitin ligation assay in the presence of purified SCF^{Skp2} components (see below). The peak fractions of activity eluted at around 30-40 mM NaCl. The peak containing factor activity was pooled, concentrated by centrifuge ultrafiltration and was subjected to the final step of gel filtration chromatography on Superdex-75 HR 10/30 column (Pharmacia) equilibrated with 20 mM Tris-HCl (pH 7.2), 150 mM NaCl, 1 mM DTT and 0.1% Brij-35. Samples of 0.5 ml were collected at a flow rate of 0.4 ml/min. Column fractions were concentrated to a volume of 50 µl by centrifuge ultrafiltration (Centricon-10, Amicon). Samples of 0.004 µl of column fractions were assayed for activity to stimulate p27-ubiquitin ligation. Results were quantified by phosphorimager analysis and were expressed as the percentage of ³⁵S-p27 converted to ubiquitin conjugates. Arrows at top indicate the elution position of molecular mass marker proteins (kDa).

Mass spectrometric sequencing The 10-kDa protein from the last step of purification was excised and digested in gel as described (Shevchenko, et al., 1996, Anal. Chem. 68:850. Mass spectrometric analysis was performed on a Sciex QSTAR mass spectrometer (MDS-Sciex, Concord, ON, Canada). A tryptic peptide at mass 2163.5 was fragmented from doubly and triply charged species to yield a complete match to residues 5-20 of human Cks1.

Assay of p27-ubiquitin ligation. Unless otherwise stated, the reaction mixture contained in a volume of 10 µl: 40 mM Tris-HCl (pH 7.6), 5 mM MgCl₂, 1 mM DTT, 10% (v/v) glycerol, 10 mM phosphocreatine, 100 µg/ml creatine phosphokinase, 0.5 mM ATP, 1

mg/ml soybean trypsin inhibitor, 1 μ M ubiquitin aldehyde, 1 mg/ml methylated ubiquitin, 1 pmol E1, 50 pmol Cdc34, 0.25 μ l Skp2/Skp1, 0.25 μ l Cul-1/ROC1, 0.1 μ l cyclin E/Cdk2, 0.5 μ l of 35 S-p27 and additions as specified. Following incubation at 30°C for 60 minutes, samples were subjected to SDS-polyacrylamide gel electrophoresis and autoradiography.

- 5 The ligation of I κ B α to ubiquitin was assayed as described (Chen, et al., 2000, J. Biol. Chem. 275:15432), except that baculovirus-expressed, purified Skp1/ β -TrCP was used (5 pmol Skp1, \sim 1 pmol β -TrCP).

- Preparation of 32 P labeled purified p27 and assay of its ubiquitylation. Purified p27 (0.18 μ g) was incubated (60 minutes at 30 °C) with Cdk2/cyclin E (0.25 μ l) in a reaction mixture containing in a volume of 10 μ l: 50 mM Tris-HCl (pH 7.6), 5 mM MgCl₂, 1 mM DTT, 10% glycerol, 1 mg/ml soybean trypsin inhibitor, 1 μ M okadaic acid and 100 μ M [32 P- γ]-ATP (\sim 50 μ Ci). This preparation is referred to as " 32 P-p27". The ligation of p27 to MeUb was assayed as described above, with the following changes: 35 S-p27 was replaced by 32 P-p27, the concentration of unlabeled ATP was increased to 2 mM (for more complete isotopic dilution of labeled ATP present in the preparation of 32 P-p27) and okadaic acid (1 μ M) was added.

- Assay of binding of p27 to Skp2/Skp1 The reaction mixture contained in a volume of 10 μ l: 40 mM Tris-HCl (pH 7.6), 2 mg/ml bovine serum albumin , 1 μ l 35 S-p27, 1 μ l Cdk2/cyclin E, 1 μ l Skp2/Skp1, as well as MgCl₂, ATP, DTT, phosphocreatine and creatine phosphokinase at concentrations similar to those described above for p27-ubiquitin ligation assay. Following incubation at 30°C for 30 min, 6 μ l of Affi-prep-Protein A beads (BioRad) to which polyclonal rabbit antibody against full length Skp2 (Carrano, et al., 1999, Nat. Cell Biol. 1:193) had been covalently linked by dimethyl pimelimidate (Harlow, E. & Lane, D., 1998, in *Antibodies. A Laboratory Manual* (eds. Harlow, E. & Lane, D.), Cold Spring Harb. LabPress, Cold Spring Harbor, NY) was added. The samples were rotated with the anti-Skp2-Protein A beads at 4°C for 2 hours, and then the beads were washed 4 times with 1-ml portions of RIPA buffer (Harlow, E. & Lane, D., 1998, in *Antibodies. A Laboratory Manual* (eds. Harlow, E. & Lane, D.), Cold Spring Harb. LabPress, Cold Spring Harbor, NY). Following elution with SDS electrophoresis sample buffer, the samples were subjected to SDS-polyacrylamide gel electrophoresis and autoradiography.

9.2 RESULTS

9.2.1 The factor from Fraction 1 is a protein

- The activity of Fraction 1 is not destroyed by heating at 90°C. However, the active factor is a protein, as indicated by the observation that incubation of heat-treated

Fraction 1 with trypsin completely destroyed its activity (FIG. 44, lane 2). Heat-treated Fraction 1 (~ 0.1 mg/ml) was incubated at 37°C for 60 min with 50 mM Tris-HCl (pH 8.0) either in the absence (lane 1) or in the presence of 0.6 mg/ml of TPCK-treated trypsin (Sigma T8642) (lane 2). Trypsin action was terminated by the addition of 2 mg/ml of soybean trypsin inhibitor (STI). In lane 3, STI was added 5 min prior to a similar incubation with trypsin. Subsequently, samples corresponding to ~50 ng of heat-treated Fraction 1 were assayed for the stimulation of p27-ubiquitin ligation. Incubation of Fraction 1 with trypsin is terminated by the addition of excess soybean trypsin inhibitor (STI), to prevent proteolytic damage to the other components of the system, added following trypsin treatment. STI indeed efficiently blocks trypsin action as is shown in a control experiment in which STI is added to heated Fraction 1 prior to incubation with trypsin (FIG. 44, lane 3). In this incubation, there is no significant decrease in p27-ubiquitin ligation.

9.2.2 The factor from Fraction 1 is not Nedd8

Podust et al. (Podust, et al., 2000, Proc. Natl. Acad. Sci. U.S.A. 97:4579) have reported that the ligation of p27 to ubiquitin requires Fraction 1, and have suggested that Nedd8 is the active component in Fraction 1. Nedd8 (called Rub-1 in yeast) is a highly conserved ubiquitin-like protein that is ligated to different cullins, including Cul-1 (Yeh, et al., 2000, Gene 248:1). The ligation of Nedd8 to Cul-1 has been shown to stimulate, though not to be absolutely required for, the activity of the SCF^{B-TrCP} complex in the ligation of ubiquitin to IκBα (Furukawa, et al., 2000, Mol. Cell Biol. 20:8185; Read, et al., 2000, Mol. Cell Biol. 20:2326; Wu, et al., 2000, J. Biol. Chem 275:32317). Since ³⁵S-labeled p27 can be produced by *in vitro* translation in reticulocyte lysates, and since reticulocyte lysates contain the enzymes required for the ligation of Nedd8 to cullins (Osaka, et al., 1998, Genes Dev. 12:2549), it is possible that under these conditions Nedd8 could be ligated to Cul-1. However, recombinant purified Nedd8 does not replace the factor from Fraction 1 in promoting p27-ubiquitin ligation (FIG. 45A). Where indicated, ~50 ng of heat-treated Fraction 1 or 100 ng of purified recombinant human Nedd8 are added to the p27-MeUb ligation assay. To further examine this problem, the enzymes that ligate Nedd8 to Cul-1 are purified by affinity chromatography on GST-Nedd8-Sepharose. Incubation of Cul-1 with Nedd8 and its purified conjugating enzymes convert about one-half of Cul-1 molecules to Nedd8-conjugated form that migrates slower in SDS-polyacrylamide gel electrophoresis (FIG. 45B). Ligation of Nedd8 to Cul-1. Cul-1/ROC1 (3 μl) is incubated with Nedd8 (10 μg) and purified Nedd8-conjugating enzymes (20 μl) in a 100-μl reaction mixture containing Tris (pH 7.6), MgCl₂, ATP, phosphocreatine, creatine phosphokinase, DTT,

glycerol and STI at concentrations similar to those described for the p27-ubiquitin ligation assay. A control preparation of Cul1/ROC1 is incubated under similar conditions, but without Nedd8 conjugating enzymes. Following incubation at 30°C for 2 hours, samples of control or Nedd8-modified preparations are separated on an 8% polyacrylamide-SDS gel and immunoblotted with an anti-Cul-1 antibody (Zymed). The slower migrating form indeed contains Nedd8 as is verified by immunoblotting with a specific antibody directed against Nedd8. The activity of these preparations of Nedd8-conjugated and unmodified Cul-1 in the p27 ubiquitinylation reaction is measured in the presence or absence of heat-treated Fraction 1. Bacterially expressed, purified p27 (20 ng) is used as the substrate rather than ³⁵S-labeled p27 translated in reticulocyte lysate, because reticulocyte lysates also contain the enzyme(s) that rapidly cleave(s) the amide linkage between Nedd8 and Cul-1. The ligation of p27 to MeUb occurs at 30C for 60 minutes and is followed by separation on a 12.5% polyacrylamide-SDS gel, transfer to nitrocellulose, and immunoblotting with a monoclonal antibody directed against p27 (Transduction Laboratories). Using this purified system and in the presence of heat-treated Fraction 1, significant formation of mono-ubiquitinated, and less of di-ubiquitinated derivatives of p27 is promoted by unmodified Cul-1 (FIG. 45C). With the purified system, conjugates with MeUb larger than the di-ubiquitinated form are not observed, as opposed to the 4-5 conjugates observed with *in vitro*-translated ³⁵S-p27 (compare with Fig. 44). With Cul-1 conjugated to Nedd8, a modest stimulation in the ubiquitinylation of p27 is observed, with a special increase in the formation of the di-ubiquitin derivative (FIG. 45, lane 3). In different preparations of Cul-1, Nedd8 ligation increases the over-all rate of p27-ubiquitin ligation by 1.5-3 fold. The basal activity of p27-ubiquitin ligation observed with unmodified Cul-1 is not due to its significant modification by Nedd8 in insect cells, from which baculovirus-expressed Cul-1 was purified, because similar activity is observed with a mutant Cul-1 in which Lys720 at its specific Nedd8-ligation site (Yeh, et al., 2000, Gene 248:1) was changed to Arg. Other investigators have also observed that elimination of Nedd8 modification by a similar mutation significantly reduced, but did not abolish the activity of SFC^{8-TrCP} in the ubiquitinylation of IκBα (Furukawa, et al., 2000, Mol. Cell Biol. 20:8185; Read, et al., 2000, Mol. Cell Biol. 20:2326; Wu, et al., 2000, J. Biol. Chem 275:32317). Importantly, the supplementation of Fraction 1 is still required for p27-MeUb ligation even in the presence of Nedd8-modified Cul-1 (FIG. 45, lanes 5 and 6). Similar results are obtained when MeUb is replaced by native ubiquitin, except that in the latter case high molecular weight polyubiquitin derivatives of p27 are formed. Thus, the data does not support the conclusions of Podust et al. (Podust et al., 2000, Proc. Natl. Acad. Sci. U.S.A. 97:4579) that the active component in Fraction 1 is Nedd8.

9.2.3 Purification of the factor and its identification as Cks1

The factor from fraction 1 is purified. FIG. 46A shows the last step of purification on a gel filtration column. The peak of active material from the MonoS step was applied to a Superdex 75 HR 10/30 column (Pharmacia) equilibrated with 20 mM Tris-HCl (pH 7.2), 150 mM NaCl, 1 mM DTT and 01% Brij-35. Samples of 0.5 ml were collected at a flow rate of 0.4 ml/min. Column fractions were concentrated to a volume of 50 μ l by centrifuge ultrafiltration (Centricon-10, Amicon). Samples of 0.004 μ l of column fractions were assayed for activity to stimulate p27-ubiquitin ligation. Results were quantified by phosphorimager analysis and were expressed as the percentage of 35 S-p27 converted to ubiquitin conjugates. Arrows at top indicate the elution position of molecular mass marker proteins (kDa). Activity eluted as a sharp peak at an apparent molecular mass of approx. 10 kDa. Electrophoresis of samples of 2.5 μ l from the indicated fractions of the Superdex 75 column on a 16% polyacrylamide-SDS gel and silver staining of column fractions show a single protein of approx. 10 kDa (FIG. 46B). Numbers on the right indicate the migration position of molecular mass marker proteins (kDa). Elution of the ~10 kDa protein peak coincided with the elution of the peak of activity in fractions 27-28. However, a similar-sized protein continues to be eluted in fractions 30-31, where activity declines markedly. To identify the protein(s), samples from fraction 28 (peak of activity) and fraction 31, subsequent to the peak of activity, are subjected to mass spectrometric sequencing of tryptic peptides. A tryptic peptide of the sequence QIYYSDKYDDEEFYR, corresponding to amino acid residues 5-20 of human Cks1, is detected in the ~10 kDa protein of both fractions. The reason for the difference in the activity of the Cks1 protein in these different fractions is not known. Possibly, the Cks1 protein in fraction 31 is a denatured conformation that may have altered exclusion properties in the gel filtration column.

9.2.4 Activity of Cks1/Suc proteins

To address whether all Cks/Suc1 proteins used in this study were functional, we have examined their action in promoting multi-phosphorylation of cyclosome/APC by protein kinase Cdk1/cyclinB was examined (Patra, D. & Dunphy, W.G., 1998, *Genes Dev.* 12:2549; Shteinberg, M. & Hershko, A., 1999, *Biochem. Biophys. Res. Commun.* 257:12). Cyclosomes from S-phase HeLa cells were partially purified (Yudkovsky, et al., 2000, *Biochem. Biophys. Res. Commun.* 271:299) and incubated with 500 units of Suc1-free Cdk1/cyclin B (Shteinberg, M. & Hershko, A., 1999, *Biochem. Biophys. Res. Commun.* 257:12), as described (Yudkovsky, et al., 2000, *Biochem. Biophys. Res. Commun.* 271:299). Where indicated, 10 ng/ μ l of the corresponding Cks/Suc1 protein was supplemented. The

samples were subjected to immunoblotting with a monoclonal antibody directed against human Cdc27 (Transduction Laboratories). As shown in FIG. 47 the Cdk1-catalyzed hyperphosphorylation of Cdc27, a subunit of the cyclosome/APC, is markedly stimulated by all three recombinant Cks/Suc1 proteins. This is indicated by the decrease in the
5 unphosphorylated form of Cdc27 and its conversion to several hyperphosphorylated forms that migrate slower in SDS-polyacrylamide gel electrophoresis (FIG. 47, lanes 3-5) This large electrophoretic shift, promoted by all recombinant Cks/Suc1 proteins, requires the action of protein kinase Cdk1/cyclin B (FIG. 47, lane 6). All three bacterially expressed Cks/Suc1 proteins used are at least 95% homogeneous, as indicated by SDS-polyacrylamide
10 gel electrophoresis and Coomassie staining.

9.2.5 Confirmation that the factor required for p27-ubiquitin ligation is Cks1

Cks1 produced by *in vitro* translation (FIG. 48B, lane 3) or bacterially expressed, purified Cks1 (FIG. 48B, lane 6) effectively replaced the factor in this reaction.
15 This action is found to be specific for Cks1 and is not shared by other members of the Cks/Suc1 family of proteins. Human Cks2, which is 81% identical and 90% similar to Cks1, as well as the fission yeast homologue, Suc1, are completely inactive in this reaction, either when produced by *in vitro* translation (FIG. 48B, lane 4) or as bacterially expressed purified proteins (FIG. 48B, lanes 7 and 8) Purified recombinant Cks2 and Suc1 do not stimulate
20 p27-ubiquitin ligation even when added at up to 50-fold higher concentrations despite their being functional, as demonstrated by their ability to promote the multi-phosphorylation of Cdc27 by Cdk1. The combined evidence thus strongly indicates that the action of Cks1 in p27-ubiquitin ligation is specific and is not shared by other members of this protein family.

25 9.2.6 Cks1 promotes the ligation of ubiquitin to P27

Cks1 does not seem to be required for the action of all mammalian SCF complexes. In the well-characterized case of SCF ^{β -TrCP}, the purified complex carries out robust ubiquitinylation of I κ B *in vitro* (Tan, et al., 1999, Mol. Cell 3:527). Furthermore, the addition of Cks1 had no observable influence on the rate of the ligation of ubiquitin to
30 phosphorylated I κ B α by purified SCF ^{β -TrCP}. It seemed more likely that Cks1 is specifically involved either in the action of the SCF^{Skp2} complex or in some other process necessary for p27-ubiquitin ligation. Since p27 has to be phosphorylated on Thr-187 by Cdk2 for recognition by the SCF^{Skp2} complex (Carrano, et al., 1999, Mat. Cell Biol. 1:1993; Tsvetkov, et al., 1999, Current Biology 661) and since Cks proteins may stimulate the protein kinase
35 activity of some, but not all, Cdk/cyclin complexes (Reynard, et al., 2000, Mol. Cell Biol.

20:5858), it seems possible that Cks1 stimulates the phosphorylation of p27 by Cdk2. However, as shown in (FIG. 49A) p27 is rapidly phosphorylated by Cdk2/cyclin E in the absence of Cks1, and the addition of Cks1 has no significant influence on this process. The conclusion that Cks1 acts at a step subsequent to the phosphorylation of p27 is corroborated
5 by the finding that when purified p27 is first phosphorylated by incubation with Cdk2/cyclin E and $^{32}\text{P-}\gamma\text{-ATP}$, its subsequent ligation to MeUb still requires Cks1 (FIG. 49B). Therefore, Cks1 greatly stimulates the binding of phosphorylated p27 to Skp2.

9.2.7 Cks1 affects the binding of phosphorylated p27 to Skp2

10 Whether the step affected by Cks1 is the binding of phosphorylated p27 to Skp2 was assessed. Skp2/Skp1 complex was used instead of Skp2, because in the absence of Skp1, recombinant Skp2 is not expressed abundantly in insect cells in a soluble form. Previously small, but significant binding of ^{35}S -labeled, *in vitro*-translated p27 to Skp2/Skp1 was detected (by immunoprecipitation with an antibody directed against Skp2), which is
15 dependent upon its phosphorylation on Thr-187 by Cdk2/cyclin E (Carrano, et al., 1999, Nat. Cell Biol 1:193). Using a similar procedure, the binding of p27 to Skp2/Skp1 is greatly stimulated by Cks1 (FIG. 49C, lanes 2 and 3). This action requires the phosphorylation of p27 on Thr-187, since binding of the non-phosphorylatable mutant Thr-187-Ala did not occur even in the presence of Cks1 (FIG. 49C, lanes 4 and 5). To examine whether this
20 action of Cks1 also occurs in a completely purified system devoid of reticulocyte lysate present in preparations of *in vitro*-translated p27, a similar experiment is performed with bacterially expressed, purified p27 that is phosphorylated by $^{32}\text{P-}\gamma\text{-ATP}$. In this case there is some non-specific binding of phosphorylated p27 to anti-Skp2-Protein A beads in the absence of Skp2. Still, a marked stimulation of the specific binding of ^{32}P -p27 to Skp2/Skp1
25 by Cks1 is observed (FIG. 49D). Therefore, Cks1 greatly stimulates the binding of phosphorylated p27 to Skp2.

As shown in FIG. 50A, a strong binding of ^{35}S -Cks1 to the Skp2/Skp1 complex was observed. Under similar conditions, no binding of ^{35}S -Cks2 to Skp2/Skp1 was seen. Since in these experiments Skp2/Skp1 complex is used (because of the lack of
30 recombinant native Skp2), it is examined whether Cks1 may bind to Skp1 in the absence of Skp2. In the experiment shown in FIG. 50B, ^{35}S -Cks1 is incubated with either His₆-Skp1 or with Skp2/His₆-Skp1 complex, and then binding to Ni-NTA-agarose beads is estimated. A strong binding of Cks1 to Skp2/His₆-Skp1 but not to His₆-Skp1 was observed. Thus, human Cks1 specifically binds to the Skp2/Skp1 complex, likely through the Skp2 protein.

35

The results presented herein demonstrate that the binding of Skp2 to phosphopeptide-Sepharose beads (but not to control beads that contained an identical but unphosphorylated p27-derived peptide) is greatly increased by Cks1 (FIG. 50C). These findings indicate that binding to this phosphopeptide can serve as a valid tool to study Cks1-assisted Skp2-p27 interaction. Using the same p27-derived peptide beads, significant binding of ³⁵S-Cks1 to phosphorylated p27 peptide, but not to unphosphorylated p27 peptide is observed FIG. 50D. These findings indicate that Cks1 binds directly to phospho-Thr187 of p27 and demonstrate that the presence of Cdk2/cyclin E is not obligatory for the binding of Skp2 to phosphorylated p27.

10

10. EXAMPLE: ASSAY TO IDENTIFY AN FBP INTERACTION WITH A CELL CYCLE REGULATORY PROTEIN (e.g., SKP2 with E2F)

The following study was conducted to identify novel substrates of the known FBP, Skp2.

As shown in Figure 44, E2F-1, but not other substrates of the ubiquitin pathway assayed, including p53 and Cyclin B, physically associates with Skp2. Extracts of insect cells infected with baculoviruses co-expressing Skp2 and E2F-1, (lanes 1,4 and 5), or Skp2 and hexa-histidine p53 (His-p53) (lanes 2,6,7,10 and 11), or Skp2 and His-Cyclin B (lanes 3,8,9,12, and 13) were either directly immunoblotted with an anti-serum to Skp2 (lanes 1 - 3) or first subjected to immunoblotted with an anti-serum to Skp2 (lanes 1 - 3) or first subjected to immunoprecipitation with the indicated antibodies and then immunoblotted with an anti-serum to Skp2 (lanes 4 - 13). Antibodies used in the immunoprecipitations are: normal purified mouse immunoglobulins (IgG) (lane 4,6,10 and 12), purified mouse monoclonal anti-E2F-1 antibody (KH-95, from Santa Cruz) (lane 5), purified mouse monoclonal anti-p53 antibody (DO-1, from Oncogene Science) (lane 7), purified rabbit IgG (lane 8), purified rabbit polyclonal anti-Cyclin B antibody (lane 9), purified mouse monoclonal anti-His antibody (clone 34660, from Qiagen) (lanes 11 and 13).

As shown in Figure 44B, Skp2 physically associates with E2F-1 but not with other substrates of the ubiquitin pathway (p53 and Cyclin B). Extracts of insect cells infected with baculoviruses co-expressing Skp2 and E2F-1 (lanes 1 - 3), or Skp2 and His-p53 (lanes 4 - 6), or Skp2 and His-Cyclin B (lanes 7 - 9) were either directly immunoblotted with antibodies to the indicated proteins (lanes 1,4 and 7) or first subjected to immunoprecipitation with the indicated anti-sera and then immunoblotted with antibodies to the indicated proteins (lanes 2,3,5,6,8 and 9). Anti-sera used in the immunoprecipitations are: anti-Skp2 serum (lanes 2,5 and 8), and normal rabbit serum (NRS) (lane 3,6 and 9).

As shown in Figure 44C, E2F-1 physically associates with Skp2 but not with another F-box protein (FBP1). Extracts of insect cells infected with baculoviruses co-expressing Skp2 and E2F-1 (lanes 1,3 and 4), or Flag-tagged-FBP1 and E2F-1 (lanes 2,5 and 6) were either directly immunoblotted with a mouse monoclonal anti-E2F-1 antibody (lanes 1 and 2) or first subjected to immunoprecipitation with the indicated antibodies and then immunoblotted with a mouse monoclonal anti-E2F-1 antibody (lanes 3 - 6). Antibodies used in the immunoprecipitations are: anti-Skp2 serum (lanes 3), NRS (lane 4), purified rabbit polyclonal anti-Flag (lane 5), purified rabbit IgG (lane 6).

The methodology used in this example can also be applied to identify novel substrates of any FBP, including, but not limited to, the FBPs of the invention, such as FBP1, FBP2, FBP3a, FBP3b, FBP4, FBP5, FBP6, FBP7, FBP8, FBP9, FBP10, FBP11, FBP12, FBP13, FBP14, FBP15, FBP16, FBP17, FBP18, FBP19, FBP20, FBP21, FBP22, FBP23, FBP24, and FBP25.

The invention is not to be limited in scope by the specific embodiments described which are intended as single illustrations of individual aspects of the invention, and functionally equivalent methods and components are within the scope of the invention. Indeed various modifications of the invention, in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and accompanying drawings. Such modifications are intended to fall within the scope of the appended claims.

All references cited herein are incorporated herein by reference for all purposes.

WHAT IS CLAIMED IS:

1. A method for screening compounds useful for the treatment of proliferative and
differentiative disorders comprising contacting a compound with a cell or a cell extract
5 expressing Skp2 and one or both of p27 and Cks1, and detecting a change in the activity of
Skp2.
2. The method of Claim 1 wherein the change in the activity of Skp2 is detected by
detecting a change in the interaction of Skp2 with either p27 or Cks1.
10
3. The method of Claim 1 wherein the change in the activity of Skp2 is detected by
detecting a change in the ubiquitination of p27 or degradation of p27 or Cks1.
4. A method for screening compounds useful for the treatment of proliferative and
15 differentiative disorders comprising adding a compound in a purified system containing Skp2
and one or both of p27 and Cks1, and detecting a change in the activity of Skp2.
5. The method of Claim 4 wherein the change in the activity of Skp2 is detected by
detecting a change in the interaction of Skp2 with either p27 or Cks1.
20
6. The method of Claim 4 wherein the change in the activity of Skp2 is detected by
detecting a change in the ubiquitination of p27 or degradation of p27 or Cks1.
7. A method for screening compounds useful for the treatment of proliferative and
25 differentiative disorders comprising adding a compound in a purified system containing Skp2
and one or both of a polypeptide corresponding to the carboxy terminus of the human p27
chain having the sequence NAGSVEWTPKKPGLRRRQT with or without a
phosphothreonine at position 187 and Cks1, and detecting a change in the activity of Skp2.
- 30 8. The method of Claim 7 wherein the change in the activity of Skp2 is detected by
detecting a change in the interaction of Skp2 with either the polypeptide or Cks1.
9. The method of Claim 7 wherein the change in the activity of Skp2 is detected by
detecting a change in the ubiquitination of the polypeptide or degradation of the polypeptide
35 or Cks1.

FIG. 1




FIG. 2

10 20 30 40 50 60
MDPAEAVLQEKALKFMNSSEREDCNNGEPPRKIIPEKNSLRQTYNSCARLCLNQETVCLA
70 80 90 100 110 120
STAMKTENCVAKTKLANGTSSMIVPKQRKLSASYEKEKELCVKYFEQWSESDQVEFVEHL
130 140 150 160 170 180
ISQMCHYQHGHSYLPMLQRDFITALPARGLDHIAENILSYLDAKSLCAAELVCKEWY
190 200 210 220 230 240
RVTS D GMLWKKLIERMVRTDSLWRGLAERRGWGQYLFKNKPPDGNAPPNSFYRALYPKII
250 260 270 280 290 300
QDIETIESNWRCGRHSLQRIHCRSETSKGVYCLQYDDQKIVSGLRDNTIKIWDKNTLECK
310 320 330 340 350 360
RILTGHTGSVLCLQYDERVIITGSSDSTVRVWDVNTGEMLNTLIHHCEAVLHLRFNNGMM
370 380 390 400 410 420
VTCSKDRSIAVWDMASPTDITLRRVLVGHRAAVNVVDFDDKYIVSASGDRTIKVWNTSTC
430 440 450 460 470 480
EFVRTLNHGKRGIAQLQYRDLVVGSSDNTIRLWDIECGACLRVLEGHEELVRCIRFDN
490 500 510 520 530 540
KRIVSGAYDGGIKVWDLVAALDPRAPAGTLCRLTLVEHSGRVFRLQFDEFQIVSSSHDDT
550 560
ILIWDFLNDPAAQAEPPRSPSRITYTISR

FIG. 3A

[illegible]

FIG. 3B

10 20 30 40 50 60
MERKDFETWLDNISVTFLSLTDLQNETLDHLISLSGAVQLRHLSSNNLETLLKRDFLKL
70 80 90 100 110 120
PLELSFYLLKWLDLPQTLLTCLVSKQWNKVISACTEVWQTACKNLGWQIDDSVQDALHWK
130 140 150 160 170 180
KVYLKAILRMKQLEDHEAFETSSLIGHSARVYALYYKDGLLCTGSDDLAKLWDVSTGQC
190 200 210 220 230 240
VYGIQTHTCAAVKFDEQKLVTSFNTVACWEWSSGARTQHFRGHTGAVFSVDYNDELDI
250 260 270 280 290 300
LVSGSADFTVKVWALSAGTCLNTLTGHTTEWVTKVVLQKCKVKSSLHSPGDYILLSADKYE
310 320 330 340 350 360
IKIWPIGREINCKCLKTLVSSEDRSICLQPRLHFDGKYIVCSSALGLYQWDFASYDILRV
370 380 390 400 410 420
IKTPEIANLALLGFGDIFALLFDNRYLYIMDLRTESLISRWPLPEYRESKRGSSFLAGEH
PG

FIG. 4A

10 20 30 40 50 60 70 80 90
ATGGAGAGAAAGGACTTTGAGACATGGCTTGATAACATTCTGTTACATTCTTCTCTGACGGACTTGCAGAAAAATGAAGCTCGGATCAACC
100 110 120 130 140 150 160 170 180
TGATTAGTCTGAGTGGGGCAGTCCAGCTCAGGCATCTCTCCAATAACCTAGAGACTCTCTCAAGCGGGACTTCTCAAACCTCCTTCCCCTGGA
190 200 210 220 230 240 250 260 270 280
GCTCAGTITTTTATTTGTTAAAAATGGCTCGATCCTCAGACTTTACTCAGATGCTGCTCTCTAAACAGTGGAAATAAGGTGATAAGTGCTGT
290 300 310 320 330 340 350 360 370
ACAGAGGTCTGGCAGACTGCATGTAAAAATTTGGGCTGGCAGATAGATGATTCTGTTTCAGGACGCTTTGCACTGGAAGAAGGTTTATTGAAGG
380 390 400 410 420 430 440 450 460 470
CTATTTTGAGAATGAAGCAACTGGAGGACCATGAAGCCTTTGAAACCTCGTCATTAAATTGGACACAGTGCCAGAGTGTATGCACCTTACTACAA
480 490 500 510 520 530 540 550 560
AGATGGACTTCTCTGTACAGGGTCAGATGACTTGTCTGCAAGCTGTGGGATGTGAGCACAGGGCAGTCCCTTTATGGCATCCAGACCCACACT
570 580 590 600 610 620 630 640 650
TGTGCAGCGGTGAAGTTTGATGAACAGAAGCTTGTGACAGGCTCTTTGACAACACTGTGGCTTGTCTGGGAATGGAGTTCGGAGCCAGGACCC
660 670 680 690 700 710 720 730 740 750
AGCACTTTCCGGGGCACACGGGGGGGTATTTAGCCTGGACTACAATGATGAAGTGGATATCTTGGTGAGCGGCTCTGCAGACTTCACTGTGAA
760 770 780 790 800 810 820 830 840
AGTATGGGCTTTATCTGCTGGACATGCTGAACACACTCACCAGGGCACAGGAATGGGTACCAAGCTAGTTTTCAGAAATGCAAGTCAAG
850 860 870 880 890 900 910 920 930 940
TCTCTCTTGACAGTCTGGAGACTACATCTCTTAAGTGCAGACAAATATGAGATTAGATTGGGCAATTGGGAGAGAAATCAACTGTAAAT
950 960 970 980 990 1000 1010 1020 1030
GCTTAAAGACATTGCTCTCTCTGAGGATAGAAGTATCTGCTGCAGCCAGACTTCATTTTGATGGCAATACATTGCTGTAGTTTCAGCACT
1040 1050 1060 1070 1080 1090 1100 1110 1120
TGGTCTCTACCACTGGGACTTTGCCAGTTATGATATTCTCAGGGTCATCAAGACTCTGAGATAGCAAACTTGGCCTTGGCTTGGCTTGGAGAT
1130 1140 1150 1160 1170 1180 1190 1200 1210 1220
ATCTTTGCCCTGCTGTTTGACAACCGCTACCTGTACATCATGGACTTGGGACAGAGAGCCTGATTAGTCCCTGCTCTGCCAGAGTACAGGG
1230 1240 1250 1260 1270 1280 1290 1300 1310
AATCAAAGAGAGGCTCAAGCTTCTGGCAGCGGAACATCTGGCTGAATGGACTGGATGGGCACAATGACACGGCTTGGCTTTTCCACCAGC
1320 1330 1340 1350 1360 1370 1380 1390 1400 1410
ATGCTTGACCACAGTATTACCTGGTGTGTGGAAGGAGCAGGGCTGACACCATGAGCCACCAACCGCTGACTGACTTTGGGTCCCGGGCTGCG
1420 1430 1440 1450 1460 1470
GGTTTGGGTGCACCTCTGCGGCACGGGACTGCATGAACCAAAGTTCTCACCTAATGGTATCATCA

FIG. 4B

10 20 30 40 50 60
MKRGGGRSDRNSSEEGTAEKSKLRTTNEHSQTCDWGNLLQDIILQVFKYLPLLDRAHAS
70 80 90 100 110 120
QVCRNWNQVFHMPDLWRCFEFELNQPATSYLKATHPELIKQIIKRHSNHLQYVSFKVDSS
130 140 150 160 170 180
KESAEACDILSQLVNCSLKTILGLISTARPSFMDLPKSHFISALTVVVFVNSKSLSSLKID
190 200 210 220 230 240
DTPVDDPSLKVLVANNSTLKLKMSSCPHVSPAGILCVADQCHGLRELALNYHLLSDEL
250 260 270 280 290 300
LLALSSEKHVRLEHLRIDVVSENPGQTHFHTIQKSSWDAFIRHSPKVNLMYFFLYEEEF
310 320 330 340 350 360
DPFFRYEIPATHLYFGRSVSKDVLGRVGMTCPRLVELVVCANGLRPLDEELIRIAERCKN
370 380 390 400 410 420
LSAIGLGECEVSCSAFVEFVKMCGGRLSQLSIMEEVLPDQKYSLEQIHWEVSKHLGRVW
FPDMPTW

FIG. 5A

10 20 30 40 50 60 70 80 90
CGGGGTGGTGTGTGGGGGAAGCCGCCCGGCGAGCAGGATGAACAGGAGGAGAGATAGTGACCGTAATTCATCAGAAGAAGAACTGCAGA
100 110 120 130 140 150 160 170 180
GAAATCCAAGAACTGAGGACTACAAATGAGCATTCTCAGACTTGTGATTTGGGTAATCTCCTTCAGGACATTATTCTCCAAGTATTTAAATAT
190 200 210 220 230 240 250 260 270 280
TTGCCCTCTCTTGACCGGGCTCATGCTTCACAAGTTTGGCGCAACTGGAAACAGGTATTTTCATGCGCTGACTTGTGGAGATGTTTGAATTTG
290 300 310 320 330 340 350 360 370
AACTGAATCAGCCAGCTACATCTTATTTGAAAGCTACCCATCCAGAGCTGATCAAAACAGATTATTAAAGACATTCAAACCATCTACAATATGT
380 390 400 410 420 430 440 450 460 470
CAGCTTCAAGGTTGGACAGCAGCAAGGAATCAGCTGAAGCAGCTTGTGATATACTATCGCAACTTGTGAATTGCTCTTTAAAAACACTTGGACTT
480 490 500 510 520 530 540 550 560
ATTTCAACTGCTCGACCAAGCTTTATGGATTTACCAAAGTCTCACTTTATCTCTGCACTGACACTTGTGTTTGTAACTCCAAATCCCTGCTCTT
570 580 590 600 610 620 630 640 650
CGCTTAAGATAGATGATCTCCAGTAGATGATCCATCTCTCAAAGTACTAGTGGCTAAACAATAGTGATACACTCAAGCTGTTGAAAAATGAGCAG
660 670 680 690 700 710 720 730 740 750
CTGCTCTCATGCTCTCCAGCAGGTATCCTTTGTGTGGCTGATCAGTGTACCGGCTTAAGAGAACTAGCCCTGAACTACCACCTTATTGAGTGAT
760 770 780 790 800 810 820 830 840
GAGTTGTTACTTGCAATTGCTCTCTGAAAAACATGTTTCGATTAGAACATTTCGCCATTGATGTAGTCACTGACAATCTCGACAGACACACTTCC
850 860 870 880 890 900 910 920 930 940
ATACTATTCAAGAGTAGCTGGGATGCTTTTCATCAGACATTCAOCCAAAGTGAACCTTAGTGATGTATTTTTTTTATATGAAGAAGAAATTTGA
950 960 970 980 990 1000 1010 1020 1030
CCCCCTCTTTTCGCTATGAATACCTGGCACCCATCTGTACTTTGGGASATCAGTAAGCAAGATGTGCTTGGCCGTGTGGGAATGACATGCCCT
1040 1050 1060 1070 1080 1090 1100 1110 1120
AGACTGCTTGAAGTAGTAGTGTGTGCAAAATGGATTACGGCCACTTGTGAAGAGTTAATTCCGATTCAGAACGTTGCAAAAAATTTGTCAAGCTA
1130 1140 1150 1160 1170 1180 1190 1200 1210 1220
TTGGACTAGGGGAATGGAAGTCTCATGTAGTGCCTTTGTTGAGTTTGTGAAGATGTGTGGTGGCCGCTATCTCAATTATCCATTATGGAAGA
1230 1240 1250 1260 1270 1280 1290 1300 1310
AGTACTAATTCCTGACCAAAAGTATAGTTTGGAGCAGATTCACTGGGAAGTGTCCAAGCATCTTGGTAGGGTGTGCTTTCCCGACATGATGCC
1320 1330 1340 1350 1360 1370 1380 1390 1400
ACTTGGTAAAAACTGCATGATGAATAGCACCTTAATTTCAAGCAATGTATTATAATTAAAGTTTATTTGCTGTAAAAA

FIG. 5B

10 20 30 40 50 60
MKRNSLSVENKIVQLSGAAKQPKVGFYSSLNQTHHTVLLDWGSLPHHVVLQIFQYLPLL
70 80 90 100 110 120
DRACASSVCRRWNEVFHISDLWRKFELNQSATSSFKSTHPDLIQQIIKKHFAHLQYVS
130 140 150 160 170 180
FKVDSSAESAEAACDILSQLVNCSIQTLGLISTAKPSFMNVSESHFVSALTVVFINSKSL
190 200 210 220 230 240
SSIKIEDTFVDDPSLKILVANNSDTLRLPKMSSCPHVSSDGILCVADRCQGLRELALNYY
250 260 270 280 290 300
ILTDELFLALSSETHVNLEHLRIDVVSSENPQGIKFHAVKKHSDALIKHSPRVNVVMHFF
310 320 330 340 350 360
LYEEEFETFFKEETPVTHLYFGRSVSKVVLGRVGLNCPRLIELVVCANDLQPLDNELICI
370 380 390 400 410 420
AEHCTNLTAALGLSKCEVSCSAFIRFVRLCERRLTQLSMEEVLIPDEDYSLDEIHTEVSK
430
YLGRVWFDPVMPWL

FIG. 6A

10 20 30 40 50 60
ACATTTCTAATGTTTACAGAATGAAGAGGAACAGTTTATCTGTTGAGAATAAAATTGTCCAGTTGTCA
70 80 90 100 110 120 130
GGAGCAGCGAAACAGCCAAAAGTTGGGTTCTACTCTTCTCTCAACCAGACTCATACACACACGGTTCTT
140 150 160 170 180 190 200
CTAGACTGGGGAGTTTGCCCTCACCATGTAGTATTACAAATTTTTCAGTATCTTCTTTACTAGATCGG
210 220 230 240 250 260 270
GCCTGTGCATCTTCTGTATGTAGGAGGTGGAATGAAGTTTTCATATTTCTGACCTTTGGAGAAAGTTT
280 290 300 310 320 330 340
GAATTGAACTGAACCAGTCACTACTTCTCTTTAAGTCCACTCATCTGATCTCATTTCAGCAGATC
350 360 370 380 390 400 410
ATTAAAAAGCATTGTGCTCATCTTCAGTATGTACAGCTTTAAGGTTGACAGTAGCGCTGAGTCAGCAGAA
420 430 440 450 460 470 480
GCTGCCCTGTGATATACTCTCTCAGCTGGTAAATGTTCCATCCAGACCTTGGGCTTGATTTCACAGCC
490 500 510 520 530 540 550
AAGCCAAGTTTCATGAATGTGTGGAGTCTCATTGTGTGTCAGCACTTACAGTTGTTTTATCAACTCA
560 570 580 590 600 610 620
AAATCATTATCATCAATCAAAATGAAGATACACCAGTGGATGATCCTTCATTGAAGATTCTTGTGGCC
630 640 650 660 670 680 690
AATAATAGTGACACTCTAAGACTCCCAAGATGAGTAGCTGTCTCATGTTTCATCTGATGGAATTCTT
700 710 720 730 740 750
TGTGTAGCTGACCGTTGTCAAGGCCCTTAGAGAACTGGCGTTGAATTATTACATCCTAAGTGAAGTT
760 770 780 790 800 810 820
TTCCTTGCACTCTCAAGCGAGACTCATGTTAACCTTGAACATCTTCGAATTGATGTTGTGAGTGAATA
830 840 850 860 870 880 890
CCTGGACAGATTAAATTTTCATGCTGTTAAAAACACAGTTGGGATGCACTTATTAACATTCCCTAGA
900 910 920 930 940 950 960
GTTAATGTTGTTATGCACTTCTTCTATATGAAGAGGAATTCGAGACGTTCTTCAAAGAAGAAACCCCT
970 980 990 1000 1010 1020 1030
GTTACTCACCTTTATTTTGGTCGTTTCAGTCAGCAAGTGGTTTTAGGACGGGTAGGTCTCAACTGTCTCT
1040 1050 1060 1070 1080 1090 1100
CGACTGATTGAGTTAGTGGTGTGTCTAATGATCTTCAGCCTCTTGATAATGAAGTTATTGTATTGCT
1110 1120 1130 1140 1150 1160 1170
GAACACTGTACAAACCTAACAGCCTTGGGCCCTCAGCAAAATGTGAAGTTAGCTGCAGTGCCTTCATCAGG
1180 1190 1200 1210 1220 1230 1240
TTTGTAAGACTGTGTGAGAGAAGGTTAACACAGCTCTCTGTAATGGAGGAAGTTTGATCCCTGATGAG
1250 1260 1270 1280 1290 1300 1310
GATTATAGCCTAGATGAATTCACACTGAAGTCTCCAAATACCTGGGAAGAGTATGGTTCCCTGATGTG
1320
ATGCCCTCTCTGG

FIG. 6B

10 20 30 40 50 60
MAGSEPRSGTNSPPPPFSDWGRLEAAILSGWKTFWQSVSKDRVARTTSREEVDEAASTLT
70 80 90 100 110 120
RLPIDVQLYILSFLSPHDLCLGSTNHYWNETVRNPILWRYFLLRDLPSWSSVDWKSPLY
130 140 150 160 170 180
LQILKKPISEVSDGAFFDYMAVYLMCCPYTRASKSSRPMYGAVTSFLHSLIIPNEPRFA
190 200 210 220 230 240
LFGPRLEQLNTSIVLSLLSSEELCPTAGLPQRQIDGIGSGVNFQLNNQHKNILILYSTT
250 260 270 280 290 300
RKERDRAREEHTSAVNKMFSTRHNEGDDRPGRSVIPQIQKLCEVVDGFIYVANAIAHAKR
310 320 330 340 350 360
HEWQDEFSSHIMAMTDPAFGSSGRPLLVLSCISQGDVKRMPCFYLAHELHLNLLNHPWLQQ
370 380 390 400 410 420
DTEAETLTGFLNGIEWILEEVESKRAR*FSFQILGTETI*NLLLR*CEYLLSQPTLSCL
430 440 450 460 470 480
FADRLSFGQL*LLCFLYYFYFLP*INYKKRVSVLVFSPKMNL*TFFW*FLYFLSF*KY*I

L

FIG. 7A

10 20 30 40 50 60
ATGGCGGGAAGCGAGCGCGCGGGAACAAATTCGCCGCCGCCGCCCTTCAGCGACTGGGGCGCGCTG
70 80 90 100 110 120 130
GAGGGCGCCATCCTCAGCGGCTGGAAGACCTTCTGGCAGTCAGTGAGCAAGGATAGGGTGGCGGTACG
140 150 160 170 180 190 200
ACCTCCCGGAGGAGGTGGATGAGGCGGCCAGCACCTTGACCGCGCTGCCGATTGATGTACAGCTATAT
210 220 230 240 250 260 270
ATTTTGTCCCTTCTTTCACCTCATGATCTGTGTCAGTTGGGAAGTACAAATCATTATTGGAATGAAACT
280 290 300 310 320 330 340
GTAAGAAATCCAATCTGTGGAGATACTTTTGTGAGGGATCTTCCTTCTTGGTCTTCTGTGACTGG
350 360 370 380 390 400 410
AAGTCTCTTCATATCTACAAATCTTAAAAAGCCTATATCTGAGGTCTCTGATGGTGCATTTTTGTGAC
420 430 440 450 460 470 480
TACATGGCAGTCTATCTAATGTGCTGTGCATACACAAGAAGAGCTTCAAAATCCAGCGCTCTATGTAT
490 500 510 520 530 540 550
GGAGCTGTCACTTCTTTTACACTCCCTGATCATTCCTCAATGAACCTCGATTGTGCTCTGTTTGGACCA
560 570 580 590 600 610 620
CGTTTGGAAACAATTGAATACCTCTTTGTGTTGAGCTTGCTGTCTTCAGAGGAACCTTGGCCCAACAGCT
630 640 650 660 670 680 690
GGTPTGCCCTCAGAGGCGAGATTGATGATGGATCAGGAGTCAATTTTCAGTTGAACAACCAACATAAA
700 710 720 730 740 750
TTCAACATTCTAATCTTATATTCAACTACCAGAAAGGAAAGAGATAGAGCAAGGGAAGAGCATACAAAT
760 770 780 790 800 810 820
GCAGTTAACAAGATGTTCACTCGACACAAATGAAGGTGATGATCGACCAGGAAGCGGTACAGTGTGATT
830 840 850 860 870 880 890
CCACAGATTCAAAAAGTGTGTAAGTTGTAGATGGGTTCATCTATGTTGCAAAATGCTGAAGCTCATAAA
900 910 920 930 940 950 960
AGACATGAATGGCAAGATGAATTTCTCATATTATGGCAATGACAGATCCAGCCCTTTGGGTCTTCGGGA
970 980 990 1000 1010 1020 1030
AGACCATTTGTTGGTTTTATCTTGTATTCTCAAGGGGATGTAAAAAGAAATGCCCTGTTTTATTGTGCT
1040 1050 1060 1070 1080 1090 1100
CATGAGCTGCATCTGAATCTTCTAAATCACCCTATGGCTGGTCCAGGATACAGAGGCTGAAACTCTGACT
1110 1120 1130 1140 1150 1160 1170
CGTTTTTTGAATGGCATTGAGTGGATTCTTGAAGAAGTGAATCTAAGCGTGCAAGATGATTCTTTTT
1180 1190 1200 1210 1220 1230 1240
CAGATCTTGGGAAGTGAACCATTTGAAATTTATTACTAAGGTGCTGATGTGAATATTGCTCAGTCAG
1250 1260 1270 1280 1290 1300 1310
CCCACCTTGTCTGCTTTTTCAGATAGGCTTTCAATTTGGACAGCTATAACTGCTGTGTTTTTATAT
1320 1330 1340 1350 1360 1370 1380
TATTTTACTTTTACCATAAATCAATTACAGAAAAGAGTTTCAGTCTAGTATTTAGCCCCAAAATG
1390 1400 1410 1420 1430 1440
AACCTTTAAACATTTTTTGGTAATTTTATATTTCTGTCTTTTAAAAATATTAAATTTTG

FIG. 7B

10 20 30 40 50 60
MSRRPCSCALRPPRCSCSASPSAVTAAGRPRPSDSCKEESSTLSVKMKCDFNCNHVHSGL
70 80 90 100 110 120
KLVKPDDIGRLVSYTPAYLEGSCKDCIKDYERLSCIGSPIVSPRIVQLETESKRLHNKEN
130 140 150 160 170 180
QHVQQTNLNSTNEIEALETSLRYEDSGYSSFSLSQGLSEHEEGSLLEENFGDSLQSCLLQI
190 200 210 220 230 240
QSPDQYPNKNLLPVLHFEKVVCSTLKKNAKRNPKVDREMLKEIIARGNFRLQNIIGRKMG
250 260 270 280 290 300
LECVDILSELFRRGLRHVLATILAQLSDMDLINVSKVSTTWKKILEDDKGAFQLYSKAIQ
310 320 330 340 350 360
RVTENNNKFSPHASTREYVMFRTPLASVQKSAQAQTSLLKDAQTKLSNQGDQKGSTYSRHN
370 380 390 400 410 420
EFSEVAKTLKKNESLKACIRCNSPAKYDCYLQRATCKREGCGFDYCTKCLCNHYHTTKDCS
430 440
DGKLLKASCKIGPLPGTKKSKKNLRRL

FIG. 8A

10 20 30 40 50 60 70 80 90
AGGTTGCTCAGTGGCCCCGGAGCGGTTCTCCACCTGAGGCAGACACCACTCGGTTGGCATGAGCCGGCCCGCTGCAGCTGCCCGCTACGG

100 110 120 130 140 150 160 170 180
CCACCCCGCTGCTCTGCAGCCCGAGCCCGAGCGGAGTGACAGCCCGGGCGCCCTCGACCCCTGGATAGTTGTAAGAAGAAAGTTCTACCC

190 200 210 220 230 240 250 260 270 280
TTTCTGTCAAAATGAAGTGTGATTTTAATTGTAAACCATGTTTCATTCGGGACTTAAACTGGTAAAACTGATGACATGGAAGACTAGTTTCTTA

290 300 310 320 330 340 350 360 370
CACCCCTGCATATCTGGAAGCTTCTCTAAAGACTGCATTAAAGACTATGAAAGGCTGTCTATGTTGGGTACCGGATTTGTAGCCCTAGGATT

380 390 400 410 420 430 440 450 460 470
GTACAACTTGAAGTGAAGCAAGCGCTTGCATAAAGGAAATCAACATGTGCAACAGACACTTAATAGTACAAATGAATAGAACACTAG

480 490 500 510 520 530 540 550 560
AGACCACTAGACTTTATGAAGACAGTGGCTATTCCTCATTTTCTCTACAAAGTGGCCCTCAGTGAACATGAAGAAGGTAGCCTCTCGAGGAGAA

570 580 590 600 610 620 630 640 650
TTTCGCTGACAGTCTACAACTCTGCTGCTACAAATACAAAGCCAGACCAATATCCCAACAAAACTTCTGCTCAGTTCTTCTATTTTGA AAAA

660 670 680 690 700 710 720 730 740 750
GTGGTTTGTTCACATTAAAAAAGATGCAAAAGCAAAATCTCTAAAGTAGATCGGGAGATGCTGAAGGAAATTAGCCAGAGAAATTTTAGAC

760 770 780 790 800 810 820 830 840
TGCAGAAATATAATTGGCAGAAAAATGGCCCTAGAAATGTGTAGATATTTCTCAGCGAACTCTTTCGAAGGGGACTCAGACATGCTTACCAACTAT

850 860 870 880 890 900 910 920 930 940
TTTAGCACAATCAGTGACATCGACTTAATCAATGTGTCTAAAGTCAGCACAATTTGGAAGAAGATCCTAGAGATGATAAGGGGGGATTCAG

950 960 970 980 990 1000 1010 1020 1030
TTGTACAGTAAAGCAATACAAAGAGTTACCGAAAAACAATAAATTTTCACCTCATGCTTCAACCAGAGAATATGTTATGTTAGAACCCAC

1040 1050 1060 1070 1080 1090 1100 1110 1120
TGGCTTCTGTTCAGAAATCAGCAGCCAGACTTCTCTCAAAAAGATGCTCAAAACCAAGTTATCCAAATCAAGGTGATCAGAAAGGTTCTACTTA

1130 1140 1150 1160 1170 1180 1190 1200 1210 1220
TAGTCGACACAATGAATTTCTGTGAGTTGCCAAGACATTCAAAAAGAACGAAAGCCTCAAGCCCTGATTTGCTGTAAATTCACCTGCAAAATAT

1230 1240 1250 1260 1270 1280 1290 1300 1310
GATTGCTATTTFACAAAGGGCAACCTGCAAAAGCAGAAAGGCTGTGGATTGATTATTGTACGAAGTGTCTCTGTAAATATCATACTACTAAAGACT

1320 1330 1340 1350 1360 1370 1380 1390 1400 1410
GTTTCAGATGCGAAGCTCTCAAGCCAGTTGTAAATAGGTCCCTGCTGCTGACAAAGAAAGCAAAAGAAATTTACGAAGATCTGTATCTCT

1420 1430 1440 1450 1460 1470 1480 1490 1500
TATTAAATCAATTTCTACTGATCATGAATGTTAGTTAGAAAATGTTAGGTTTAACTTAAAAAAATTTGATTGTGATTTTCAATTTTATGTTG

1510 1520 1530 1540 1550 1560 1570 1580 1590
AAATCGGTGTAGTATCTGAGGTTTTTTTCCCGCCAGAAAGATAAGAGGATAGACAACTCTTAAAAATTTTACAAATTTAATGAGAAAAAGT

1600 1610 1620 1630 1640 1650 1660 1670 1680 1690
TTAAAAATCTCAATACAAATCAAAATTTAAATATTTTAAAGAAAAAGGAAAGTAGATAGTACTGAGGGTAAAAAAATTTGATTCAA

1700 1710 1720 1730 1740 1750 1760 1770 1780
TTTTATGTTAAAGGAAACCCATGCAATTTTACCTAGACAGTCTTAAATATGTTCTGTTTCCATCTGTTAGCATTTTACAGATTTTATGTTCT

1790 1800 1810 1820 1830 1840 1850 1860 1870 1880
CTTACTCAATGATACCAACAGAAATATCAACTTCTGGAGTCTATTAAATGTTGTTGACCTTTCTAAAGCTTTTTCATTTGTTCTATTTC

1890 1900 1910 1920 1930 1940 1950 1960 1970
CAAGAAAGTATCTTTGTAAAACTTGTCTTTTCTCTATTCTGAAATCTTTTAAATATTTTGTATACATGTAATATTTCTGTATTTTT

1980 1990 2000 2010 2020 2030 2040 2050 2060
TATATGTCAAAGAATATGTTCTTGTATGTACATATAAAAAATAATTTGCTCAATAAAATTTGTAAGCTTAAAAAAATTTTAACTCGAG

2070
ACTAGTGC

FIG. 8B

10 20 30 40 50 60
ARSGASALRRRRVQVWLSRPPPGGDSFRTTRPQRGPGPGGSQAMDAPHSKAALDSINE
70 80 90 100 110 120
LPDNILLELFTHVPARQLLLNCRSLVCSLWRDLIDLLTLWKRKCLRKGFITKDWDQPVADW
130 140 150 160 170 180
KIFYFLRSLHRNLLRNPCAENDMFAWQIDFNGGDRWKVDSLPGAHGTEFPDPKVKKSFVT
190 200 210 220 230 240
SYELCLKWELVDLLADRYWEELLDTRPDIVVKDWFAARADCGCTYQLKVQLASADYFVL
250 260 270 280 290 300
ASFEPPTIQQWNNATWTEVSYTFSFYPRGVRYILFQHGGRDTQYWAGWYGPRTNSSI
310 320 330
VVSPKMTNRNQASSEAQPGQKHGQEEAAQSPYGAVVQIF

FIG. 9A

10 20 30 40 50 60 70 80 90
GCGCGTTGGGAGCTTCGGCCCTGCGTAGGAGGCGGGTGACAGGTGTGGGTCTGAGCGCGCCCGCCCTGGAGGGGGAGACAGCTTCAGGACAC
100 110 120 130 140 150 160 170 180
GCAGGCGCGACGAGGGGCGGGGCGGGGATCCAGGCCATGGACGCTGCCCACTCCAAAGCAGCCCTGGACAGCATTAAAGAGCTGCCCGA
190 200 210 220 230 240 250 260 270 280
TAACATCTCTGCTGGAGCTGTTACGACAGCTGCCCGCGCGCCAGCTGCTGCTGAACTGCCGCTGCTGCGAGCCTCTGGCGGGACCTCATCGAC
290 300 310 320 330 340 350 360 370
CTCCTGACCCCTCTGGAAGCGCAAGTGCTGCGAAAGGGCTTCATCACCAGGACTGGGACACCCCGTGGCGGACTGGAAATCTTCTACTTCC
380 390 400 410 420 430 440 450 460 470
TACGAGCCTGCATAGGAACCTCTCGCGCAACCCGTGTGCTGAAACGATATGTTTGCATGGCAAATTGATTCAATGGTGGGACCGCTGGAA
480 490 500 510 520 530 540 550 560
GGTGGATAGCTCCCTGCGCCACCGACAGAAATTTCTGACCCCAAGTCAAGAACTCTTTTGTACATCCTACGAACTGTGCTCAAGTGG
570 580 590 600 610 620 630 640 650
GAGTGTGTGGACCTTCTAGCCGACCGCTACTGGGAGGAGCTACTAGACACATTCGCGCCGACATCGTGGTTAAGGACTGGTTGTCTGCCAGAG
660 670 680 690 700 710 720 730 740 750
CCGACTGTGCTGCACCTACCAACTCAAAGTGCAGCTGGCGCTCGGCTGACTACTTGTGTGTGGCCTCTTCGAGCCCCACCTGTGACCATCCA
760 770 780 790 800 810 820 830 840
ACAGTGAACAAATGCCACATGGACAGAGGTCTCCTACACCTTCTCAGACTACCCCGGGGTGTCCGCTACATCTCTTCAGCATGGGGGCAGG
850 860 870 880 890 900 910 920 930 940
GACACCCAGTACTGGGAGGCTGGTATGGGCCCCGAGTACCAACAGCAGCATTTGCTGTAGCCCCAAGATGACCGGAACAGGCTCTCGTCCG
950 960 970 980 990 1000 1010 1020 1030
AGGCTCAGCCTGGGAGAGCATGGACAGGAGGAGGCTGCCCAATCGCCCTACGGAGCTGTGTCCAGATTTCAGACAGCTGTCCATCTGTG
1040 1050 1060 1070 1080 1090 1100 1110 1120
TCTGGGTGAGGAGGTTCTCCAGGAGGAGCTGAGCATGGGGTGGGAGTGGGTCCTGTACCAGGACTCCTGCCCGGTTCAACCCTA
1130 1140 1150 1160 1170 1180 1190 1200 1210 1220
CCAGCTTGTGGTAACTTACTGTACATAGCTCTGACGTTTGTGTGTAATAATGTTTTCAGCCCGGCACTGTGGCTCAGCCTGTAATCCAG
1230 1240 1250 1260 1270 1280 1290 1300 1310
CACTTTGGGAGACCGAGGAGGTGGATCAGGAGCTCAGGAGACAGAGACCATCTGCCCAACAGGTTGAAACCTGTCTCTACTAAAAATACAA
1320 1330 1340 1350 1360 1370 1380 1390 1400 1410
AAAAATTAGCCGGGCTGGTGGCGGGCGCTGTAGTCCAGCTACTCGGGAGGCTGATGCAGAAATGGCGTGAACCGGAAGGAGAGCTTGC
1420 1430 1440 1450 1460 1470 1480 1490 1500
AGTGAACCGAGATCAGCCACTGCAGCCTGGGTGACAGAGGAGACTGTGCTCATAAAAATAATAATAATAATAATAATAATAATA
1510 1520 1530
AATGGTTTTCAGTAAAAA

FIG. 9B

10 20 30 40 50 60
MSNTRFTITLNYKDPLTGDEETLASYGIVSGDLICLILHDDIPPPNIPSTDSEHSSLQN
70 80 90 100 110 120
NEQPSLATSSNQTSIQDEQPSDSFQGGAAQSGVWNDDSM LGPSQNFEAESIQDNAHMAEG
130 140 150 160 170 180
TGFYPSEPLLCSESVEGQVPHSLETLYQSADCS DANDALIVLIHLLMESGYIPQGTEAK
190 200 210 220 230 240
ALSLPEKWKLSGVYKLQYMHHLCEGSSATLTCVPLGNLIVVNATLKINNEIRSVKRLQLL
250 260 270 280 290 300
PESFICKEKLGENVANIYKDLQKLSRLFKDQLVYPLLAFT RQALNLPNVFGLVVLPLELK
310 320 330 340 350 360
LRIFRLLDVRSVLSLSAVCRDLFTASNDPLLWRFLYLRDFRDNTVRVQD TDWKELYRKRH
370 380 390 400 410 420
IQRKESPKGRFVLLLPSSHTIPFYPNPLHPRPFPSSRLPPGIIGGEYDQRPTLPYVGDP
430 440 450 460 470 480
ISSLIPGPGETPSQLPPLRPREDPVGPLPGPNPILPGRGGPNDRFPFRPSRGRPTDGRLS

FM

FIG. 10A

10 20 30 40 50 60 70 80 90
TGGAAATGCCATGGACCATGTCTAATACCCGATTTCATTTACATTGAACTACAAGGATCCCTCTCACTGGAGATGAAGAGACCTTGGCTTCATA
100 110 120 130 140 150 160 170 180
TGGGATGTGTTCTGGGGACTTGATATGTTTGATTCTCTCAGGATGACATTCCACCGCCTAATATACCTTCATCCACAGATTACAGGACATTCTTCA
190 200 210 220 230 240 250 260 270 280
CTCCAGAACAAATGAGCAACCTCTTTGGCCACCAGCTCCAATCAGACTAGCATACAGGATGAACAACCAAGTGATTTCATCCAGGACAGGCAG
290 300 310 320 330 340 350 360 370
CCAGTCTGGTGTGTTGGAATGACCACAGTATGTTAGGGCTAGTCAAAATTTTGAAGCTGAATCAATTCAAGATAATGGCCATATGGCAGAGGG
380 390 400 410 420 430 440 450 460 470
CACAGGTTTCTATCCCTCAGAACCCCTGCTCTGTAGTGAATCGGTGGAAGGGCAAGTCCACATTTCATTAGAGACCTTGATCAATCAGCTGAC
480 490 500 510 520 530 540 550 560
TGTCTGATGCCAATGATGCGTTGATGTTGATACATCTTCTCATGTTGGAGTCAGGTTACATACCTCAGGGCACCAGCAAGCAAGCACTGT
570 580 590 600 610 620 630 640 650
CCCTGCCGAGAAAGTGGAAAGTTGAGCGGGGTGTATAAGCTGCAGTACATGCATCATCTCTGCGAGGGCAGCTCCGCTACTCTCACCTGTGTGCC
660 670 680 690 700 710 720 730 740 750
TTTGGGAAACCTGATTGTTGTAATGCTACACTAAAAATCAACAATGAGATTAGAAGTGTGAAAGATTGCAGCTGCTACCAGAACTTTTATT
760 770 780 790 800 810 820 830 840
TGCAAGAGAGAACTAGGGGAAAATGTAGCCAAACATATACAAAGATCTTCAGAAACTCTCTCGCTCTTTAAAGACCAGCTGGTGTATCCTCTTC
850 860 870 880 890 900 910 920 930 940
TGGCTTTTACCCGACAAGCACTGAACCTACCAAAATGTATTGGGTGCTGCTCTCCATTGGAACTGAAACTACGGATCTTCCGACTTCTGGA
950 960 970 980 990 1000 1010 1020 1030
TGTCTGTTCCGTCTGTCTTTGTCTGCGGTGTTGCTGACCTCTTTACTGCTTCAAATGACCCACTCTGTGGAGGTTTTTATATCTGCGTGAT
1040 1050 1060 1070 1080 1090 1100 1110 1120
TTTCGAGACAACTACTGTCTAGATTCAAGACACAGATTGGAAAGAACTGTACAGGAAGAGGCACATACAAAGAAAGAAATCCCGAAAGGGCGGT
1130 1140 1150 1160 1170 1180 1190 1200 1210 1220
TTGTGCTGCTCTGCTGCTCAACCCACACCATTCATTCTATCCCAACCCCTTGCACCCTAGGCCATTTCCTAGCTCCCGCTTCTCTCAGG
1230 1240 1250 1260 1270 1280 1290 1300 1310
AATTATCGGGGGTGAATATGACCAAGACCAACACTTCCCTATGTTGGAGACCAATCAGTTCACTCATTCCTGGTCTGGGGAGACCCCGAGC
1320 1330 1340 1350 1360 1370 1380 1390 1400 1410
CAGTTACCTCCACTGAGACCACGCTTTGATCCAGTTGGCCCACTTCCAGGACCTAACCCCATCTTGCCAGGGCGAGGGGCCCCAATGACAGAT
1420 1430 1440 1450 1460 1470 1480 1490 1500
TTCCCTTTAGACCCAGCAGGGGTGGGCAACTGATGGCGCTGTCAATTCATGTCATGATTGTAATTTCAATTTCTGGAGCTCCATTGTTTT
1510 1520 1530 1540 1550 1560 1570 1580 1590
TGTTTCTAAACTACAGATGTCATCTCTGGGGTGTGATCTCGAGTGTATTCTGATTGCTGTTGAGAGTTGCACTCCAGAAACCTTTT
1600 1610 1620 1630 1640 1650 1660 1670 1680 1690
AAGAGATACATTTATAGCCCTAGGGGTGGTATGACCAAGGTTCTCTGTGACAAGGTGGGCTTGGGAATAGTTGGCTGCCAATCTCCCTGC
1700 1710 1720 1730 1740 1750 1760
TCTTGGTTCTCTCTAGATTGAAGTTTGTGTTCTGATGCTGTTCTTACCAGATTAAGAAAAAGTGTAAATT

FIG. 10B

10 20 30 40 50 60
ETSKLG*SAVLAPAAGGTLSEGRSAVSGILIAVTSTGVDK*SLNQLLHGLGTSSRLSHF

70 80 90 100 110 120
PFG*KSPPRGQFVAAAVEIAGRSGLQMGQGLWRVVRNQQLQEGYSEQGYLTREQSRMA

130 140 150 160 170 180
ASNISNTNHRKQVQGGIDYHLLKARKSKEQEGFINLEMLPPELSFTILSYLNATDLCLA

190 200 210 220 230 240
SCVWQDLANDELLWQGLCKSTWGHCSIYNKNPPLGFSFRKXYMQLDEGSLTFNANPDEGV

250 260 270 280 290 300
NYFMSKGILDDSPKEIAKFIFCTRTLNWKKLRIYLDERRDVLDDLVTLHNFNRNQFLPNAL

310 320 330 340 350 360
REFFRHIHAPEERGEYLETLITKFSHRFCACNPDLRELGLSPDAVYVLCYSLILLSIDL

370 380 390 400 410 420
TSPHVKNKMSKREFIRNTRAAQNISEDFVGHLYDNIYLIGHVAA*KAQLLGLQFLLQTK

430 440 450 460 470 480
ATQGLSRYGGYISAGHCSSLISQSSFSVQPFFLLPFSILVISLGN*IILQNFS*FCLSRFA

490 500 510 520 530 540
QSRATV*HSC*RMIN*HYTLKDGVEVH*ICLKNFIHFHSLYKYHVMCTYLTKEIYSHNYF

550 560 570 580 590 600
IVKIILTKVFPFLSN*VLKFI*F*SETIVXVKVRSDFRQKPIPASFSFKL*RVLICYYITM

610 620 630 640 650
QNWQLFL*YKFII*FFILKTGLIKSR*VL*TI*DF*NIKIYDLHS*E*NKIXLELW

FIG. 11A

10 20 30 40 50 60 70 80 90
GGAAACGTCAAAATTTGGGATAGTCGGCAGTTCTGGGCCCTGCAGCTGGAGGTACCCCTGAGTTCTGAGGTCGTAGTCTCTTTCTGCTATTCTTC
100 110 120 130 140 150 160 170 180
ATCGCGGTCACTCTTACCGGTGTGGACAAGTAAAGTTTGAATCAGCTTCTCCATGGCCTGGGCACCAGTTCCCGGCTGAGCCATTTTCTTTTTC
190 200 210 220 230 240 250 260 270 280
GCTAAAGTCCCGCCAGAGGCCAATTGCTCGCGCGCGGTGGAGATCGCAGGTGCTCAGGCTTGCAGATGGGTCAAGGGTTGTGGAGAGT
290 300 310 320 330 340 350 360 370
GGTCAGAAACAGCAGCTGCAACAAGAGGCTACAGTCAGCAAGGCTACCTCACCAGAGAGCAGAGCAGGAGAATGGCTGGCAGCAACATTTCT
380 390 400 410 420 430 440 450 460 470
AACACCAATCATCGTAAACAAGTCCAAGGAGGCAATTGACATATATCATCTTTTGAAGGCAAGGAATCGAAAGAACAGGAAGGATTTCATTAATT
480 490 500 510 520 530 540 550 560
TGGAAATGTTGGCTCTGAGCTAAGCTTTACCATCTTGTCTACCTGAATGCAACTGACCTTTGCTTGGCTTCATGTGTTTGGCAGGACCTTGC
570 580 590 600 610 620 630 640 650
GAATGATGAATCTCTCGCAAGGGTTGTGCAAAATCCACTTGGGCTCACTGTTCCATATACAATAAGAACCCACCTTTAGGATTTTCTTTTAGA
660 670 680 690 700 710 720 730 740 750
AAATGTATATGCACTGCGATGAAGGCAGCCTCACTTTAATGCCAACCCAGATGAGGAGTGAACACTTTTATGTCCAAGGGTATCCTGGATG
760 770 780 790 800 810 820 830 840
ATTCCGCCAAAGAAATAGCAAGTTTATCTTCTGTACAAGAACTAAATTGGAAAAAAGTGAATCTATCTTGTGAAGGAGAGATGTCTT
850 860 870 880 890 900 910 920 930 940
GGATGACCTTGTAACTTGCATAATTTTAGAAATCAGTTCTTGCACAAATGCACTGAGAGAATTTTTTGTCTATCCATGCCCTGAAGAGCGT
950 960 970 980 990 1000 1010 1020 1030
GGAGAGTATCTTGAACCTTTATAACAAGTTCTCAGATAGATTCTGTGCTTGCACCCCTGATTTAATGCGAGAACTTGGCCTTACTCTGTATG
1040 1050 1060 1070 1080 1090 1100 1110 1120
CTGTCTATGTACTTGTCTACTCTTTGATTCTACTTTCCATTGACCTCACTAGCCCTCATGTGAAGAATAAAATGTCAAAAAGGGAATTTATTTC
1130 1140 1150 1160 1170 1180 1190 1200 1210 1220
AAATACCCGTGCGGCTGCTCAAAATATTAGTGAAGATTTGTAGGGCATCTTTATGACAATATCTACCTTATTGGCCATGTGGCTGCATAAAAA
1230 1240 1250 1260 1270 1280 1290 1300 1310
GCACAATGCTAGGACTTCAGTTTTTACTTTCAGACTAAAGCTACCCAGGACTTACAGATATCGGGCTTACATCAGTGTGGTCAATTGTAGCC
1320 1330 1340 1350 1360 1370 1380 1390 1400 1410
TGAGTATACAAATCAAGCTTCAGTGTGCAACCTTTTTTTCTTTTGGCATTCTTATTTTAGTAATTTCTTGGGGAACATAAATTTTGCAGAA
1420 1430 1440 1450 1460 1470 1480 1490 1500
TTTTTCTTAATTTTGTATTATCAGTTTTCACAAACAGAGCCACTGTCTAACACAGCTGTTAACGAATGATAAACTGACATTATACTCTAAAA
1510 1520 1530 1540 1550 1560 1570 1580 1590
GATGCTGATTTTGTGATTAGATTGCTGAAAACTTTATCCATTTCATTCTTATACAAATACCATGTAATGTGTACATATTTAACTAAAG
1600 1610 1620 1630 1640 1650 1660 1670 1680 1690
AGATTTATAGTCATAATTATTTTATGTAAAGATTTAACTAAAGTTTTCCTTTCTCTCAAACTGAGTTCTGAAATTTATTTGATTCTGATC
1700 1710 1720 1730 1740 1750 1760 1770 1780
TCAAACTATTCTCTTCGTAAGTTAGATCTGACTTCAGRCAGAAACCAATACCAGCTTCTTTCTTTTAACTTTGAAGAGTGTGATTGT
1790 1800 1810 1820 1830 1840 1850 1860 1870 1880
TACTATATTACTATGCAAACTGGCAGTTATTTTATAATATAAATTTATAATTGATTTTTTATTTTAAAACTGGCTTAATCAAGTCTCGGT
1890 1900 1910 1920 1930 1940 1950 1960 1970
AAGTCTTTTAAACCTTTAGGATTTTAAACATCAAAATTTATGATTTACATTATAGGAATAAAATATAATATTAGAACTCTGGT

FIG. 11B

10 20 30 40 50 60
MAAAVDSAMEVVPALAEAAPEVAGLSCLVNLPGEVLEYILCCGSLTAADIGRVSSSTR

70 80 90 100 110 120
RLRELCSGSKVWKEQFRVRWPSLMKHYSPTDYVNWLEEKVRQKAGLEARKIVASFSCR

130 140 150 160 170 180
FFSEHVPCNGFSDIENLEGPEIFFEDELVCILNMEGRKALTWKYYAKKILYYLRQQKILN

190 200 210 220 230 240
NLKAFLLQPPDDYESYLEGAVYIDQYCNPLSDISLKDIQAQIDSIVELVCKTLRGINSRHP

250 260 270 280 290 300
SLAFKAGESSMIMEIELQSQVLDAMNYVLYDQLKFKGNRMDYYNALNLYMHQVLRRTGI

310 320 330 340 350 360
PISMSLLYLTARQLGVPLEPVNFP SHFLLRWCQGAEGATLDIFDYIYIDAFGKGKQLTV

370 380 390 400 410 420
KECEYLIGQHVTAAALYGVVNVKKVLQRMVGNLLSLGKREGIDQSYQLLRDSL DLYLAMYP

430 440 450 460 470 480
DQVQLLLLQARLYFHLGIWPEKVL DILQHIQTLDPGQHGA VG YLVQHTLEHIERKKEEVG

490 500 510 520 530 540
VEVKLRSDDEKHRDVCYSIGLIMKHKRYGYN CVIYGWDPTCMMGHEWIRNMNVHSLPHGHH

550 560 570 580 590 600
QPFYNVLVEDGSCRYAAQENLEYNVEPQEISHPDVGRYFSEFTGTHYIPNAELEIRYPED

610 620
LEFVYETVQNIYSAKKENIDE

FIG. 12A

FIG. 12B

10	20	30	40	50	60
RSTGFRRAGEEWSR*XLAASPGXLRRPAXTFVLSNLAEVVERVLTFLPAKALLRVACVCR					
70	80	90			
LWRECVRRVLRTHRSVTWISAGLAEAGHLXGH					

FIG. 13A

10 20 30 40 50 60
CCGTAGTACTGGNTTCCGGCGGGCTGGTGAGGAATGGAGCCGGTAGNTGCTTGCGGCGAG
70 80 90 100 110 120
TCCCGGGNTCCTCCGTAGACCCGCGGANACCTTCGTGTTGAGTAACCTGGCGGAGGTGGT
130 140 150 160 170 180
GGAGCGTGTGCTCACCTTCCTGCCCCCAAGGCGTTGCTGCGGGTGGCCTGCGTGTGCCG
190 200 210 220 230 240
CTTATGGAGGGAGTGTGTGCGCAGAGTATTGCGGACCCATCGGAGCGTAACCTGGATCTC
250 260 270
CGCAGGCCTGGCGGAGGCCGCCACCTGGNGGGGCATT

FIG. 13B

10 20 30 40 50 60
RPRPVQQQQQQPPQQPPPPQQPPQQPPPPPPQQQQQQPPPPPPPPPLPQERNNVG

70 80 90 100 110 120
ERDDDVPAADMVAEESGPGAQNSPYQLRRKTL LPKRTACPTKNSMEGASTSTTENFGHRAK

130 140 150 160 170 180
RARVSGKSQDL SAAPAEQYLQEKL PDEVVLKIFSYLLEQDLCRAACVCKRFSELANDPNL

190
WKRLYMEVF EYTRPMMH

FIG. 14A

10 20 30 40 50 60
GCGGCCGCGCCCGGTGCAGCAACAGCAGCAGCAGCCCCCGCAGCAGCCCGCCCGCAGCC

70 80 90 100 110 120
GCCCCAGCAGCAGCCGCCCCAGCAGCAGCCTCCGCCCGCCGCGCAGCAGCAGCAGCA

130 140 150 160 170 180
GCAGCCTCCGCCCGCCGCCACCGCCGCTCCGCCGCTGCCTCAGGAGCGGAACAACGTCCG

190 200 210 220 230 240
CGAGCGGGATGATGATGTGCCTGCAGATATGGTTGCAGAAGAATCAGGTCCTGGTGCACA

250 260 270 280 290 300
AAATAGTCCATACCAACTTCGTAGAAAACTCTTTTGCCGAAAAGAACAGCGTGTCCCAC

310 320 330 340 350 360
AAAGAACAGTATGGAGGGCGCCTCAACTTCAACTACAGAAAACCTTGGTCATCGTGCAAA

370 380 390 400 410 420
ACGTGCAAGAGTGTCTGGAAAATCACAAGATCTATCAGCAGCACCTGCTGAACAGTATCT

430 440 450 460 470 480
TCAGGAGAAACTGCCAGATGAAGTGGTTCTAAAAATCTTCTCTTACTTGCTGGAACAGGA

490 500 510 520 530 540
TCTTTGTAGAGCAGCTTGTGTATGTAAACGCTTCAGTGAACCTTGCTAATGATCCCAATTT

550 560 570 580 590
GTGGAACGATTATATATGGAAGTATTTGAATATACTCGCCCTATGATGCAT

FIG. 14B

10 20 30 40 50 60
RPRPGLRGGRAPCEVTMEAGGLPLELWRMILAYLHLPDLGRCSLVCRAWYELILSLDSTR
70 80 90 100 110 120
WRQLCLGCTECHRPNWPNQPDVEPESWREAFKQHYLASKTWTKNALDLESSICFSLFRRR
130 140 150 160 170
RERRTL SVGP GREFD SLGSALAMASLYDRIVLFPGVYEEQGEIILKVPVEIVGQGLG

FIG. 15A

10 20 30 40 50 60
GCGGCCGCGGCCCGGACTCCGCGGTGGGCGAGCGCCCTGTGAGGTGACCATGGAGGCTGG
70 80 90 100 110 120
TGGCCTCCCTTGGAGCTGTGGCGCATGATCTTAGCCTACTTGCACCTTCCCGACCTGGG
130 140 150 160 170 180
CCGCTGCAGCCTGGTATGCAGGGCCTGGTATGAACTGATCCTCAGTCTCGACAGCACCCG
190 200 210 220 230 240
CTGGCGGCAGCTGTGTCTGGGTGACCCGAGTGCCGCCATCCCAATTGGCCCAACCAGCC
250 260 270 280 290 300
AGATGTGGAGCCTGAGTCTTGGAGAGAAGCCTTCAAGCAGCATTACCTTGCATCCAAGAC
310 320 330 340 350 360
ATGGACCAAGAATGCCTTGGACTTGGAGTCTTCCATCTGCTTTTCTCTATTCCGCCGGAG
370 380 390 400 410 420
GAGGGAACGACGTACCCTGAGTGTGGGCCAGGCCGTGAGTTTGACAGCCTGGGCAGTGC
430 440 450 460 470 480
CTTGGCCATGGCCAGCCTGTATGACCGAATTGTGCTCTTCCCAGGTGTGTACGAAGAGCA
490 500 510 520 530
AGGTGAAATCATCTTGAAGGTGCCTGTGGAGATTGTAGGGCAGGGGAAGTTGGGTGA

FIG. 15B

10 20 30 40 50 60
ETETAPLTLES LPTDPLLLILSF LDYRDLINCCYVSRRLSQLSSH DPLWRRHCKKYWLIS

70 80 90 100 110 120
EEETQKNQCWKS LFDITYSDVGRYIDHYAAIKKASGMISRNIWSPGV LGWVLSLKEGCS

130 140 150 160 170 180
RGRPRCCGSADWAASFLDDYRCSYRIHNGQKL VGSGYWEAWHCLITIVLKIC*TSIQLP

190 200 210 220 230 240
EIPAETGTEILSPFNFCIHTGLSQYIAVEAAEG*NKNEVFYQCQTVERVFKYGIKMCSDG

250
CINGMH*VFS

FIG. 16A

10 20 30 40 50 60
GAGACCGAGACGGCGCCGCTGACCCCTAGAGTCGCTGCCACCGATCCCCTGCTCCTCATC
70 80 90 100 110 120
TTATCCTTTTTTGGACTATCGGGATCTAATCAACTGTTGTTATGTCAGTCGAAGATTAAGC
130 140 150 160 170 180
CAGCTATCAAGTCATGATCCGCTGTGGAGAAGACATTGCAAAAAATACTGGCTGATATCT
190 200 210 220 230 240
GAGGAAGAGAAAACACAGAAGAATCAGTGTGGAAATCTCTCTTCATAGATACTTACTCT
250 260 270 280 290 300
GATGTAGGAAGATACATTGACCATTATGCTGCTATTAAAAAGGCCCTCGGGAATGATCTCA
310 320 330 340 350 360
AGAAATATTTGGAGCCCAGGTGTCCTCGGATGGGTTTTATCTCTGAAAGAGGGGTGCTCG
370 380 390 400 410 420
AGAGGAAGACCTCGATGCTGTGGAAGCGCAGATTGGGCTGCAAGTTTCCTGGACGATTAT
430 440 450 460 470 480
CGATGTTTCATACCGAATTCACAATGGACAGAAGTTAGTTGGTTCTGGGGTTATTGGGAA
490 500 510 520 530 540
GCATGGCACTGTCTAATCACTATCGTTCTGAAGATTGTGTTAGACGTCGATACAGCTGCCG
550 560 570 580 590 600
GAGATTCCAGCAGAGACAGGGACTGAAATACTGTCTCCCTTTAACTTTTGACATACATACT
610 620 630 640 650 660
GGTTTGAGTCAGTACATAGCAGTGGAAGCTGCAGAGGGTTGAAACAAAAATGAAGTTTTC
670 680 690 700 710 720
TACCAATGTCAGACAGTAGAACGTGTGTTTAAATATGGCATTAAAGATGTGTTCTGATGGT
730 740 750
TGTATAAATGGCATGCATTAGGTATTTTCAG

FIG. 16B

WO 02/055665

31/80

PCT/US02/00311

10 20 30 40 50 60
GSGFRAGGWPLTMPGKHQHFQEPVGGCCGKYFLFGFNIVFWVLGALFLAIGLWAWGEKGV
70 80 90 100 110 120
LSNISALTDLGGLDPVWLVCGSWRRHVGAGLCWAAIGALRENTFLLKFFXXFLGLIFFLE
LA

FIG. 17A

10 20 30 40 50 60
GGCTCCGGTTTCCGGGCCGGCGGGTGGCCGCTCACCATGCCCGGNAAGCACCAGCATTTC

70 80 90 100 110 120
CAGGAACCTGAGGTCGGCTGCTGCGGGAAATACTTCCTGTTTGGCTTCAACATTGTCTTC

130 140 150 160 170 180
TGGGTGCTGGGAGCCCTGTTCTCTGGCTATCGGCCTCTGGGCCTGGGGTGAGAAGGGCGTT

190 200 210 220 230 240
CTCTCGAACATCTCAGCGCTGACAGATCTGGGAGGCCTTGACCCCGTGTGGCTTGTTTGT

250 260 270 280 290 300
GGTAGTTGGAGGCGTCATGTCGGTGCTGGGCTTTGCTGGGCTGCAATTGGGGCCCTCCGG

310 320 330 340 350 360
GAGAACACCTTCCTGCTCAAGTTTTTCTNCGNSTTCCTCGGTCTCATCTTCTTCCTGGAG

CTGGCAAC

FIG. 17B

10 20 30 40 50 60
AAAAAYLDELPEPLLLRVLAALPAAELVQACRLVCLRWKELVDGAPLWLLKCQQEGLVP
70 80 90 100 110 120
EGGVEEERDHWQQFYFLSKRRRNLLRNPCGEEDLEGWCDVEHGGDGWRVEELPGDSGVEF
130 140 150 160 170 180
THDESVKKYFASSFEWCRKAQVIDLQAEGYWEELDDTTQPAIVVKDWYSGRSDAGCLYEL
190 200 210 220 230 240
TVKLLSEHENVLAEFSSGQVAVPQSDGGGWMEISHTFTDYGPGVRFVRFEHGGQGSVYW
250
KGWFGARVTNSSVWVEP*

FIG. 18A

10 20 30 40 50 60
GCGGCGGCCCGCGCGCTACCTGGACGAGCTGCCCGAGCGCTGCTGCGCGTCTGGCCGCACTG
70 80 90 100 110 120 130
CCGGCCCGCGAGCTGGTGCAGGCCTGCCGCTGCTGCTGCGCTGGAAGGAGCTGGTGGACGCGGCC
140 150 160 170 180 190 200
CCGCTGTGGCTGCTCAAGTGCCAGCAGGAGGGGCTGGTGCCCGAGGGCGCGTGGAGGAGGAGCGCGAC
210 220 230 240 250 260 270
CACTGGCAGCAGTTCTACTTCCTGAGCAAGCGCGCCGCAACCTTCTGCGTAACCCGTGTGGGAAGAG
280 290 300 310 320 330 340
GACTTGAAGGCTGGTGTGACGTGGAGCATGGTGGGGACGGCTGGAGGGTGGAGGAGCTGCCTGGAGAC
350 360 370 380 390 400 410
AGTGGGGTGGAGTTCACCCACGATGAGAGCGTCAAGAAGTACTTCGCCCTCCTCTTGTAGTGGTGTCCG
420 430 440 450 460 470 480
AAAGCACAGGTCAATTGACCTGCAGGCTGAGGGCTACTGGGAGGAGCTGCTGGACACGACTCAGCCGGCC
490 500 510 520 530 540 550
ATCGTGGTGAAGGACTGGTACTCGGGCCGCGAGCGCTGGTTGCCCTCTACGAGCTCACCCTTAAGCTA
560 570 580 590 600 610 620
CTGTCCGAGCACGAGAACGTGCTGGCTGAGTTCAAGCAGCGGGCAGGTGGCAGTGCCCCAAGACAGTGAC
630 640 650 660 670 680 690
GGCGGGGGCTGGATGGAGATCTCCACACCTTCACCGACTACGGGGCCGGCGTCCGCTTCGTCCGCTTC
700 710 720 730 740 750
GAGCACGGGGGGCAGGGCTCCGTCTACTGGAAGGGCTGGTTCGGGGCCCGGGTGACCAACAGCAGCGTG
760 770
TGGGTAGAACCCTGA

FIG. 18B

10 20 30 40 50 60
MGEKAVPLLRRRRVKRSCPCSGSELGVEEKRGKGNPISIQLFPPPELVEHIISFLPVRDLV

70 80 90 100 110 120
ALGQTCRYFHEVCDGEGVWRRICRRLSPRLQDQDTKGLYFQAFGGRRRCLSKSVAPLLAH

130 140 150 160 170 180
GYRRFIPTKDHVFILDYVGTLLFFLKNALVSTLGQMOWKRACRYVVLRCGAKDFASDPRCD

190 200 210 220 230 240
TVYRKLYVLATREPQEVVGTSSRACDCVEVYLQSSGQRVFKMTFHHSMTFKQIVLVGQ

250 260 270 280 290 300
ETQRALLLLTEEGKIYSLVVNETQLDQPRSYTVQLALRKVSHYLPFLRVACMTSNQSSTL

310
YVTDPILCSWLQPPWPGG

FIG. 19A

10 20 30 40 50 60
ATGGGGCAGAAGGCGGTCCCTTTGCTAAGGAGGAGGCGGGTGAAGAGAAGCTGCCCTTCTGTGGCTCG
70 80 90 100 110 120 130
GAGCTTGGGGTTGAAGAGAAGAGGGGAAAGGAAATCCGATTTCATCCAGTTGTTCCCCCAGAGCTG
140 150 160 170 180 190 200
GTGGAGCATATCATCTCATTCCTCCAGTCAGAGACCTTGTTGCCCTCGGCCAGACCTGCCGCTACTTC
210 220 230 240 250 260 270
CACGAAGTGTGCGATGGGGAAGGCGTGTGGAGACGCATCTGTGCGAGACTCAGTCCGCGCCTCCAAGAT
280 290 300 310 320 330 340
CAGGACACGAAGGGCCTGTATTTCCAGGCATTGAGAGGCCCGCCGATGTCTCAGCAAGAGCGTGGCC
350 360 370 380 390 400 410
CCCTTGCTAGCCACGGCTACCGCCGCTTCTTGCCCAAGGATCAGTCTTCATTCTTGACTACGTG
420 430 440 450 460 470 480
GGGACCTCTTCTCCTCAAAAATGCCCTGGTCTCCACCCTCGGCCAGATGCAGTGAAGCGGGCCTGT
490 500 510 520 530 540 550
CGCTATGTTGTGTTGTGTCGTGGAGCCAAGGATTTGCCTCGGACCCAAGGTGTGACACAGTTTACCGT
560 570 580 590 600 610 620
AAATACCTCTACGTCTTGGCCACTCGGGAGCCGCAGGAAGTGGTGGGTACCACCAGCAGCCGGGCCTGT
630 640 650 660 670 680 690
GACTGTGTTGAGGTCTATCTGCAGTCTAGTGGGCAGCGGTCTTCAAGATGACATTCCACCACTCAATG
700 710 720 730 740 750
ACCTTCAAGCAGATCGTGCTGGTGGTGCAGGAGACCCAGCGGGCTCTACTGCTCCTCACAGAGGAAGGA
760 770 780 790 800 810 820
AAGATCTACTCTTTGGTAGTGAATGAGACCCAGCTTGACCAGCCACGCTCCTACAGGTTTCAGCTGGCC
830 840 850 860 870 880 890
CTGAGGAAGGTGTCCCACTACCTGCCTCACCTGCGCGTGGCCTGCATGACTTCCAACCAGAGCAGCACC
900 910 920 930 940 950
CTCTACGTACAGATCCTATCTGTGCTCTTGGCTACAACCACCTTGGCCTGGTGGATGA

FIG. 19B

10 20 30 40 50 60
RGGSEGRGRGREKRARGARRKRKQGGREARAADGEGGSGPGAEGARTRPREEAEGGGSV
70 80 90 100 110 120
EEGARGIIKGDEGSVGAGKEAQGRKYGKEEWRVRARRREGARPGRVQQGGQVWAYIPGT
130 140 150 160 170 180
GAAMAAAAREEEEEEAARESAACPAAGPALWRLPEVLLHMC SYLDMRALGR LAQVYRWLW
190 200 210 220 230 240
HFTNCDLLRRQIAWASLNSGFTRLGTNLMTSVPVKVSQNWIVGCCREGILLKWRC SQMPW
250 260 270 280 290 300
MQLEDDALYISQANFILAYQFRPDGASLNRQPLGVSAGHDEDVCHFVLATSHIVSAGGDG
310 320 330 340 350 360
KIGLGKIHSTFAAKYWAHEQEVNVCVCKGGIISFGSRDRTAKVWPLASGQLGQCLYTIQT
370 380 390 400 410 420
EDQIWSVAIRPLLSSFVTGTACCGHFSPLKIWDLNSGQLMTHLDRDFPPRAGVLDVIYES
430 440 450 460 470 480
PFALLSCGYDTYVRYWDCRTSVRKCVMEWEEPHNSTLYCLQTDGNHLLATGSSSFYSVVRL
490 500 510 520 530
WDRHQACPHTFPLTSTRLGSPVYCLHLTTKHLAALSYNLHVLDIQNP*

FIG. 20A

FIG. 20B

LILTSVLLFQRHGYCTLGEAFNRDLDFSSAIQDIRTFNYVVKLLQLIAKSQLTSLSGVAQK
70 80 90 100 110 120
NYFNILDKIVQKVLDDHHNPRLIKDLLQDLSSTLCILIRGVGKSVLVGNINIWICRLETI
130 140 150 160 170 180
LAWQQQLQDLQMTKQVNNGLTSLDLPLHMLNNILYRFS DGWDIITLGQVTPTLYMLSEDR
190 200 210 220 230 240
QLWKKLCQYHFAEKQFCRHLILSEKGHIEWKLMYFALQKHYPAKEQYGDTLHFCHCSIL
250 260 270
FWKDSGHPCTAADPDSCFTPVSPQHFDLDFKF

FIG. 21A

10 20 30 40 50 60
GCATTGCTATAATTTTACTATACTCTCATCTAAATCTAAAATCAGTCTTCAAAATAAAAACAAATTGTC
70 80 90 100 110 120 130
CTTGCCAAAAATTTTAAATCGCACAAATTAATTGACATTAAGTCCAATTCTTTTGGCTAATTGAC
140 150 160 170 180 190 200
TAATTTTAACTTCTGTGTGCTTTTCCAGAGGCATGGCTATTGCACCTTGGGAGAAGCCTTTAATCGGT
210 220 230 240 250 260 270
TAGACTTCTCAAGTGCAATTCAAGATATCCGAACGTTCAATTATGTGGTCAAACGTTGCAGCTAATTG
280 290 300 310 320 330 340
CAAAATCCAGTTAACTTCATTGAGTGGCGTGGCACAGAAGAATTACTTCAACATTTTGGATAAAATCG
350 360 370 380 390 400 410
TTCAAAAGGTTCTTGATGACCACCACAATCCTCGCTTAATCAAAGATCTTCTGCAAGACCTAAGCTCTA
420 430 440 450 460 470 480
CCCTCTGCATTCTTATTAGAGGAGTAGGGAAGTCTGTATTAGTGGGAAACATCAATATTGGATTGGCC
490 500 510 520 530 540 550
GATTAGAACTATTCTCGCCTGGCAACAACAGCTACAGGATCTTCAGATGACTAAGCAAGTGAACAATG
560 570 580 590 600 610 620
GCCTCACCCCTCAGTGACCTTCCTCTGCACATGCTGAACACATCCTATACCGGTTCTCAGACGGATGGG
630 640 650 660 670 680 690
ACATCATCACCTTAGGCCAGGTGACCCCAAGTGTATATGCTTAGTGAAGACAGACAGCTGTGGAAGA
700 710 720 730 740 750
AGCTTTGTCTAGTACCATTTTGTCTGAAAAGCAGTTTGTAGACATTTGATCCTTTAGAAAAAGGTCATA
760 770 780 790 800 810 820
TTGAATGGAAGTTGATGTACTTTGCACTTCAGAAACATTACCCAGCGAAGGAGCAGTACGGAGACACAC
830 840 850 860 870 880 890
TGCATTTCTGTGGCACTGCAGCATTCTCTTTTGAAGGACTCAGGACAQCCCTGCACGGCGGCCGACC
900 910 920 930 940 950 960
CTGACAGCTGCTTCACGCCCTGTGTCTCCGCAGCACTTCATCGACCTCTTCAAGTTTTAAGGGCTGCCCC
970 980 990 1000 1010 1020 1030
TGCCATCCCTATTGGAGATTGTGAATCCTGTGTCTGTGCAGGGCTCATAGTGAGTGTCTGTGAGGTG
1040 1050 1060 1070 1080 1090 1100
GGTGAGACTCCTCGGAAGCCCTGCTTCCAGAAAGCCTGGGAAGAACTGCCCTTCTGCAAAGGGGGGA
1110 1120 1130 1140 1150 1160 1170
CTGCATGGTTGCATTTTCATCACTGAAAGTCAGAGGCCAAGGAATCATTTCTACTTCTTTAAAACTC
1180 1190 1200 1210
CTTCTAAGCATATTAAAAATGTGAAATTTTGGCTACTCTCTC

FIG. 21B

10	20	30	40	50	60
YGSEGGKSSSISSDVSSSTDHTPTKAQKNVATSESDSLSMRTLSTPSPALICPPNLPGFQ					
70	80	90	100	110	120
NGRGSSSTSSSITGETVAMVHSPPPTRLTHPLIRLASRPQKEQASIDRLPDHSMVQIFSF					
130	140	150	160	170	180
LPTNQLCRCARVCRRWYNLAWDPRWLWRTIRLTGETINVDRALKVLTRRLCQDTPNVCLML					
190	200	210	220	230	240
ETVTVSGCRRLTDGRGLYTIAQCCPELRRLEVSGCYNISNEAVFDVVS LCPNLEHLDVSGC					
250	260	270	280	290	300
SKVTCISLTREASIKLSPLHGKQISIRYLDMTDCFVLEDEGLHTIAAHCTQLTHLYLRRC					
310	320	330	340	350	360
VRLTDEGLRYLVIIYCASIKELSVSDCRFVSDFGLREIAKLESRLRYLSIAHCGRVTDVGI					
370	380	390	400	410	420
RYVAKYCSKRLRYLNARGCEGITDHGVEYLAKNCTKLKSLDIGKCPLVSDTGLECLALNCF					
430	440	450	460	470	480
NLKRLSLKSCESITGQGLQIVAANCFDLQTLNVQDCEVSVEALRFVVKRHCKRCVIEHTNP					

AFF

FIG. 22A

FIG. 22B

10 20 30 40 50 60
AAAPAPAPAPTPTPEEGPDAGWGDRIPLEILVQIFGLLVAADGMPFPGAARVCRRWQE

70 80 90 100 110 120
AASQPALWHTVTLSSPLVGRPAKGGVKAEEKLLASLEWLMPNRFSQLQRLTLHWKSQVH

130 140 150 160 170 180
PVLKLVGECCPRLTFLKLSGCHGVTADALVMLAKACQLHSLDLQHSMVESTAVVSFLEE

190 200 210 220 230 240
AGSRMRKLWLTYSQTTAILGALLGCCPQLQVLEVSTGINRNSIPLQLPVEALQKGCPQ

250 260 270 280
LQVLRLLNLMWLKPPGRGVAPGPGFPSLEELCLASSTCNFVS

FIG. 23A

10 20 30 40 50 60
TGCGGCGCGCGCCGACCCGACCCGGCACCCACGCCCGAGGAAGGGCCGACGCGGGCTGGGG
70 80 90 100 110 120 130
AGACCGCATTCCCTTGGAATCCTGGTGCAGATTTCGGGTGTTGGTGGCGCGACGCGCCCATGCC
140 150 160 170 180 190 200
CTTCCTGGGCAGGGCTGCGCGGTGTGCGCCGCTGGCAGGAGGCGCTTCCCAACCCGCGCTCTGGCA
210 220 230 240 250 260 270
CACCGTGACCCTGTCTGCCCGCTGGTGGCCGCGCTGCCAAGGGCGGGGTCAAGGCGGAGAAGAAGCT
280 290 300 310 320 330 340
CCTTGCTTCCTGGAGTGGCTTATGCCCAATCGGTTTTTCACAGCTCCAGAGGCTGACCCTCATCCACTG
350 360 370 380 390 400 410
GAAGTCTCAGGTACACCCCGTGTGAAGCTGGTAGGTGAGTGCTGTCTCGGCTCACTTTCCTCAAGCT
420 430 440 450 460 470 480
CTCCGGCTGCCACGGTGTGACTGCTGACGCTCTGGTCATGCTAGCCAAAGCCTGCTGCCAGCTCCATAG
490 500 510 520 530 540 550
CCTGGACCTACAGCACTCCATGGTGGAGTCCACAGCTGTGGTGAGCTTCTTGGAGGAGGCAGGGTCCCG
560 570 580 590 600 610 620
AATGCGCAAGTTGTGGCTGACCTACAGCTCCCAGACGACAGCCATCCTGGGCGCATTGCTGGGCAGCTG
630 640 650 660 670 680 690
CTGCCCCCAGCTCCAGGTCTGGAGGTGAGCACCGGCATCAACCGTAATAGCATTCCTTCAGCTGCC
700 710 720 730 740 750
TGTGAGGCTCTGCAGAAAGGCTGCCCTCAGCTCCAGGTGCTGCGGCTGTGAACCTGATGTGGCTGCC
760 770 780 790 800 810 820
CAAGCCTCCGGGACGAGGGGTGGCTCCCGGACCAGGCTTCCCTAGCCTAGAGGAGCTCTGCCTGGCGAG
830 840 850
CTCAACCTGCAACTTTGTGAGC

FIG. 23B

10 20 30 40 50 60
QHCSQKDTAELLRGLSLWNHAEERQKFFKYSVDEKSDKEAEVSEHSTGITHLPPEVMLSI
70 80 90 100 110 120
FSYLN PQELCRCSQVSMKWSQLTKTGSLWKHLYPVHWARGDWYSGPATELDTPEDDWVK
130 140 150 160 170 180
NRKDESRAPEHWEDEDADIDEESESAEESIAISIAQMEKRLHGLIHNVL PYVGT SVKTLV
190 200 210 220 230 240
LAYSSAVSSKMVRQILELCPNLEHLDLTQTDISDSAFDSWSWLGCCQSLRHLDSLGCCKI
250 260 270 280 290 300
TDVALEKISRALGILTSHQSGFLKTSTSKITSTAWKNKDITMQSTKQYACLHDLTNKGIG
310 320 330 340 350 360
EEIDNEHPWTKPVSSSENFTSPYVWMLDAEDLADIEDTVEWRHRNVESLCVMETASNFSCS
370 380 390 400 410 420
TSGCFSKDIVGLRTSVCWQQHCASPAFAYCGHSFCCTGTALRTMSSLPESSAMCRKAART
430 440 450 460 470 480
RLPRGKDLIYFGSEKSDQETGRVLLFLSLSGCYQITDHGLRVLTGGGLPYLEHLNLSCG
490 500 510 520 530 540
LTITGAGLQDLVSACPSLNDEYFYCDNINGPHADTASGCQNLCGFRACCRSGE*PLTS
550 560 570 580 590
DLCLHLAEQAFFHALYS*HISCVNHPFLSVTCFGPIXYNFRNLNYQXIVML

FIG. 24A

10 20 30 40 50 60 70 80 90
ACAACACTGCTCTCAGAAAGGATACTGCAGAACTCCTTAGAGGTCTTAGCCTATGGAATCATGCTGAAGAGCGACAGAAATTTTTAAATATTC
100 110 120 130 140 150 160 170 180
GTGGATGAAAAAGTCAGATAAAGAAGCAGAAGTGTGAGAACTCCACAGGTATAACCCATCTTCTCTCAGGTAATGCTGTCAATTTTCAGCT
190 200 210 220 230 240 250 260 270 280
ATCTTAATCCTCAAGAGTTATGTCGATGCAGTCAAGTAAGCATGAAATGGTCTCAGCTGACAAAAACGGGATCGCTTTGGAAACATCTTTACCC
290 300 310 320 330 340 350 360 370
TGTTCATTTGGCCAGAGGTGACTGCTATAGTGGTCCGCAACTGAACTTGATACTGAACCTGATGATGAATGGGTGAAAAATAGGAAGATGAA
380 390 400 410 420 430 440 450 460 470
AGTCGTGCTTTTCATGAGTGGGATGAAGATCTGACATTGATGAATCTGAAGAGTCTGCCGAGGAATCAATTGCTATCAGCATTCACAAATGG
480 490 500 510 520 530 540 550 560
AAAAAGCTTTACTCCATGGCTTAATTCATAACGTTCTACCATATGTTGGTACTTCTGTAAAAACCTTAGTATTAGCATACAGCTCTGCACTTTG
570 580 590 600 610 620 630 640 650
CAGCAAAATGGTTAGGCAGATTTTAGAGCTTTTCTTAACCTGGAGCATCTGGATCTTACCCAGACTGACATTTGAGATTTGACAGT
660 670 680 690 700 710 720 730 740 750
TGGTCTTGGCTTGGTTGCTGCCAGAGTCTTCCGCATCTTGATCTGCTGGTTGTGAGAAAAATCAGAGATGTGGCCCTAGAGAAGATTTCCAGAG
760 770 780 790 800 810 820 830 840
CTCTTGAATTTGACATCTCATCAAGTGGCTTTTGAACATCTTACAAGCAAAATTAATTCAACTGCGTGGAAAAATAAGACATTACCAT
850 860 870 880 890 900 910 920 930 940
GCAGTCCACCAAGCAGTATGCTCTTTGACAGATTTAATAACAAGGGCATTTGAGAGAGAAATAGATAATGAACACCCCTGGACTAAGCGTTT
950 960 970 980 990 1000 1010 1020 1030
TCTTCTGAGAAATTCACATCTCTCTTATGCTGGATGTTAGATGCTCAAGATTTGGCTGATATTGAAGATACTGTGGAATGGAGACATAGAAATG
1040 1050 1060 1070 1080 1090 1100 1110 1120
TTGAAAGTCTTTGTCTAATGGAAACAGCATCCAACCTTTAGTTGTTCCACCTCTGGTTGTTTGTAGTAAGGACATGTTGGACTAAGGACTAGTGT
1130 1140 1150 1160 1170 1180 1190 1200 1210 1220
CTGTTGGCAGCAGATTTGCTCTCTCCAGCCTTTCCGTATTGTGCTCACTCATTTTGTGTTGACAGAACAGCTTTAAGAACTATGTCATCACTC
1230 1240 1250 1260 1270 1280 1290 1300 1310
CCAGAACTTTCTGCAATGTGTAGAAAAAGCAAGGACTAGATTGCTAGGGGAAAGACTTAATTTACTTTGGGAGTGAAAAATCTGATCAAG
1320 1330 1340 1350 1360 1370 1380 1390 1400 1410
AGACTGGACGTGACTTCTCTCTCAGTTTATCTGATGTTATCAGATCAGACCATGGTCTCAGGGTTTGGACTCTGGGAGGAGGCTGCC
1420 1430 1440 1450 1460 1470 1480 1490 1500
TTATTTGAGGACCTTAATCTCTCTGTTGCTTACTATACTGCTGACGGCTGACAGGATTTGGTTTCAGCATGCTCTTCTCTGAATGATGAA
1510 1520 1530 1540 1550 1560 1570 1580 1590
TACTTTTACTACTGTGACAACTTAACGGTCTCATGCTGATACCGCCAGTGGATGCCAGAAATTTGCAAGTGGTTTTCAGGCTGCTGCCGT
1600 1610 1620 1630 1640 1650 1660 1670 1680 1690
CTGGCCAAATGACCCCTGACTTCTGATCTTTGTCTACTTTCATTAGCTGAGCAGGCTTCTTTCATGCACITTTACTCATAGCACATTTCTTGTGT
1700 1710 1720 1730 1740 1750 1760 1770
TAACCATCCCTTTTGGAGCTGACTTCTTTTGGGCCATTNITTACAACCTCAGAAATCTTAATTACCAAGTGRATTGTAATGTTG

FIG. 24B

10 20 30 40 50 60
RVTSGCGLARGSSAMVFSNNDEGLINKKLPKELLRRIFSFLDIVTLCRCAQISKAWNILA

70 80 90 100 110 120
LDGSNWQRIDLFNFQIDVEGRVVENISKRCVGFRLKLSLRGCIGVGDSSLKTFAQNCRNI

130 140 150 160 170 180
EHLNLNGCTPKITDSTCYSLSRFCSKHLXLTSCV SITNSSLKGISEGCRNLEYLNLSWC

190 200 210 220 230 240
DQITKDGIEALVRGCRGLKALLRGCTQLEDEALKHIQNYCHELVSLNLQSCSRITDEGV

250 260 270 280 290 300
VQICRGCHRLQALCLSGCSNLTDASLTALGLNCPRLQILEAARCSHLTDAGFTLLARNCH

310 320 330 340 350 360
ELEKMDLEXCILITDSTLIQLSIHCPKLQALSLSHCELIXDDGILHLSNSTCGHERLRVL

370 380 390 400 410 420
ELDNCLLITDVALXHLNCRGLERLELYDCQQVTRAGIKRMRAQLPHVKVHAYFAPVTPP

430 440 450 460 470 480
TAVAGSGQRLCRCCVIL*QQLPGPKG*GILSSRRPESS*PTPSPNLLILHWERHLQFP

490 500 510 520 530 540
NRHLRSFKNGEDKKGFISNI*HHIVT*NMALT*LVLLLPSSLMSSLTSTHLL*YL*RLI

550
ILKTDQTGPASKYINCVQ*

FIG. 25A

10 20 30 40 50 60 70 80 90
TTTACTGTACACAGTGTGATGTATTTTGATGCTGGGCTGTCTGGTCTGTCTTGAGGATTATTAACTTTAGAGGTATCAGAGAAGCAATGGG
100 110 120 130 140 150 160 170 180
TACTGTGTAGGGCTGCTCATTAGGGGAAGAGGGCAAAAGGAGCACTAGCTAGGTGAGGCCATGTTTCAGGTCACAATGTGATGTCTAGATGTTGCT
190 200 210 220 230 240 250 260 270 280
TATAAATCCTTTCTGTCTTCGCCATTCTTAAATCTTGATAGGTGCTGTTGGGAAACTGTAAATGCCTTTCCCAATGGAGAATCAACAGATTG
290 300 310 320 330 340 350 360 370
GGTGATGGTGGAGTCGGTCAGGAAGACTCAGGTCTTCTAGAGGAAAGGATGCTCATCACCCTTNGGCCAGGCAGCTGCTGTCAGAGAATGA
380 390 400 410 420 430 440 450 460 470
CACAGCACCTGCACAGTCGCTGCCACTTCTGCACTGCTGTCCGTGGGGTACGGGAGCAAAAGTAGGGCTGGACTTTGACATGAGGGAGCTG
480 490 500 510 520 530 540 550 560
AGCCCGATCCGCTTGATGCTGACGGGTAACTGCTGGCAGTGTACAGCTCGAGGGCTCCAGGCTCGGCAGTTCTCTAGGTGTTCAGG
570 580 590 600 610 620 630 640 650
GCCACATCAGTGATGAGGAGGCAGTTGTCCAATCCAGTACCCGACGCTCTCATGCCCCACAGGTACTGTGCTCAGGTGCAGGATCCCATCAT
660 670 680 690 700 710 720 730 740 750
CTGKGATGAGTTACAGTGGGACAGGCTCAGGGCTTGCAGTTTAGGACAGTGAANTGGAGAGCTGGATGAGTGTCTCTCGTTATCAGGATGCA
760 770 780 790 800 810 820 830 840
WTCTTCAAGATCCATCTTCTCCAATTCTGTCGAATTCAGGCTAAAAGTGTAAAACCTGCGTCAAGTCAAAATGGGAGCATCGGGACGCTCCAAA
850 860 870 880 890 900 910 920 930 940
ATTTGCAGTCGGGACAGCTTCAAAACCCAGGGCTGTAAGAGAGGCATCTGTGAGGTGTGCTGCAACCCGAAAGGCAGAGAGCTGTAGCCGGTGAC
950 960 970 980 990 1000 1010 1020 1030
AGCCCTGCATATCTGACCCACACCTTCATCCGTGATACGTGAGCAGGACTGCAAGTTGAGGCTCACAAGCTCATGGCAGTAATTCTGAATGTG
1040 1050 1060 1070 1080 1090 1100 1110 1120
TTTCAGAGCTTCATCTTCTAACTGTGTGACGCCCTCAGGAGCAGGGCTTTCAGGCTCGACAACTCGCACCACTGCTCGATGCCATCCTTC
1130 1140 1150 1160 1170 1180 1190 1200 1210 1220
GTGATCTGATCACACCAAGAGAGGTTTCAGTACTCCAGTTTCGGCAGCCCTCACTGATCCCTTCAAGGAGCTGTTTGTATAGACACACAGG
1230 1240 1250 1260 1270 1280 1290 1300 1310
AGGTCAGAWCCAGATGTTTCAGCTTGGAAACAGAACTGTCTAAGGCTATAACACGTGCTGTCAAGTATTTTGTGATCCATTGAGGTTCAAATG
1320 1330 1340 1350 1360 1370 1380 1390 1400 1410
TTCAATGTTTCGGCAGTTCTGTGCAAAAGGCTTCAAGGAGGAATCCCAACACCAATCCAGCCTCGCAAGCTGAGCTTCTCAGGAATCCAAAG
1420 1430 1440 1450 1460 1470 1480 1490 1500
CATCGCTTCGAGATATTTCCACCACTCGACCTCTACATCTATTTGAAAAGTTAAAAAGATCTATTCTTTGCCAGTTGCTTCCATCCAGGGCTA
1510 1520 1530 1540 1550 1560 1570 1580 1590
AGAATGTTCCAAAGCTTGGAAATCTGTGCACATCGGCACAAAGTTACTATATCCAAGAAGGAAATATTCTTAACAGAAGTTCTTTGGGTAACCT
1600 1610 1620 1630 1640 1650 1660 1670 1680
TTTGTTAATAAGGCTTCATCATTTGTTTGAAGAAACCATGGCCGAAGAGCCCGAGCCGAGCCACAGCCCGAAGTCAACACGGC

FIG. 25B

10 20 30 40 50 60
MSPVFPMLTVLTMFYIICLRRRARTATRGEMMNTHRAIESNSQTSPLNAEVVQYAKEVVD

70 80 90 100 110 120
FSSHYGSENSMSYTMWNLAVPNVFPSSGDFQTAVFRTYGTWWDQCPSASLPFKRTPFN

130 140 150 160 170 180
FQSQDYVELTFEQQVYPTAVHVLETYHPGAVIRILACSANPYSNPPAEVRWEILWSERP

190 200 210 220 230 240
TKVNASQARQFKPCIKQINFPTNLIRLEVNSSLEYYTELDAVVLHGVKDKPVLSLKTSL

250 260 270 280 290 300
IDMNDIEDDAYAEKDGCGMDSLNNKFSSAVLGEGPNNGYFDKLPYELIQLILNHLTLPDL

310 320 330 340 350 360
CRLAQTCKLSSQHCCDPLQYIHLNLQPYWAKLDDTSLEFLQSRCTLVQWLNL SWTGNRGF

370 380 390 400 410 420
ISVAGFSRFLKVCSELVRLELSCSHFLNETCLEVISEMCPNLQALNLSSCDKLPPQAFN

430 440 450 460 470 480
HIAKLCSLKRVLVYRTKVEQTALLSILNFCSELQHLSLGSCVMIEDYDVIA SMIGAKCKK

490 500 510 520 530 540
LRTL DLWRCKNITENGIAELASGCPLLEELDLGWQPTLQSSTGCFTRLAHQLPNLQKFL

550 560 570 580 590 600
TANRSVCDTDIDELACNCTRLQQLDILGTRMVSPASLRKLLSCKDLSLLDVSFCSQIDN

610 620
RAVLELNASF PKVFIKKSFTQ

FIG. 26A

10 20 30 40 50 60 70 80 90
ATGTCACCGGTCCTTCCCATGTTAACAGTTCTGACCATGTTTTATTATATATGCTTCGGCGCGAGCCAGGACAGCTACAAGAGGAGAAATGA
100 110 120 130 140 150 160 170 180
TGAACACCCATAGAGCTATAGAATCAAACAGCCAGACTTCCCTCTCAATGCGAGAGGTAGTCCAGTATGCCAAAGAACTAGTGGATTTCAGTTTC
190 200 210 220 230 240 250 260 270 280
CCATTATGGAAGTGAGAAATAGTATGTCCTATAGTATGTTGGAATTTGGCTGGTGTACCAAATGTATTCCCAAGTTCTGTGACTTTACTCAGACA
290 300 310 320 330 340 350 360 370
GCTGTGTTTGAACCTTATGGGACATGGTGGGATCAGTGTCTTAGTGTCTTCCATTCAAGAGGACGCCACCTAATTTTCAGAGCCAGGACT
380 390 400 410 420 430 440 450 460 470
ATGTGGAACCTTACTTTTGAACAACAGGTGTATCTTACAGCTGTACATGTTCTAGAAACCTATCATCCCGAGCAGTCATTAGAAATTCCTGCTTG
480 490 500 510 520 530 540 550 560
TTCTGCAAACTCTTATTCGCCAAATCCACAGCTGAAGTAAAGATGGGAGATTCTTTGGTCAAGAGAGCTACGAAGGTGAATGCTTCCCAAGCT
570 580 590 600 610 620 630 640 650
CCCCAGTTTAAACCTTGTATTAAAGCAGATAAATTTCCCCACAAATCTTATACGACTGGAAAGTAAATAGTTCTCTTCTGGAATATTACACTGAAT
660 670 680 690 700 710 720 730 740 750
TAGATGCGAGTTGTGCTACATGCTGTGAAGGACAAGCCAGTCTTCTCTCAAGACTTCACTTATTGACATGAATCATATAGAGATGATGCTTA
760 770 780 790 800 810 820 830 840
TGCAGAAAGGATGCTTCTGGAATGGACAGTCTTAACAAAAGTTTAGCAGTCTGTCTCGGGGAAGGGCCAAATATGGGTATTTTGATAAA
850 860 870 880 890 900 910 920 930 940
CTACCTTATGAOCTTATTCAGCTGATTCTGAATCATCTTACACTACGACACCTGTGTAGATTAGCACAGACTTGCAAACTACTGAGCCAGCATT
950 960 970 980 990 1000 1010 1020 1030
GCTGTGATCTCTGCAATACATCCACCTCAATCTGCAACCATACTGGGCAAACTAGATGACACTTCTCTGGAATTTCTACAGTCTCGCTGCAC
1040 1050 1060 1070 1080 1090 1100 1110 1120
TCTGTCCAGTGGCTTAATTTATCTTGGAGTGGCAATAGAGGCTTCATCTCTGTTCAGGAGTTTAGCAGGTTTCTGAAGGTTTGTGGATCCGAA
1130 1140 1150 1160 1170 1180 1190 1200 1210 1220
TTAGTAGCGCTTGAATTTGCTTTCAGCCACTTTCTTAAAGAACTTCTTGAAGTTATTTCTGAGATGTGTCCAAATCTACAGGCTTAAATC
1230 1240 1250 1260 1270 1280 1290 1300 1310
TCTCTCTCTGTGATAAGCTTACCACTCAAGCTTTCAACCACATTCCTCAAGTTATGCGAGCTTAAACGACTTGTCTCTATCGAACAAAGTAGA
1320 1330 1340 1350 1360 1370 1380 1390 1400 1410
GCAACAGCACTGCTCAGCATTTTGAACCTTCTGTTTCAAGGCTTCAGCACTCAGTTTAGGCAGTTGTGTCATGATTGAAGACTATGATGTGATA
1420 1430 1440 1450 1460 1470 1480 1490 1500
GCTAGCATGATAGGAGCCAAAGTGTAAAAAAGTCCGGACCTGGATCTGTGAGAGATGTAGAATATTACTGAGAATGGAATAGCAGAACTGGCTT
1510 1520 1530 1540 1550 1560 1570 1580 1590
CTGGGTGTCCACTACTGGAGGAGCTTGACCTTGGCTGGTCCCAACTCTGCAGAGCAGCACCGGGGTCTTACCAGACTGGCACCCAGCTCCC
1600 1610 1620 1630 1640 1650 1660 1670 1680 1690
AAACTTGCAAAAAGCTTTCTTACAGCTAATAGATCTGTGTGTGACACAGACATTGATGAATGGCATGTAATTTGTACCAGCTTACAGCAGCTG
1700 1710 1720 1730 1740 1750 1760 1770 1780
GACATATTAGGAACAAGAATGGTAAGTCCGGCATCTTAAAGAAAGTCTCTGGAATCTTGTAAAGATCTTCTTACTTGTATGTCTCTCTGTT
1790 1800 1810 1820 1830 1840 1850 1860
CGCAGATTGATAACAGAGCTGTGCTAGAACTGAATGCAAGCTTTCCAAAAGTGTTCATAAAAAAGAGCTTTACTCAGTGA

FIG. 26B

10 20 30 40 50 60
MQLVPDIEFKITYTRSPDGDGVGNSYIEDNDDSKMADLLSYFQQQLTFQESVLKLCQPE

70 80 90 100 110 120
LESSQIHISVLPMEVLMYIFRWVSSDLDLRSLEQLSLVCRGFYICARDPEIWRACLKV

130 140 150 160 170 180
WGRSCIKLVPYTSWREMFLERPRVRFDGVYISKTTYIRQGEQSLDGFYRAWHQVEYYRYI

190 200 210 220 230 240
RFFPDGHVMMLTTPPEEPQSIVPRLRTRNTRTDAILLGHYRLSQDTDNQTKVFAVITKKKE

250 260 270 280 290 300
EKPLDYKYRYFRVPVQEADQSFHVGLQLCSSGHQRFNKLIWIHHSCHITYKSTGETAVS

310 320
AFEIDKMYTPLFFARVRSYTAFSERPL

FIG. 27A

10 20 30 40 50 60
ATGCAACTTGTACCTGATATAGAGTTCAAGATTACTTATACCCGGTCTCCAGATGGTGATGGCGTTGGA
70 80 90 100 110 120 130
AACAGCTACATTGAAGATAATGATGATGACAGCAAAATGGCAGATCTCTTGTCCTACTTCCAGCAGCAA
140 150 160 170 180 190 200
CTCACATTTTCAGGAGTCTGTGCTTAACTGTGTGAGCCTGAGCTTGAGAGCAGTCAGATTCACATATCA
210 220 230 240 250 260 270
GTGCTGCCAATGGAGGTCCTGATGTACATCTTCCGATGGGTGGTGTCTAGTGAAGTGGACCTCAGATCA
280 290 300 310 320 330 340
TTGGAGCAGTTGTGCTGGTGTGCAGAGGATTCTACATCTGTGCCAGAGACCCTGAAATATGGCGTCTG
350 360 370 380 390 400 410
GCCCTGCTTGAAAGTTTGGGGCAGAAGCTGTATTAACTTGTTCCGTACACGTCCTGGAGAGAGATGTTT
420 430 440 450 460 470 480
TTAGAACGGCCTCGTGTTCGGTTTGATGGCGTGTATATCAGTAAAACCATATATTCGTCAAGGGGAA
490 500 510 520 530 540 550
CAGTCTCTTGATGGTTTCTATAGAGCCTGGCACCAAGTGAATATTACAGGTACATAAGATTCTTTCCT
560 570 580 590 600 610 620
GATGGCCATGTGATGATGTTGACAACCCCTGAAGAGCCTCAGTCCATTGTTCCACGTTTAAGAACTAGG
630 640 650 660 670 680 690
AATACCAGGACTGATGCAATTCTACTGGGTCACTATCGCTTGTGACAAGACACAGACAATCAGACCAA
700 710 720 730 740 750
GTATTTGCTGTAATAACTAAGAAAAAAGAAGAAAAACCACTTGACTATAAATACAGATATTTTCGTCGT
760 770 780 790 800 810 820
GTCCCTGTACAAGAAGCAGATCAGAGTTTTTCATGTGGGGCTACAGCTATGTTCCAGTGGTCACCAGAGG
830 840 850 860 870 880 890
TTCAACAACTCATCTGGATACATCATTCTTGTACATTACTTACAAATCAACTGGTGAGACTGCAGTC
900 910 920 930 940 950 960
AGTGCTTTTGAGATTGACAAGATGTACACCCCTTGTTCCTCGCCAGAGTAAGGAGCTACACAGCTTTC
970 980
TCAGAAAGGCCTCTGTAG

FIG. 27B

10	20	30	40	50	60
AALDPDLENDDFFVRKTGAFHANPYVLRAFEDFRKFSEQDDSVERRDIILQCREGELVLPD					
70	80	90	100	110	120
LEKDDMIVRRIPAQKKEVPLSGAPDRYHPVFPPEPWTLPPEIQAKFLCVLERTCPSKEKS					
130	140	150	160	170	180
NSCRILVPSYRQKKDDMLTRKIQSWKLGTTVPPISTPGPCSEADLKRWEAIREASRLRH					
190	200	210	220	230	240
KKRLMVERLFQKIYGENGSKSMSDVSAEDVQNLRLRYEEMQKIKSQLKEQDQKWQDDLA					
250					
KWKDRRKSYTSDLQK					

FIG. 28A

10 20 30 40 50 60
GCAGCCCTGGATCCTGACTTAGAGAATGATGATTCTTTGTCAGAAAGACTGGGGCTTTCCATGCAAAT
70 80 90 100 110 120 130
CCATATGTTCTCCGAGCTTTTGAAGACTTTAGAAAGTTCTCTGAGCAAGATGATTCTGTAGAGCGAGAT
140 150 160 170 180 190 200
ATAATTTTACAGTGTAGAGAAGGTGAACCTTGTACTTCCGGATTGGAAGATGATATGATTGTTCCG
210 220 230 240 250 260 270
CGAATCCCAGCACAGAAGAAAGAGTCCCGCTGTCTGGGGCCCCAGATAGATACCACCCAGTCCCTTTT
280 290 300 310 320 330 340
CCCGAACCCTGGACTCTTCTCCAGAAATTCAGCAAAATTTCTCTGTGTACTTGAAAGGACATGCCCA
350 360 370 380 390 400 410
TCCAAAGAAAAAGTAATAGCTGTAGAATATTAGTTCTTCATATCGGCAGAGAAAGATGACATGCTG
420 430 440 450 460 470 480
ACACGTAAGATTCACTCCTGGAACTGGGAACTACCGTGCCTCCCATCAGTTTCACNCTGGCCCCCTGC
490 500 510 520 530 540 550
AGTGAGGCTGACTTGAAGAGATGGGAGGCCATCCGGGAGGCCAGCAGACTCAGGCACAAGAAAAGGCTG
560 570 580 590 600 610 620
ATGGTGGAGAGACTCTTTCAAAGATTTATGGTGAGAATGGGAGTAAGTCCATGAGTGATGTCAGCGCA
630 640 650 660 670 680 690
GAAGATGTTCAAACCTGCGTCAGCTGCGTTACGAGGAGATGCAGAAAATAAATCACAATTAAAAGAA
700 710 720 730 740 750
CAAGATCAGAAATGGCAGGATGACCTTGCAAAATGGAAAGATCGTCGAAAAAGTTACACTTCAGATCTG
760
CAGAAG

FIG. 28B




FIG. 29




FIG. 30




FIG. 31




FIG. 32




FIG. 33




FIG. 34 A-E




FIG. 35 A-C




FIG. 36 A-B




FIG. 37 A-C




FIG. 38




FIG. 39 A-B




FIG. 40 A-C



B




FIG. 41 A-B




FIG. 42




FIG. 43 A-C




FIG. 44





FIG. 45

A.



B.




FIG. 46




FIG. 47


A

FIG. 48




FIG. 48



A**B**

FIG. 49





C**D**

FIG. 49

A**B****FIG. 5D**



C**D**

FIG. 5D




FIG. 51 A-C

SEQUENCE LISTING

<110> Pagano, M.

<120> METHODS TO IDENTIFY COMPOUNDS USEFUL FOR THE TREATMENT OF
PROLIFERATIVE AND DIFFERENTIATIVE DISORDERS

<130> 5914-090-228

<140> To be assigned

<141> 2002-1-07

<150> 60/260,179

<151> 2001-01-5

<160> 89

<170> PatentIn Ver. 2.0

<210> 1

<211> 2151

<212> DNA

<213> Homo sapiens

<400> 1

```
tgcgttggt ggcgcctggc accaaagggg cggcccccggc ggagagcgga cccagtggcc 60
tcggcgatta tggaccggc cgaggcggtg ctgcaagaga aggcactcaa gtttatgaat 120
tcctcagaga gagaagactg taataatggc gaacccctta ggaagataat accagagaag 180
aattcactta gacagacata caacagctgt gccagactct gcttaaacca agaaacagta 240
tgtttagcaa gcactgctat gaagactgag aattgtgtgg ccaaaacaaa acttgccaat 300
ggcacttoca gtatgattgt gcccaagcaa cggaaactct cagcaagcta tgaaaaggaa 360
aaggaaactgt gtgtcaataa ctttgagcag tggtcagagt cagatcaagt ggaatttgtg 420
gancatctta tatcccaaat gtgtcattac caacatgggc acataaactc gtatcttaaa 480
cctatgttgc agagagattt cataactgct ctgccagctc ggggatttga tcatatcgct 540
gagaacattc tgcataacct ggatgccaaa tcaactatgt ctgctgaact tgtgtgcaag 600
gaatggtaac gagtgacctc tgatggcatg ctgtggaaga agcttatcga gagaatggtc 660
aggacagatt ctctgtggag aggcctggca gaacgaagag gatggggaca gtatttatc 720
aaaaacaaac ctctgacggg gaatgctcct cccaactott ttatagagc actttatcct 780
aaaattatac aagacattga gacaatagaa tctaattgga gatgtggaag acatagttaa 840
cagagaattc actgcgaag tgaaacaagc aaaggagtct actgtttaca gtatgatgat 900
cagaaaatag taagcggcct tcgagacaac acaatcaaga tctgggataa aaacacatg 960
gaatgcaagc gaattctcac aggcataca ggttcagtc tctgtctcca gtatgatgat 1020
agagtgcata taacaggatc atcggattcc acggtcagag tgtgggatgt aaatacaggt 1080
gaaatgctaa acacgttgat tcaccattgt gaagcagttc tgcacttgcg tttcaataat 1140
ggcatgatgg tgacctgctc caaagatcgt tccattgctg tatgggatat ggccctccca 1200
actgacatta cctccggag ggtgctggtc ggacaccgag ctgctgtcaa tgtttagac 1260
tttgatgaca agtacattgt ttctgcatct ggggatagaa ctataaaggt atggaacaca 1320
agtactltgt aatttgtaag gaccttaaat ggacacaaac gaggcattgc ctgtttgcag 1380
tacagggaca ggctggtagt gagtggctca tctgacaaca ctatcagatt atgggacata 1440
gaatgtggcg catgtttacg agtgttagaa ggccatgagg aattgggtgcg ttgtattcga 1500
tttgataaca agaggatagt cagtggggcc tatgatggaa aaattaaagt gtgggatctt 1560
gtggctgctt tggacccccg tgetcctgca gggacactct gtctacggac ccttgtggag 1620
cattccggaa gagtttttgc actacagttt gatgaattcc agattgtcag tagttcacat 1680
gatgacacaa tctcatctg ggaacttcta aatgatccag ctgcccgaagc tgaaccccc 1740
cgttccctt ctcgaacata cactacatc tccagataaa taaccataca ctgacctcat 1800
acttgcccag gaccatttaa agttgcggta tttaacgtat ctgccaatac caggatgagc 1860
aacaacagta acaatcaaac tactgccag tttccctgga ctagccgagg agcagggctt 1920
tgagactcct gttgggacac agttggctcg cagtcggccc aggaaggctc actcagcaca 1980
actgactgct tcagtgtgc tatcagaaga tgtctctat caattgtgaa tgattggaac 2040
ttttaaacct cccctctct cctccttca cctctgcacc tagttttttc ccattgggtc 2100
cagacaaaagg tgacttataa atatatttag tgttttgcca gaaaaaaaaa a 2151
```

<210> 2
 <211> 569
 <212> PRT
 <213> Homo sapiens

 <400> 2
 Met Asp Pro Ala Glu Ala Val Leu Gln Glu Lys Ala Leu Lys Phe Met
 1 5 10 15
 Asn Ser Ser Glu Arg Glu Asp Cys Asn Asn Gly Glu Pro Pro Arg Lys
 20 25 30
 Ile Ile Pro Glu Lys Asn Ser Leu Arg Gln Thr Tyr Asn Ser Cys Ala
 35 40 45
 Arg Leu Cys Leu Asn Gln Glu Thr Val Cys Leu Ala Ser Thr Ala Met
 50 55 60
 Lys Thr Glu Asn Cys Val Ala Lys Thr Lys Leu Ala Asn Gly Thr Ser
 65 70 75 80
 Ser Met Ile Val Pro Lys Gln Arg Lys Leu Ser Ala Ser Tyr Glu Lys
 85 90 95
 Glu Lys Glu Leu Cys Val Lys Tyr Phe Glu Gln Trp Ser Glu Ser Asp
 100 105 110
 Gln Val Glu Phe Val Glu His Leu Ile Ser Gln Met Cys His Tyr Gln
 115 120 125
 His Gly His Ile Asn Ser Tyr Leu Lys Pro Met Leu Gln Arg Asp Phe
 130 135 140
 Ile Thr Ala Leu Pro Ala Arg Gly Leu Asp His Ile Ala Glu Asn Ile
 145 150 155 160
 Leu Ser Tyr Leu Asp Ala Lys Ser Leu Cys Ala Ala Glu Leu Val Cys
 165 170 175
 Lys Glu Trp Tyr Arg Val Thr Ser Asp Gly Met Leu Trp Lys Lys Leu
 180 185 190
 Ile Glu Arg Met Val Arg Thr Asp Ser Leu Trp Arg Gly Leu Ala Glu
 195 200 205
 Arg Arg Gly Trp Gly Gln Tyr Leu Phe Lys Asn Lys Pro Pro Asp Gly
 210 215 220
 Asn Ala Pro Pro Asn Ser Phe Tyr Arg Ala Leu Tyr Pro Lys Ile Ile
 225 230 235 240
 Gln Asp Ile Glu Thr Ile Glu Ser Asn Trp Arg Cys Gly Arg His Ser
 245 250 255
 Leu Gln Arg Ile His Cys Arg Ser Glu Thr Ser Lys Gly Val Tyr Cys
 260 265 270
 Leu Gln Tyr Asp Asp Gln Lys Ile Val Ser Gly Leu Arg Asp Asn Thr
 275 280 285
 Ile Lys Ile Trp Asp Lys Asn Thr Leu Glu Cys Lys Arg Ile Leu Thr
 290 295 300

3/57

Gly His Thr Gly Ser Val Leu Cys Leu Gln Tyr Asp Glu Arg Val Ile
 305 310 315 320
 Ile Thr Gly Ser Ser Asp Ser Thr Val Arg Val Trp Asp Val Asn Thr
 325 330 335
 Gly Glu Met Leu Asn Thr Leu Ile His His Cys Glu Ala Val Leu His
 340 345 350
 Leu Arg Phe Asn Asn Gly Met Met Val Thr Cys Ser Lys Asp Arg Ser
 355 360 365
 Ile Ala Val Trp Asp Met Ala Ser Pro Thr Asp Ile Thr Leu Arg Arg
 370 375 380
 Val Leu Val Gly His Arg Ala Ala Val Asn Val Val Asp Phe Asp Asp
 385 390 395 400
 Lys Tyr Ile Val Ser Ala Ser Gly Asp Arg Thr Ile Lys Val Trp Asn
 405 410 415
 Thr Ser Thr Cys Glu Phe Val Arg Thr Leu Asn Gly His Lys Arg Gly
 420 425 430
 Ile Ala Cys Leu Gln Tyr Arg Asp Arg Leu Val Val Ser Gly Ser Ser
 435 440 445
 Asp Asn Thr Ile Arg Leu Trp Asp Ile Glu Cys Gly Ala Cys Leu Arg
 450 455 460
 Val Leu Glu Gly His Glu Glu Leu Val Arg Cys Ile Arg Phe Asp Asn
 465 470 475 480
 Lys Arg Ile Val Ser Gly Ala Tyr Asp Gly Lys Ile Lys Val Trp Asp
 485 490 495
 Leu Val Ala Ala Leu Asp Pro Arg Ala Pro Ala Gly Thr Leu Cys Leu
 500 505 510
 Arg Thr Leu Val Glu His Ser Gly Arg Val Phe Arg Leu Gln Phe Asp
 515 520 525
 Glu Phe Gln Ile Val Ser Ser Ser His Asp Asp Thr Ile Leu Ile Trp
 530 535 540
 Asp Phe Leu Asn Asp Pro Ala Ala Gln Ala Glu Pro Pro Arg Ser Pro
 545 550 555 560
 Ser Arg Thr Tyr Thr Ile Ser Arg
 565

<210> 3
 <211> 1476
 <212> DNA
 <213> Homo sapiens

<400> 3
 atggagagaa aggactttga gacatggcctt gataacattt ctgttacatt tctttctctg 60
 acggacttgc agaaaaatga aactotggat cacctgatta gtctgagtgg ggcagtccag 120
 ctcaggcatc tctccaataa cctagagact ctctcaagc gggacttctt caaactcctt 180
 cccctggagc tcagttttta ttgttaaaa tggctcgatc ctcagacttt actcacatgc 240
 tgccctcgtc ctaaacagtg gaataagggtg ataagtgcct gtacagaggt gtggcagact 300

```

gcacgtaaaa atttgggctg gcagatagat gattctgttc aggacgcttt gcactggaag 360
aagctttatt tgaaggctat tttgagaatg aagcaactgg aggaccatga agcctttgaa 420
acctcgatcat taattggaca cagtggcaga gtgtatgcac ttactacaa agatggactt 480
ctctgtacag ggtcagatga cttgtctgca aagctgtggg atgtgagcac agggcagtgc 540
gtttatggca tccagacca cacttgtgca gcggtgaagt ttgatgaaca gaagcttgtg 600
acaggctcct ttgacaacac tgtggcttgc tgggaatgga gttccggagc caggaccag 660
cactttcggg ggcacacggg ggcggtatct agcgtggact acaatgatga actggatgc 720
ttgggtgagcg gctctgcaga cttcactgtg aaagtatggg ctttatctgc tgggacatgc 780
ctgaacacac tcaccgggca cagggaatgg gtcaccaagg tagttttgca gaagtgc 840
atcaagtctc tcttgacacg tcttgagac tacatctct taagtgcaga caaatatgag 900
attaagattt ggccaattgg gagagaaatc aactgtaagt gcttaaagac attgtctgtc 960
tctgaggata gaagtatctg cctgcagcca agacttcatt ttgatggcaa atacattgtc 1020
tgtagttcag cacttggctc ctaccagtgg gactttgcca gttatgatat tctcagggtc 1080
atcaagactc ctgagatagc aaacttggcc ttgcttggct ttggagatat ctttgcctg 1140
ctggttgaca accgctacct gtacatcatg gacttgcgga cagagagcct gattgtctgc 1200
tggcctctgc cagagtacag ggaatcaaag agaggctcaa gcttcttggc aggcgaacat 1260
cctggtgaa tggactggat gggcacaatg acacgggctt ggtctttgcc accagcatgc 1320
ctgaccacag tattcacctg gtgttgtgga aggagcacgg ctgacaccat gagccaccac 1380
cgctgactga ctttgggtgc cggggtgctg ggttttgggt gcacctctgc ggcacgcgac 1440
tgcacgaacc aaagtcttca cctaattgta tcatca 1476

```

<210> 4
 <211> 422
 <212> PRT
 <213> Homo sapiens

<400> 4
 Met Glu Arg Lys Asp Phe Glu Thr Trp Leu Asp Asn Ile Ser Val Thr
 1 5 10 15
 Phe Leu Ser Leu Thr Asp Leu Gln Lys Asn Glu Thr Leu Asp His Leu
 20 25 30
 Ile Ser Leu Ser Gly Ala Val Gln Leu Arg His Leu Ser Asn Asn Leu
 35 40 45
 Glu Thr Leu Leu Lys Arg Asp Phe Leu Lys Leu Leu Pro Leu Glu Leu
 50 55 60
 Ser Phe Tyr Leu Leu Lys Trp Leu Asp Pro Gln Thr Leu Leu Thr Cys
 65 70 75 80
 Cys Leu Val Ser Lys Gln Trp Asn Lys Val Ile Ser Ala Cys Thr Glu
 85 90 95
 Val Trp Gln Thr Ala Cys Lys Asn Leu Gly Trp Gln Ile Asp Asp Ser
 100 105 110
 Val Gln Asp Ala Leu His Trp Lys Lys Val Tyr Leu Lys Ala Ile Leu
 115 120 125
 Arg Met Lys Gln Leu Glu Asp His Glu Ala Phe Glu Thr Ser Ser Leu
 130 135 140
 Ile Gly His Ser Ala Arg Val Tyr Ala Leu Tyr Tyr Lys Asp Gly Leu
 145 150 155 160
 Leu Cys Thr Gly Ser Asp Asp Leu Ser Ala Lys Leu Trp Asp Val Ser
 165 170 175
 Thr Gly Gln Cys Val Tyr Gly Ile Gln Thr His Thr Cys Ala Ala Val
 180 185 190

5/57

Lys Phe Asp Glu Gln Lys Leu Val Thr Gly Ser Phe Asp Asn Thr Val
 195 200 205
 Ala Cys Trp Glu Trp Ser Ser Gly Ala Arg Thr Gln His Phe Arg Gly
 210 215 220
 His Thr Gly Ala Val Phe Ser Val Asp Tyr Asn Asp Glu Leu Asp Ile
 225 230 235 240
 Leu Val Ser Gly Ser Ala Asp Phe Thr Val Lys Val Trp Ala Leu Ser
 245 250 255
 Ala Gly Thr Cys Leu Asn Thr Leu Thr Gly His Thr Glu Trp Val Thr
 260 265 270
 Lys Val Val Leu Gln Lys Cys Lys Val Lys Ser Leu Leu His Ser Pro
 275 280 285
 Gly Asp Tyr Ile Leu Leu Ser Ala Asp Lys Tyr Glu Ile Lys Ile Trp
 290 295 300
 Pro Ile Gly Arg Glu Ile Asn Cys Lys Cys Leu Lys Thr Leu Ser Val
 305 310 315 320
 Ser Glu Asp Arg Ser Ile Cys Leu Gln Pro Arg Leu His Phe Asp Gly
 325 330 335
 Lys Tyr Ile Val Cys Ser Ser Ala Leu Gly Leu Tyr Gln Trp Asp Phe
 340 345 350
 Ala Ser Tyr Asp Ile Leu Arg Val Ile Lys Thr Pro Glu Ile Ala Asn
 355 360 365
 Leu Ala Leu Leu Gly Phe Gly Asp Ile Phe Ala Leu Leu Phe Asp Asn
 370 375 380
 Arg Tyr Leu Tyr Ile Met Asp Leu Arg Thr Glu Ser Leu Ile Ser Arg
 385 390 395 400
 Trp Pro Leu Pro Glu Tyr Arg Glu Ser Lys Arg Gly Ser Ser Phe Leu
 405 410 415
 Ala Gly Glu His Pro Gly
 420

<210> 5

<211> 1407

<212> DNA

<213> Homo sapiens

<400> 5

```

cgggggtggtg tgtgggggaa gccgcccccg gcagcaggat gaaacgagga ggaagagata 60
gtgaccgtaa ttcacagaa gaaggaactg cagagaaatc caagaaactg aggactacaa 120
atgagcattc tcagacttgt gattggggta atctccttca ggacattatt ctccaagtat 180
ttaaataatt gccctttctt gaccgggctc atgcttcaca agtttgccgc aactggaacc 240
aggtatttca catgcttgac ttgtggagat gttttgaatt tgaactgaat cagccagcta 300
catcttattt gaaagctacc catccagagc tgatcaaaca gattattaaa agacattcaa 360
accatctaca atatgtcagc ttcaagggtg acagcagcaa ggaatcagct gaagcagctt 420
gtgatatact atcgcaactt gtgaattgct ctttaaaaac acttggaact atttcaactg 480
ctcgaccaag ctttatggat ttaccaaagt ctactttat ctctgcactg acagtttgtt 540
tcgtaaaact caaatcctg ttttcgttga agatagatga tactocagta gatgatccat 600
ctctcaaagt actagtggcc aacaatagt atactctaa gctgttgaaa atgagcagct 660

```

```

gtctcatgt ctctccagca ggtatccttt gtgtggctga tcagtgtcac ggcttaagag 720
aactagccct gaactaccac ttattgagtg atgagttgtt acttgcatg tcttctgaaa 780
aacatgttgc attagaacat ttgcgcattg atgtagtcag tgagaatcct ggacagacac 840
acttccatac tattcagaag agtagctggg atgctttcat cagacattca cccaaagtga 900
acttagtgat gtattttttt ttatatgaag aagaatttga ccccttcttt cgctatgaaa 960
taactgccac ccactgtgac ttggggagat cagtaagcaa agatgtgctt ggccgtgtgg 1020
gaatgacatg ccctagactg gttgaactag tagtgtgtgc aaatggatta cggccacttg 1080
atgaagagtt aattcgcatg gcagaacgtt gcaaaaaattt gtcagctatt ggactagggg 1140
aatgtgaagt ctcatgtagt ccctttgttg agtttgtgaa gatgtgtggt ggccgcctat 1200
ctcaattatc cattatggaa gaagtactaa ttccctgacca aaagtatagt ttggagcaga 1260
ttcactggga agtgtccaag catcttggta ggggtgtggt tcccgacatg atgcccactt 1320
ggtaaaaact gcatgatgaa tagcacctta atttcaagca aatgtattat aattaaagtt 1380
ttatttgctg taaaaaaaaa aaaaaaa 1407

```

<210> 6

<211> 428

<212> PRT

<213> Homo sapiens

<400> 6

```

Met Lys Arg Gly Gly Arg Asp Ser Asp Arg Asn Ser Ser Glu Glu Gly
  1             5             10             15

Thr Ala Glu Lys Ser Lys Lys Leu Arg Thr Thr Asn Glu His Ser Gln
          20             25             30

Thr Cys Asp Trp Gly Asn Leu Leu Gln Asp Ile Ile Leu Gln Val Phe
          35             40             45

Lys Tyr Leu Pro Leu Leu Asp Arg Ala His Ala Ser Gln Val Cys Arg
          50             55             60

Asn Trp Asn Gln Val Phe His Met Pro Asp Leu Trp Arg Cys Phe Glu
          65             70             75             80

Phe Glu Leu Asn Gln Pro Ala Thr Ser Tyr Leu Lys Ala Thr His Pro
          85             90             95

Glu Leu Ile Lys Gln Ile Ile Lys Arg His Ser Asn His Leu Gln Tyr
          100            105            110

Val Ser Phe Lys Val Asp Ser Ser Lys Glu Ser Ala Glu Ala Ala Cys
          115            120            125

Asp Ile Leu Ser Gln Leu Val Asn Cys Ser Leu Lys Thr Leu Gly Leu
          130            135            140

Ile Ser Thr Ala Arg Pro Ser Phe Met Asp Leu Pro Lys Ser His Phe
          145            150            155            160

Ile Ser Ala Leu Thr Val Val Phe Val Asn Ser Lys Ser Leu Ser Ser
          165            170            175

Leu Lys Ile Asp Asp Thr Pro Val Asp Asp Pro Ser Leu Lys Val Leu
          180            185            190

Val Ala Asn Asn Ser Asp Thr Leu Lys Leu Leu Lys Met Ser Ser Cys
          195            200            205

Pro His Val Ser Pro Ala Gly Ile Leu Cys Val Ala Asp Gln Cys His
          210            215            220

Gly Leu Arg Glu Leu Ala Leu Asn Tyr His Leu Leu Ser Asp Glu Leu

```

7/57

225 230 235 240
 Leu Leu Ala Leu Ser Glu Lys His Val Arg Leu Glu His Leu Arg
 245 250 255
 Ile Asp Val Val Ser Glu Asn Pro Gly Gln Thr His Phe His Thr Ile
 260 265 270
 Gln Lys Ser Ser Trp Asp Ala Phe Ile Arg His Ser Pro Lys Val Asn
 275 280 285
 Leu Val Met Tyr Phe Phe Leu Tyr Glu Glu Glu Phe Asp Pro Phe Phe
 290 295 300
 Arg Tyr Glu Ile Pro Ala Thr His Leu Tyr Phe Gly Arg Ser Val Ser
 305 310 315 320
 Lys Asp Val Leu Gly Arg Val Gly Met Thr Cys Pro Arg Leu Val Glu
 325 330 335
 Leu Val Val Cys Ala Asn Gly Leu Arg Pro Leu Asp Glu Glu Leu Ile
 340 345 350
 Arg Ile Ala Glu Arg Cys Lys Asn Leu Ser Ala Ile Gly Leu Gly Glu
 355 360 365
 Cys Glu Val Ser Cys Ser Ala Phe Val Glu Phe Val Lys Met Cys Gly
 370 375 380
 Gly Arg Leu Ser Gln Leu Ser Ile Met Glu Glu Val Leu Ile Pro Asp
 385 390 395 400
 Gln Lys Tyr Ser Leu Glu Gln Ile His Trp Glu Val Ser Lys His Leu
 405 410 415
 Gly Arg Val Trp Phe Pro Asp Met Met Pro Thr Trp
 420 425

<210> 7
 <211> 1444
 <212> DNA
 <213> Homo sapiens

<400> 7
 atggcgggaa ggcgagcgcg cagcggaaca aattcgccgc cgcgcgccct cagcgactgg 60
 ggccgccttg agcgcgccat cctcagcggc tggagagcct tctggcagtc agtgagcaag 120
 gatagggttg cgcgtacgac ctcccgggag gaggtggatg aggcggccag caccctgacg 180
 cggctgcgga ttgatgtaca gctatatatt ttgtcccttc ttccacctca tgatctgtgt 240
 cagttgggaa gtacaaatca ttattggaat gaaactgtaa gaaatccaat tctgtggaga 300
 tactttttgt tgagggatct tcttctttgg tcttctgttg actggaagtc tcttccatat 360
 ctacaaatct taaaaaagcc tataatctgag gtctctgatg gtgcattttt tgactacatg 420
 gcagtctatc taatgtgctg tccatacaca agaagagctt caaaatccag ccgtccctatg 480
 tatggagctg tcacttcttt tttaactctc ctgacatctc ccaatgaacc tggatttgct 540
 ctggtttggac cacgtttgga acaattgaat acctcttttg tgttgagctt gctgtcttca 600
 gaggaacttt gcccaacagc tgggtttgct cagaggcaga ttgatggtat tggatcagga 660
 gtcaattttc agttgaacaa ccaacataaa ttcaacatct taatcttata ttcaactacc 720
 agaaaggaaa gagatagagc aagggaagag catacaagtg cagttaacaa gatgttcagt 780
 cgacacaatg aaggtgatga tgcaccagga agccgggtaca gtgtgattcc acagattcaa 840
 aaactgtgtg aagttgtaga tgggttcacg tatgttgcaa atgctgaagc tcataaaaaga 900
 catgaatggc aagatgaatt ttctcatatt atggcaatga cagatccagc ctttgggtct 960
 tcgggaagac cattgttgtt ttatcttgtt atttctcaag gggatgtaaa aagaatgccc 1020
 tgtttttatt tggctcatga gctgcactcg aatcttctaa atcaccatg gctgggtccag 1080


```

gatacagagg ctgaaactct gactgggttt ttgaatggca ttgagtggat tcttgaagaa 1140
gtggaatcta agcgtgcaag atgattctct tticagatct tgggaactga aaccatttga 1200
aatttattac taaggctcgtg atgtgaatat ttgctcagtc agcccacctt gtcctgcctt 1260
tttgcagata ggctttcatt tggacagcta taactgctgt gttttttata ttatttttac 1320
tttttaccat aaatcaatta caagaaaaga gtttcagtc tagtatttag ccccaaaatg 1380
aacctttaa ctttttttg gtaattttta tattttctgt ctttttaaaa atattaaatt 1440
ttgg                                     1444

```

<210> 8
 <211> 472
 <212> PRT
 <213> Homo sapiens

<400> 8
 Met Ala Gly Ser Glu Pro Arg Ser Gly Thr Asn Ser Pro Pro Pro Pro
 1 5 10 15
 Phe Ser Asp Trp Gly Arg Leu Glu Ala Ala Ile Leu Ser Gly Trp Lys
 20 25 30
 Thr Phe Trp Gln Ser Val Ser Lys Asp Arg Val Ala Arg Thr Thr Ser
 35 40 45
 Arg Glu Glu Val Asp Glu Ala Ala Ser Thr Leu Thr Arg Leu Pro Ile
 50 55 60
 Asp Val Gln Leu Tyr Ile Leu Ser Phe Leu Ser Pro His Asp Leu Cys
 65 70 75 80
 Gln Leu Gly Ser Thr Asn His Tyr Trp Asn Glu Thr Val Arg Asn Pro
 85 90 95
 Ile Leu Trp Arg Tyr Phe Leu Leu Arg Asp Leu Pro Ser Trp Ser Ser
 100 105 110
 Val Asp Trp Lys Ser Leu Pro Tyr Leu Gln Ile Leu Lys Lys Pro Ile
 115 120 125
 Ser Glu Val Ser Asp Gly Ala Phe Phe Asp Tyr Met Ala Val Tyr Leu
 130 135 140
 Met Cys Cys Pro Tyr Thr Arg Arg Ala Ser Lys Ser Ser Arg Pro Met
 145 150 155 160
 Tyr Gly Ala Val Thr Ser Phe Leu His Ser Leu Ile Ile Pro Asn Glu
 165 170 175
 Pro Arg Phe Ala Leu Phe Gly Pro Arg Leu Glu Gln Leu Asn Thr Ser
 180 185 190
 Leu Val Leu Ser Leu Leu Ser Ser Glu Glu Leu Cys Pro Thr Ala Gly
 195 200 205
 Leu Pro Gln Arg Gln Ile Asp Gly Ile Gly Ser Gly Val Asn Phe Gln
 210 215 220
 Leu Asn Asn Gln His Lys Phe Asn Ile Leu Ile Leu Tyr Ser Thr Thr
 225 230 235 240
 Arg Lys Glu Arg Asp Arg Ala Arg Glu Glu His Thr Ser Ala Val Asn
 245 250 255
 Lys Met Phe Ser Arg His Asn Glu Gly Asp Asp Arg Pro Gly Ser Arg

9/57

260	265	270
Tyr Ser Val Ile Pro Gln Ile Gln Lys Leu Cys Glu Val Val Asp Gly		
275	280	285
Phe Ile Tyr Val Ala Asn Ala Glu Ala His Lys Arg His Glu Trp Gln		
290	295	300
Asp Glu Phe Ser His Ile Met Ala Met Thr Asp Pro Ala Phe Gly Ser		
305	310	315
Ser Gly Arg Pro Leu Leu Val Leu Ser Cys Ile Ser Gln Gly Asp Val		
325	330	335
Lys Arg Met Pro Cys Phe Tyr Leu Ala His Glu Leu His Leu Asn Leu		
340	345	350
Leu Asn His Pro Trp Leu Val Gln Asp Thr Glu Ala Glu Thr Leu Thr		
355	360	365
Gly Phe Leu Asn Gly Ile Glu Trp Ile Leu Glu Glu Val Glu Ser Lys		
370	375	380
Arg Ala Arg Phe Ser Phe Gln Ile Leu Gly Thr Glu Thr Ile Asn Leu		
385	390	395
Leu Leu Arg Ser Cys Glu Tyr Leu Leu Ser Gln Pro Thr Leu Ser Cys		
405	410	415
Leu Phe Ala Asp Arg Leu Ser Phe Gly Gln Leu Leu Leu Cys Phe Leu		
420	425	430
Tyr Tyr Phe Tyr Phe Leu Pro Ile Asn Tyr Lys Lys Arg Val Ser Val		
435	440	445
Leu Val Phe Ser Pro Lys Met Asn Leu Thr Phe Phe Trp Phe Leu Tyr		
450	455	460
Phe Leu Ser Phe Lys Tyr Ile Leu		
465	470	

<210> 9

<211> 2076

<212> DNA

<213> Homo sapiens

<400> 9

```

aggttgctca gctgcccccg gagcgggttcc tccacctgag gcagacacca cctcggttgg 60
catgagccgg cgcccttgca gctgcgccct acggccaccc cgtgctcct gcagcgccag 120
ccccagcgca gtgacagccg ccggggcgccc tcgacctcg gatagttgta aagaagaaa 180
ttctaccctt tctgtcaaaa tgaagtgtga ttttaattgt aacctgttc attccggact 240
taaactggta aaacctgatg acattggaag actagtttcc tacaccctg catatctgga 300
aggttcctgt aaagactgca ttaaagacta tgaaaggctg tcatgtattg ggtcaccgat 360
tgtgagccct aggattgtac aacttgaanc tgaaagcaag cgcttgcata acaaggaaaa 420
tcaacatgtg caacagacac ttaatagtac aaatgaaata gaagcactag agaccagtag 480
actttatgaa gacagtggct attcctcatt ttctctacaa agtggcctca gtgaacatga 540
agaaggtagc ctccctggagg agaatttcgg tgacagtcta caatcctgcc tgctacaaat 600
acaaagccca gaccaatata ccaacaaaaa cttgctgcca gttcttcatt ttgaaaaagt 660
ggtttgttca acattaaaaa agaattgcada acgaaatcct aaagtagatc gggagatgct 720
gaaggaaaatt atagccagag gaaatttttag actgcagaat ataattggca gaaaaatggg 780
cctagaatgt gtagatatcc tcagcgaaact ctttcgaagg ggactcagac atgtcttagc 840
aactatttta gcacaactca gtgacatgga cttaatcaat gtgtctaaag tgagcacaa 900

```

10/57

```

ttggaagaag atcclagaag atgataaggg ggcattccag ttgtacagta aagcaatata 960
aagagttacc gaaaaacaaca ataaattttc acctcatgct tcaaccagag aatatgttat 1020
gttcagaacc ccactggctt ctgttcagaa atcagcagcc cagacttctc tcaaaaaaga 1080
tgotcacaacc aagttatcca atcaagggtga tcagaaagggt tctacttata gtgcacacaa 1140
tgaattctct gaggttgcca agacattgaa aaagaacgaa agcctcaaag cctgtattcg 1200
ctgtaattca cctgcacaaat atgattgcta tttacaacgg gcaacctgca aacgagaagg 1260
ctgtggattt gattattgta cgaagtgtct ctgtaattat cataactacta aagactgttc 1320
agatggcaag ctctcacaag ccagttgtaa aatagggtccc ctgcttggtg caaagaaaag 1380
caaaaagaat ttacgaagat tgtgatctct tattaataca attgttactg atcatgaatg 1440
ttagttagaa aatgttaggt tttaacctaa aaaaaattgt attgtgattt tcaattttat 1500
gttgaaatcg gtgtagtatc ctgaggtttt tttcccccca gaagataaag aggatagaca 1560
acctcttaaa atatttttac aatttaatga gaaaaagttt aaaattctca atacaaatca 1620
aacaatttaa atattttaag aaaaaaggaa aagtagatag tgatactgag ggtaaaaaaa 1680
aaattgattc aattttatgg taaaggaaac ccattgcaatt ttacctagac agtcttaaat 1740
atgtctgggt ttccatctgt tagcatttca gacattttat gttcctctta ctcaattgat 1800
accaacagaa atatcaactt ctggagtcta ttaaatgtgt tgtcaccttt ctaaagcttt 1860
ttttcattgt gtgtattttc caagaaaagta tcctttgtaa aaacttgctt gttttcctta 1920
ttctgaaat ctgttttaat atttttgtat acatgtaaat atttctgtat tttttatatg 1980
tcaagaataa tgtctcttgt atgtacatat aaaaataaat tltgtcfaat aaaattgtaa 2040
gcttaaaaaa aaaaaaaaaa aactcgagac tagtgc 2076

```

<210> 10

<211> 447

<212> PRT

<213> Homo sapiens

<400> 10

```

Met Ser Arg Arg Pro Cys Ser Cys Ala Leu Arg Pro Pro Arg Cys Ser
  1             5             10             15

Cys Ser Ala Ser Pro Ser Ala Val Thr Ala Ala Gly Arg Pro Arg Pro
      20             25             30

Ser Asp Ser Cys Lys Glu Glu Ser Ser Thr Leu Ser Val Lys Met Lys
      35             40             45

Cys Asp Phe Asn Cys Asn His Val His Ser Gly Leu Lys Leu Val Lys
      50             55             60

Pro Asp Asp Ile Gly Arg Leu Val Ser Tyr Thr Pro Ala Tyr Leu Glu
      65             70             75             80

Gly Ser Cys Lys Asp Cys Ile Lys Asp Tyr Glu Arg Leu Ser Cys Ile
      85             90             95

Gly Ser Pro Ile Val Ser Pro Arg Ile Val Gln Leu Glu Thr Glu Ser
      100            105            110

Lys Arg Leu His Asn Lys Glu Asn Gln His Val Gln Gln Thr Leu Asn
      115            120            125

Ser Thr Asn Glu Ile Glu Ala Leu Glu Thr Ser Arg Leu Tyr Glu Asp
      130            135            140

Ser Gly Tyr Ser Ser Phe Ser Leu Gln Ser Gly Leu Ser Glu His Glu
      145            150            155            160

Glu Gly Ser Leu Leu Glu Glu Asn Phe Gly Asp Ser Leu Gln Ser Cys
      165            170            175

Leu Leu Gln Ile Gln Ser Pro Asp Gln Tyr Pro Asn Lys Asn Leu Leu
      180            185            190

```

11/57

Pro Val Leu His Phe Glu Lys Val Val Cys Ser Thr Leu Lys Lys Asn
195 200 205

Ala Lys Arg Asn Pro Lys Val Asp Arg Glu Met Leu Lys Glu Ile Ile
210 215 220

Ala Arg Gly Asn Phe Arg Leu Gln Asn Ile Ile Gly Arg Lys Met Gly
225 230 235 240

Leu Glu Cys Val Asp Ile Leu Ser Glu Leu Phe Arg Arg Gly Leu Arg
245 250 255

His Val Leu Ala Thr Ile Leu Ala Gln Leu Ser Asp Met Asp Leu Ile
260 265 270

Asn Val Ser Lys Val Ser Thr Thr Trp Lys Lys Ile Leu Glu Asp Asp
275 280 285

Lys Gly Ala Phe Gln Leu Tyr Ser Lys Ala Ile Gln Arg Val Thr Glu
290 295 300

Asn Asn Asn Lys Phe Ser Pro His Ala Ser Thr Arg Glu Tyr Val Met
305 310 315 320

Phe Arg Thr Pro Leu Ala Ser Val Gln Lys Ser Ala Ala Gln Thr Ser
325 330 335

Leu Lys Lys Asp Ala Gln Thr Lys Leu Ser Asn Gln Gly Asp Gln Lys
340 345 350

Gly Ser Thr Tyr Ser Arg His Asn Glu Phe Ser Glu Val Ala Lys Thr
355 360 365

Leu Lys Lys Asn Glu Ser Leu Lys Ala Cys Ile Arg Cys Asn Ser Pro
370 375 380

Ala Lys Tyr Asp Cys Tyr Leu Gln Arg Ala Thr Cys Lys Arg Glu Gly
385 390 395 400

Cys Gly Phe Asp Tyr Cys Thr Lys Cys Leu Cys Asn Tyr His Thr Thr
405 410 415

Lys Asp Cys Ser Asp Gly Lys Leu Leu Lys Ala Ser Cys Lys Ile Gly
420 425 430

Pro Leu Pro Gly Thr Lys Lys Ser Lys Lys Asn Leu Arg Arg Leu
435 440 445

<210> 11
 <211> 1535
 <212> DNA
 <213> Homo sapiens

<400> 11
 gcgcgttcgg gagcttcggc cctgcgtagg aggcgggtgc aggtgtgggt gctgagccgc 60
 ccgcgcctcg gagggggaga cagcttcagg acacgcaggc cgcagcgagg gcccgggccc 120
 gggggatccc aggcacatga cgtcccccac tccaaagcag ccttggacag cattaacgag 180
 ctgcccagata acatcctgct ggagctgttc acgcacgtgc ccgcccggca gctgctgctg 240
 aactgcccgc tggctctgcag cctctggcgg gacctcatcg acctcctgac cctctggaaa 300
 cgcaagtgc tgcgaaaggg ctccatcacc aaggactggg accagcccggt ggcgcactgg 360
 aaaatcttct acttctacg gagcctgcat aggaacctcc tgcgcaacct gtgtgctgaa 420
 aacgatatgt ttgcatggca aattgatttc aatggtgggg accgctggaa ggtggatagc 480

12/57

```

ctccctggag cccacgggac agaatttctt gaccccaaag tcaagaagtc ttttgtcaca 540
tctacgaac tgtgctcaa gtgggagctg gtggaccttc tagccgaccg ctactgggag 600
gagctactag acacattccg gccggacatc gtggttaagg actgggtttgc tgccagagcc 660
gactgtggct gcacctacca actcaaatg agctggcct cggttgacta cttcgtgttg 720
gctcctctcg agccccacc tgtgaccatc caacagtggg acaatgccac atggacagag 780
gtctcttaca cttctcaga ctacccccgg ggtgtccgct acatcctctt ccagcatggg 840
ggcaggggaca cccagtactg ggcaggctgg tatgggcccc gagtccacaa cagcagcatt 900
gtcgtcagcc ccaagatgac caggaaccag gcctcgccg aggcacagcc tgggcagaag 960
catggacagg aggaggctgc ccaatcgccc tacggagctg ttgtccagat tttctgacag 1020
ctgtccatcc tgtgtctggg tcagccagag gttcctccag gcaggagctg agcatgggt 1080
gggcagtgaq gtccctgtac cagcgactcc tgccccgggt caacctacc agcttgtgg 1140
aacttaactgt cacatagctc tgacgttttg ttgtaataaa tgttttcagg ccgggactg 1200
tggtcagcgc ctgtaatccc agcactttg gagaccgagg caggtggatc acgaggtcag 1260
gagacagaga ccactctggc caacacggtg aaacctgtc tctactaaaa atacaaaaaa 1320
ttagccgggc gtggtggcgg gcgcctgtag tcccagctac tcgggaggct gatgcagaag 1380
aatggcgtga acccggaagg cagagcttgc agtgagccga gatcacgcca ctgactcca 1440
gcctgggtga cagagcgaga ctctggtca taaaataata ataataata ataaataaaa 1500
ataaatggt tttcagtaaa aaaaaaaaaa aaaaa 1535

```

<210> 12

<211> 338

<212> PRT

<213> Homo sapiens

<400> 12

```

Ala Arg Ser Gly Ala Ser Ala Leu Arg Arg Arg Val Gln Val Trp
  1           5           10           15
Val Leu Ser Arg Pro Pro Pro Gly Gly Gly Asp Ser Phe Arg Thr Arg
          20           25           30
Arg Pro Gln Arg Gly Pro Gly Pro Gly Gly Ser Gln Ala Met Asp Ala
          35           40           45
Pro His Ser Lys Ala Ala Leu Asp Ser Ile Asn Glu Leu Pro Asp Asn
          50           55           60
Ile Leu Leu Glu Leu Phe Thr His Val Pro Ala Arg Gln Leu Leu Leu
          65           70           75           80
Asn Cys Arg Leu Val Cys Ser Leu Trp Arg Asp Leu Ile Asp Leu Leu
          85           90           95
Thr Leu Trp Lys Arg Lys Cys Leu Arg Lys Gly Phe Ile Thr Lys Asp
          100          105          110
Trp Asp Gln Pro Val Ala Asp Trp Lys Ile Phe Tyr Phe Leu Arg Ser
          115          120          125
Leu His Arg Asn Leu Leu Arg Asn Pro Cys Ala Glu Asn Asp Met Phe
          130          135          140
Ala Trp Gln Ile Asp Phe Asn Gly Gly Asp Arg Trp Lys Val Asp Ser
          145          150          155          160
Leu Pro Gly Ala His Gly Thr Glu Phe Pro Asp Pro Lys Val Lys Lys
          165          170          175
Ser Phe Val Thr Ser Tyr Glu Leu Cys Leu Lys Trp Glu Leu Val Asp
          180          185          190
Leu Leu Ala Asp Arg Tyr Trp Glu Glu Leu Leu Asp Thr Phe Arg Pro
          195          200          205

```

13/57

Asp Ile Val Val Lys Asp Trp Phe Ala Ala Arg Ala Asp Cys Gly Cys
 210 215 220
 Thr Tyr Gln Leu Lys Val Gln Leu Ala Ser Ala Asp Tyr Phe Val Leu
 225 230 235 240
 Ala Ser Phe Glu Pro Pro Pro Val Thr Ile Gln Gln Trp Asn Asn Ala
 245 250 255
 Thr Trp Thr Glu Val Ser Tyr Thr Phe Ser Asp Tyr Pro Arg Gly Val
 260 265 270
 Arg Tyr Ile Leu Phe Gln His Gly Gly Arg Asp Thr Gln Tyr Trp Ala
 275 280 285
 Gly Trp Tyr Gly Pro Arg Val Thr Asn Ser Ser Ile Val Val Ser Pro
 290 295 300
 Lys Met Thr Arg Asn Gln Ala Ser Ser Glu Ala Gln Pro Gly Gln Lys
 305 310 315 320
 His Gly Gln Glu Glu Ala Ala Gln Ser Pro Tyr Gly Ala Val Val Gln
 325 330 335
 Ile Phe

<210> 13
 <211> 1763
 <212> DNA
 <213> Homo sapiens

<400> 13
 tgggaattccc atggaccatg tctaataccc gattlacaat tacattgaac tacaaggatc 60
 cccctacttg agatgaagag accttggctt catatgggat tgtttctggg gacttgatat 120
 gtttgattct tcacgatgac attccaccgc ctaatatacc ttcattccaca gattcagagc 180
 attcttcact ccagaacaal gagcaacct ctttggccac cagctccaat cagactagca 240
 tacaggatga acaaccaagt gattcattcc aaggacaggc agcccagtct ggtggttggg 300
 atgcacagag tatgttaggg cctagtcaaa attttgaagc tgagtcaatt caagataatg 360
 cgtcatgggc agagggcaca ggtttctatc cctcagaacc cctgctctgt agtgaatcgg 420
 tgggaaggca agtgccacat tcattagaga ccttgtatca atcagctgac tgttctgatg 480
 ccaatgatgc gttgatagtg ttgatacatc ttctcatggt ggagtcaggt tacataacct 540
 agggcaccca agccaaagca ctgtccctgc cggagaagtg gaagttgagc ggggtgtata 600
 agctgcagta catgcatcat ctctgcgagg gcagctccgc tactctcacc tgtgtgcctt 660
 tgggaaacct gattgttgta aatgctacac taataatcaa caatgagatt agaagtgtga 720
 aaagattgca gctgctacca gaatctttta ttgcaaaga gaaactaggg gaaaatgtag 780
 ccaacatata caaagatctt cagaaactct ctgcctctt taagaccag ctggtgtatc 840
 ctcttctggc ttttaccoga caagcaactga acctaccaaa tgtatttggg ttggtcgtcc 900
 tcccattgga actgaaacta cggatcttcc gacttctgga tgttctgtcc gtcttctctt 960
 tgtctgcggt ttgtcgtgac ctctttactg cttcaaatga cccactcctg tggaggtttt 1020
 tatatctgcg tgattttoga gacaatactg tcagagttca agacacagat tggaaagaac 1080
 tgtacaggaa gaggcacata caaagaaaag aatcccccga agggcggttt gtgctgtccc 1140
 tgccatcgct aacccacacc attccattct atcccaaccc cttgcacctt aggccatttc 1200
 ctagctcccg ccttctccca ggaattatcg ggggtgaata tgacaaaaga ccaacacttc 1260
 cctatgttgg agaccacaac agttcactca ttcttggtcc tggggagacg cccagccagt 1320
 tacctccact gagaccaagc ttgatccag ttggcccact tccaggacct aaccccatct 1380
 tgccagggcg aggcggcccc aatgacagat ttccctttag acccagcagg ggtcggccaa 1440
 ctgatggcg cctgtcattc atgtgattga ttgttaattt catttctgga gctccatttg 1500
 tttttgttcc taaactacag atgtcactcc ttgggtggt gatctcaggt gttattttct 1560
 gattgtggtg ttgagagttg cactcccaga aaccttttaa gagatacatt tatagcccta 1620
 ggggtggtat gaccaaaagg ttctctgtg acaagggttg ccttgggaat agttggctgc 1680
 caatctccct gctcttggtt ctctctaga ttgaagtttg ttttctgatg ctgttcttcc 1740

cagattaaaa aaaagtgttaa att

1763

<210> 14

<211> 482

<212> PRT

<213> Homo sapiens

<400> 14

Met Ser Asn Thr Arg Phe Thr Ile Thr Leu Asn Tyr Lys Asp Pro Leu
 1 5 10 15

Thr Gly Asp Glu Glu Thr Leu Ala Ser Tyr Gly Ile Val Ser Gly Asp
 20 25 30

Leu Ile Cys Leu Ile Leu His Asp Asp Ile Pro Pro Pro Asn Ile Pro
 35 40 45

Ser Ser Thr Asp Ser Glu His Ser Ser Leu Gln Asn Asn Glu Gln Pro
 50 55 60

Ser Leu Ala Thr Ser Ser Asn Gln Thr Ser Ile Gln Asp Glu Gln Pro
 65 70 75 80

Ser Asp Ser Phe Gln Gly Gln Ala Ala Gln Ser Gly Val Trp Asn Asp
 85 90 95

Asp Ser Met Leu Gly Pro Ser Gln Asn Phe Glu Ala Glu Ser Ile Gln
 100 105 110

Asp Asn Ala His Met Ala Glu Gly Thr Gly Phe Tyr Pro Ser Glu Pro
 115 120 125

Leu Leu Cys Ser Glu Ser Val Glu Gly Gln Val Pro His Ser Leu Glu
 130 135 140

Thr Leu Tyr Gln Ser Ala Asp Cys Ser Asp Ala Asn Asp Ala Leu Ile
 145 150 155 160

Val Leu Ile His Leu Leu Met Leu Glu Ser Gly Tyr Ile Pro Gln Gly
 165 170 175

Thr Glu Ala Lys Ala Leu Ser Leu Pro Glu Lys Trp Lys Leu Ser Gly
 180 185 190

Val Tyr Lys Leu Gln Tyr Met His His Leu Cys Glu Gly Ser Ser Ala
 195 200 205

Thr Leu Thr Cys Val Pro Leu Gly Asn Leu Ile Val Val Asn Ala Thr
 210 215 220

Leu Lys Ile Asn Asn Glu Ile Arg Ser Val Lys Arg Leu Gln Leu Leu
 225 230 235 240

Pro Glu Ser Phe Ile Cys Lys Glu Lys Leu Gly Glu Asn Val Ala Asn
 245 250 255

Ile Tyr Lys Asp Leu Gln Lys Leu Ser Arg Leu Phe Lys Asp Gln Leu
 260 265 270

Val Tyr Pro Leu Leu Ala Phe Thr Arg Gln Ala Leu Asn Leu Pro Asn
 275 280 285

Val Phe Gly Leu Val Val Leu Pro Leu Glu Leu Lys Leu Arg Ile Phe

290 295 300
 Arg Leu Leu Asp Val Arg Ser Val Leu Ser Leu Ser Ala Val Cys Arg
 305 310 315 320
 Asp Leu Phe Thr Ala Ser Asn Asp Pro Leu Leu Trp Arg Phe Leu Tyr
 325 330 335
 Leu Arg Asp Phe Arg Asp Asn Thr Val Arg Val Gln Asp Thr Asp Trp
 340 345 350
 Lys Glu Leu Tyr Arg Lys Arg His Ile Gln Arg Lys Glu Ser Pro Lys
 355 360 365
 Gly Arg Phe Val Leu Leu Leu Pro Ser Ser Thr His Thr Ile Pro Phe
 370 375 380
 Tyr Pro Asn Pro Leu His Pro Arg Pro Phe Pro Ser Ser Arg Leu Pro
 385 390 395 400
 Pro Gly Ile Ile Gly Gly Glu Tyr Asp Gln Arg Pro Thr Leu Pro Tyr
 405 410 415
 Val Gly Asp Pro Ile Ser Ser Leu Ile Pro Gly Pro Gly Glu Thr Pro
 420 425 430
 Ser Gln Leu Pro Pro Leu Arg Pro Arg Phe Asp Pro Val Gly Pro Leu
 435 440 445
 Pro Gly Pro Asn Pro Ile Leu Pro Gly Arg Gly Gly Pro Asn Asp Arg
 450 455 460
 Phe Pro Phe Arg Pro Ser Arg Gly Arg Pro Thr Asp Gly Arg Leu Ser
 465 470 475 480
 Phe Met

<210> 15
 <211> 43
 <212> PRT
 <213> Homo sapiens

<400> 15
 Leu Pro Ala Arg Gly Leu Asp His Ile Ala Glu Asn Ile Leu Ser Tyr
 1 5 10 15
 Leu Asp Ala Lys Ser Leu Cys Ala Ala Glu Leu Val Cys Lys Glu Trp
 20 25 30
 Tyr Arg Val Thr Ser Asp Gly Met Leu Trp Lys
 35 40

<210> 16
 <211> 40
 <212> PRT
 <213> Homo sapiens

<400> 16
 Leu Pro Leu Glu Leu Ser Phe Tyr Leu Leu Lys Trp Leu Asp Pro Gln
 1 5 10 15

16/57

Thr Leu Leu Thr Cys Cys Leu Val Ser Lys Gln Trp Asn Lys Val Ile
 20 25 30

Ser Ala Cys Thr Glu Val Trp Gln
 35 40

<210> 17
 <211> 39
 <212> PRT
 <213> Homo sapiens

<400> 17
 Leu Leu Gln Asp Ile Ile Leu Gln Val Phe Lys Tyr Leu Pro Leu Leu
 1 5 10 15

Asp Arg Ala His Ala Ser Gln Val Cys Arg Asn Trp Asn Gln Val Phe
 20 25 30

His Met Pro Asp Leu Trp Arg
 35

<210> 18
 <211> 39
 <212> PRT
 <213> Homo sapiens

<400> 18
 Leu Pro Ile Asp Val Gln Leu Tyr Ile Leu Ser Phe Leu Ser Pro His
 1 5 10 15

Asp Leu Cys Gln Leu Gly Ser Thr Asn His Tyr Trp Asn Glu Thr Val
 20 25 30

Arg Asn Pro Ile Leu Trp Arg
 35

<210> 19
 <211> 39
 <212> PRT
 <213> Homo sapiens

<400> 19
 Leu Arg His Val Leu Ala Thr Ile Leu Ala Gln Leu Ser Asp Met Asp
 1 5 10 15

Leu Ile Asn Val Ser Lys Val Ser Thr Trp Lys Lys Ile Leu Glu
 20 25 30

Asp Asp Lys Gly Ala Phe Gln
 35

<210> 20
 <211> 40
 <212> PRT
 <213> Homo sapiens

<400> 20
 Leu Pro Asp Asn Ile Leu Leu Glu Leu Phe Thr His Val Pro Ala Arg
 1 5 10 15

17/57

Gln Leu Leu Leu Asn Cys Arg Leu Val Cys Ser Leu Trp Arg Asp Leu
 20 25 30

Ile Asp Leu Leu Thr Leu Trp Lys
 35 40

<210> 21
 <211> 39
 <212> PRT
 <213> Homo sapiens

<400> 21
 Leu Pro Leu Glu Leu Lys Leu Arg Ile Phe Arg Leu Leu Asp Val Arg
 1 5 10 15

Ser Val Leu Ser Leu Ser Ala Val Cys Arg Asp Leu Phe Thr Ala Ser
 20 25 30

Asn Asp Pro Leu Leu Trp Arg
 35

<210> 22
 <211> 39
 <212> PRT
 <213> Homo sapiens

<400> 22
 Leu Pro Asp Glu Leu Leu Leu Gly Ile Phe Ser Cys Leu Cys Leu Pro
 1 5 10 15

Glu Leu Leu Lys Val Ser Gly Val Cys Lys Arg Trp Tyr Arg Leu Ala
 20 25 30

Ser Asp Glu Ser Leu Trp Gln
 35

<210> 23
 <211> 1323
 <212> DNA
 <213> Homo sapiens

<400> 23
 acattttcta atgtttacag aatgaagagg aacagtttat ctgttgagaa taaaattgtc 60
 cagttgtcag gagcagcgaa acagccaaaa gttgggttct acttttctct caaccagact 120
 catacacaca cggttcttct agactggggg agtttgccct accatgtagt attacaaatt 180
 ttccagtatc ttccctttact agatcggggc tgtgcatctt ctgtatgtag gaggtggaat 240
 gaagtttttc atattttctga cctttggaga aagtttgaat ttgaactgaa ccagtcagct 300
 acttcatctt ttaagtccac tcctcctgat ctcattcagc agatcattaa aaagcatttt 360
 gtcctatctt agtatgtcag ctttaagggt gacagtagcg ctgagtcagc agaagctgcc 420
 tgtgatatac tctctcagct ggtaaattgt tccatccaga ccttgggctt gatttcaaca 480
 gccaaagcaa gtttcatgaa tgtgtcggag totcattttg tgtcagcaat tacagtgtgt 540
 tttatcaact caaaatcatt atcatcaatc aaaattgaag atacaccagt ggatgatcct 600
 tcattgaaga ttcttgtggc caataatagt gacactctaa gactcccaa gatgagttagc 660
 tgtcctcatg ttctcatgga tgggaattctt tgtgtagctg accgttgtca aggccttaga 720
 gaactggcgt tgaattatta catcctaact gatgaacttt tccttgcact ctcaagcgag 780
 actcatgtta accttgaaca tcttcgaatt gatgttgtga gtgaaaatcc tggacagatt 840
 aaatttcatg ctgttaaaaa acacagttgg gatgcactta ttaaacattc ccctagagtt 900
 aatgttggtta tgcacttctt tctatatgaa gaggaattcg agacgttctt caaagaagaa 960
 acccctgtta ctcaacctta ttttggtcgt tcagtcagca aagtgggttt aggacgggta 1020
 ggtctcaact gtccctcagc gattgagtta gtggtgtgtg ctaatgatct tcagcctctt 1080

18/57

gataatgaac ttatttgtat tgcctgaacac tgtacaaacc taacagcctt gggcctcage 1140
 aaatgtgaag ttagctgcag tgccttcac aggtttgtaa gactgtgtga gagaaggtaa 1200
 acacagctct ctgtaatgga ggaagttttg atccctgatg aggattatag cctagatgaa 1260
 attcacactg aagctctcaa atacctggga agagtatggt tcctgatgt gatgcctctc 1320
 tgg 1323

<210> 24

<211> 434

<212> PRT

<213> Homo sapiens

<400> 24

Met Lys Arg Asn Ser Leu Ser Val Glu Asn Lys Ile Val Gln Leu Ser
 1 5 10 15
 Gly Ala Ala Lys Gln Pro Lys Val Gly Phe Tyr Ser Ser Leu Asn Gln
 20 25 30
 Thr His Thr His Thr Val Leu Leu Asp Trp Gly Ser Leu Pro His His
 35 40 45
 Val Val Leu Gln Ile Phe Gln Tyr Leu Pro Leu Leu Asp Arg Ala Cys
 50 55 60
 Ala Ser Ser Val Cys Arg Arg Trp Asn Glu Val Phe His Ile Ser Asp
 65 70 75 80
 Leu Trp Arg Lys Phe Glu Phe Glu Leu Asn Gln Ser Ala Thr Ser Ser
 85 90 95
 Phe Lys Ser Thr His Pro Asp Leu Ile Gln Gln Ile Ile Lys Lys His
 100 105 110
 Phe Ala His Leu Gln Tyr Val Ser Phe Lys Val Asp Ser Ser Ala Glu
 115 120 125
 Ser Ala Glu Ala Ala Cys Asp Ile Leu Ser Gln Leu Val Asn Cys Ser
 130 135 140
 Ile Gln Thr Leu Gly Leu Ile Ser Thr Ala Lys Pro Ser Phe Met Asn
 145 150 155 160
 Val Ser Glu Ser His Phe Val Ser Ala Leu Thr Val Val Phe Ile Asn
 165 170 175
 Ser Lys Ser Leu Ser Ser Ile Lys Ile Glu Asp Thr Pro Val Asp Asp
 180 185 190
 Pro Ser Leu Lys Ile Leu Val Ala Asn Asn Ser Asp Thr Leu Arg Leu
 195 200 205
 Pro Lys Met Ser Ser Cys Pro His Val Ser Ser Asp Gly Ile Leu Cys
 210 215 220
 Val Ala Asp Arg Cys Gln Gly Leu Arg Glu Leu Ala Leu Asn Tyr Tyr
 225 230 235 240
 Ile Leu Thr Asp Glu Leu Phe Leu Ala Leu Ser Ser Glu Thr His Val
 245 250 255
 Asn Leu Glu His Leu Arg Ile Asp Val Val Ser Glu Asn Pro Gly Gln
 260 265 270

19/57

Ile Lys Phe His Ala Val Lys Lys His Ser Trp Asp Ala Leu Ile Lys
 275 280 285
 His Ser Pro Arg Val Asn Val Val Met His Phe Phe Leu Tyr Glu Glu
 290 295 300
 Glu Phe Glu Thr Phe Phe Lys Glu Glu Thr Pro Val Thr His Leu Tyr
 305 310 315 320
 Phe Gly Arg Ser Val Ser Lys Val Val Leu Gly Arg Val Gly Leu Asn
 325 330 335
 Cys Pro Arg Leu Ile Glu Leu Val Val Cys Ala Asn Asp Leu Gln Pro
 340 345 350
 Leu Asp Asn Glu Leu Ile Cys Ile Ala Glu His Cys Thr Asn Leu Thr
 355 360 365
 Ala Leu Gly Leu Ser Lys Cys Glu Val Ser Cys Ser Ala Phe Ile Arg
 370 375 380
 Phe Val Arg Leu Cys Glu Arg Arg Leu Thr Gln Leu Ser Val Met Glu
 385 390 395 400
 Glu Val Leu Ile Pro Asp Glu Asp Tyr Ser Leu Asp Glu Ile His Thr
 405 410 415
 Glu Val Ser Lys Tyr Leu Gly Arg Val Trp Phe Pro Asp Val Met Pro
 420 425 430
 Leu Trp

<210> 25
 <211> 1970
 <212> DNA
 <213> Homo sapiens

<400> 25
 ggaaacgtca aaattgggat agtcggcagt tctggcccoct gcagctggag gtaccctgag 60
 lictgagggt cgtagtgtctg ttcttggtat tctcatcgcg gtcacctcta ccggtgtgga 120
 caagtaaaagt ttgaatcagc ttctccatgg cctgggcacc agttcccggc tgagccattt 180
 tctttttggc taaaagtcct cgcacagagg ccaattcgtc gcggcggcgg tggagatcgc 240
 aggtcgctca ggcttcgaga tgggtcaagg gttgtggaga gtggtcagaa accagcagct 300
 gcaacaagaa ggctacagtg agcaaggcta cctcaccaga gacagagca ggagaatggc 360
 tgcgagcaac atttctaaca ccaatcatcg taaacaagtc caaggaggca ttgacatata 420
 tcatcttttg aaggcaagga aatcgaaaga acaggaagga ttcattaatt tggaaatggt 480
 gctcctgag ctaagcttta ccatcttgct ctacctgaat gcaactgacc ttgcttggc 540
 ttcatgtgtt tggcaggacc ttgcgaatga tgaactctc tggcaagggt tgtgcaaatc 600
 cacttggggt cactgttcca tatacaataa gaaccacct ttaggatttt cttttagaaa 660
 aktgtatatg cagctgggat aaggcagcct cacttttaat gccaacccag atgagggagt 720
 gaactacttt atgtccaagg gtatctctgga tgattcgcca aaggaaatag caaagtttat 780
 ctctgttaca agaactactaa attggaaaaa actgagaatc tatcttgatg aaaggagaga 840
 tgtcttgatg gacctgttaa cattgcataa ttttagaaaat cagttcttgc caaatgcact 900
 gagagaattt ttctgtcata tccatgcccc tgaagagcgt ggagagtatc ttgaaactct 960
 tataacaag ttctcacata gattctgtgc ttgcaacctt gatttaatgc gagaacttgg 1020
 ccttagtctt gatgtgtct atgtactgtg ctactctttg attctacttt ccattgacct 1080
 cactagccct catgtgaaga ataaaatgtc aaaaagggaa tttattcgaa ataccgctcg 1140
 cgcgtgtcaa aatattagtg aagattttgt agggcatctt tatgacaata tctaccttat 1200
 tggccatgtg gctgcataaa aagcacaatt gctaggactt cagtttttac ttcagactaa 1260
 agctacccaa ggacttagca gatatggggg ttacatcagt gctggtcatt gtagcctgag 1320
 tataacaatc agcttcagtg tgcaaccttt tttcttttg ccattttcta ttttagtaat 1380

20/57

```

ttccttgggg aactaaataa ttttgcagaa tttttcctaa ttttgtttat cacgttttgc 1440
acaaagcaga gccactgtct aacacagctg ttaacgaatg ataaactgac attatactct 1500
aaaagatggt gtatttgggc attagatttg cctgaaaaac tttatccatt tccattcttt 1560
atacaaatac catgtaatgt gtacatattt aactaaagag atttatagtc ataattattt 1620
tattgtaaag attttaacta aagtttttcc tttctctca aactgagttc tgaaatttat 1680
ttgattctga tctgaaacta ttgtctycgt aaaagttaga tctgacttca gtcagaaacc 1740
aataccagct tcttttttct ttaaaacttg aagagtgttg atttgttact atattactat 1800
gcaaaactgg cagtattttt tataatataa atttataatt tgatttttta ttttaaaaaac 1860
tgggttaatc aagtcctggg aagtccttta aaccatttag gatttttaaa acatcaaaat 1920
ttatgattta cattcatagg aataaaataa aatatyatta gaactctggt 1970

```

<210> 26

<211> 634

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> all Xaa positions

<223> Xaa=unknown amino acid residue

<400> 26

```

Glu Thr Ser Lys Leu Gly Ser Ala Val Leu Ala Pro Ala Ala Gly Gly
 1             5             10             15

Thr Leu Ser Ser Glu Gly Arg Ser Ala Val Ser Gly Ile Leu Ile Ala
          20             25             30

Val Thr Ser Thr Gly Val Asp Lys Ser Leu Asn Gln Leu Leu His Gly
      35             40             45

Leu Gly Thr Ser Ser Arg Leu Ser His Phe Pro Phe Gly Lys Ser Pro
 50             55             60

Pro Arg Gly Gln Phe Val Ala Ala Ala Val Glu Ile Ala Gly Arg Ser
 65             70             75             80

Gly Leu Gln Met Gly Gln Gly Leu Trp Arg Val Val Arg Asn Gln Gln
          85             90             95

Leu Gln Gln Glu Gly Tyr Ser Glu Gln Gly Tyr Leu Thr Arg Glu Gln
 100             105             110

Ser Arg Arg Met Ala Ala Ser Asn Ile Ser Asn Thr Asn His Arg Lys
 115             120             125

Gln Val Gln Gly Gly Ile Asp Ile Tyr His Leu Leu Lys Ala Arg Lys
 130             135             140

Ser Lys Glu Gln Glu Gly Phe Ile Asn Leu Glu Met Leu Pro Pro Glu
 145             150             155             160

Leu Ser Phe Thr Ile Leu Ser Tyr Leu Asn Ala Thr Asp Leu Cys Leu
 165             170             175

Ala Ser Cys Val Trp Gln Asp Leu Ala Asn Asp Glu Leu Leu Trp Gln
 180             185             190

Gly Leu Cys Lys Ser Thr Trp Gly His Cys Ser Ile Tyr Asn Lys Asn
 195             200             205

Pro Pro Leu Gly Phe Ser Phe Arg Lys Xaa Tyr Met Gln Leu Asp Glu
 210             215             220

```

21/57

Gly Ser Leu Thr Phe Asn Ala Asn Pro Asp Glu Gly Val Asn Tyr Phe
 225 230 235 240
 Met Ser Lys Gly Ile Leu Asp Asp Ser Pro Lys Glu Ile Ala Lys Phe
 245 250 255
 Ile Phe Cys Thr Arg Thr Leu Asn Trp Lys Lys Leu Arg Ile Tyr Leu
 260 265 270
 Asp Glu Arg Arg Asp Val Leu Asp Asp Leu Val Thr Leu His Asn Phe
 275 280 285
 Arg Asn Gln Phe Leu Pro Asn Ala Leu Arg Glu Phe Phe Arg His Ile
 290 295 300
 His Ala Pro Glu Glu Arg Gly Glu Tyr Leu Glu Thr Leu Ile Thr Lys
 305 310 315 320
 Phe Ser His Arg Phe Cys Ala Cys Asn Pro Asp Leu Met Arg Glu Leu
 325 330 335
 Gly Leu Ser Pro Asp Ala Val Tyr Val Leu Cys Tyr Ser Leu Ile Leu
 340 345 350
 Leu Ser Ile Asp Leu Thr Ser Pro His Val Lys Asn Lys Met Ser Lys
 355 360 365
 Arg Glu Phe Ile Arg Asn Thr Arg Arg Ala Ala Gln Asn Ile Ser Glu
 370 375 380
 Asp Phe Val Gly His Leu Tyr Asp Asn Ile Tyr Leu Ile Gly His Val
 385 390 395 400
 Ala Ala Lys Ala Gln Leu Leu Gly Leu Gln Phe Leu Leu Gln Thr Lys
 405 410 415
 Ala Thr Gln Gly Leu Ser Arg Tyr Gly Gly Tyr Ile Ser Ala Gly His
 420 425 430
 Cys Ser Leu Ser Ile Gln Ser Ser Phe Ser Val Gln Pro Phe Phe Leu
 435 440 445
 Leu Pro Phe Ser Ile Leu Val Ile Ser Leu Gly Asn Ile Ile Leu Gln
 450 455 460
 Asn Phe Ser Phe Cys Leu Ser Arg Phe Ala Gln Ser Arg Ala Thr Val
 465 470 475 480
 His Ser Cys Arg Met Ile Asn His Tyr Thr Leu Lys Asp Gly Val Phe
 485 490 495
 Val His Ile Cys Leu Lys Asn Phe Ile His Phe His Ser Leu Tyr Lys
 500 505 510
 Tyr His Val Met Cys Thr Tyr Leu Thr Lys Glu Ile Tyr Ser His Asn
 515 520 525
 Tyr Phe Ile Val Lys Ile Leu Thr Lys Val Phe Pro Phe Leu Ser Asn
 530 535 540
 Val Leu Lys Phe Ile Phe Ser Glu Thr Ile Val Xaa Val Lys Val Arg
 545 550 555 560

22/57

Ser Asp Phe Arg Gln Lys Pro Ile Pro Ala Ser Phe Ser Phe Lys Leu
565 570 575

Arg Val Leu Ile Cys Tyr Tyr Ile Thr Met Gln Asn Trp Gln Leu Phe
580 585 590

Leu Tyr Lys Phe Ile Ile Phe Phe Ile Leu Lys Thr Gly Leu Ile Lys
595 600 605

Ser Arg Val Leu Thr Ile Asp Phe Asn Ile Lys Ile Tyr Asp Leu His
610 615 620

Ser Glu Asn Lys Ile Xaa Leu Glu Leu Trp
625 630

<210> 27

<211> 4168

<212> DNA

<213> Homo sapiens

<400> 27

```

gatggcggcg gcagcagtcg acagcgcgat ggaggtggtg ccggcgctgg cggaggaggc 60
cgcgccggag gtagcgggcc tcagctgcct cgccaacctg ccgggtgagg tgctggagta 120
catcctgtgc tgcggctgcg tgacggccgc cgacatcgcc cgtgtctcca gcacctgccg 180
gcggtgcgcg gagctgtgcc agagcagcgg gaaggtgtgg aaggagcagt tccgggtgag 240
gtggccttcc ctatgaaac actacagccc caccgactac gtcaattggg tggaagagta 300
taaagttcgg caaaaagctg ggttagaagc gcggaagatt gtagcctcgt tctcaaagag 360
gttcttttca gagcacgttc cttgtaatgg cttcagtgac attgagaacc ttgaaggacc 420
agagattttt tttgaggatg aactggtgtg tatcctaaat atggaaggaa gaaaagcttt 480
gacctggaaa tactacgcaa aaaaaattct ttactacctg cggcaacaga agatcttaaa 540
taatcttaag gcctttcttc agcagccaga tgactatgag tcttatcttg aagggtgctgt 600
atatattgac cagtactgca atcctctctc cgacatcagc ctcaaagaca tccaggccca 660
aattgacagc atcgtggagc ttgtttgcaa aacctctcgg gccataaaca gtgcgccccc 720
cagcttggcc ttcaaggcag gtgaatcatc catgataatg gaaatagaac tccagagcca 780
ggtgctggat gccatgaact atgtccttta cgaccaactg aagttcaagg ggaatcgaat 840
ggataactat aatgcctcca acttatatat gcactcaggtt ttgattcgca gaacaggaat 900
cccaatcagc atgtctctgc tctatttgac aattgctcgg cagttgggag tccacttga 960
gcctgtcaac ttcccaagtc acttcttatt aaggtggtgc caaggcgcag aaggggcgac 1020
cctggacatc tttgactaca tctacataga tgcttttggg aaaggcaagc agctgacagt 1080
gaaagaatgc gactacttga tgggccagca cgtgactgca gcactgtatg ggggtgtcaa 1140
tgtcaagaag gtgtttacaga gaatggtggg aaacctgtta agcctgggga agcgggaagg 1200
catcgaccag tcataccagc tccagagaga ctgcgtggat ctctatctgg caatgtacc 1260
ggaccaggtg cagctctctc tccccaagc caggctttac ttccacctgg gaatctggcc 1320
agagaaggtg cttgacatcc tccagcacat ccaaacccca gaccgggggc agcacggggc 1380
ggtgggctac ctggtgcagc aactctaga gcacattgag cgcaaaaagg aggaggtggg 1440
cgtagaggtg aagctgcgct ccgatgagaa gcacagagat gtctgctact ccacgggct 1500
cattatgaag cataagaggt atggctataa ctgtgtgac tacggtctgg accccacctg 1560
catgatggga cagcagtgga tccggaacat gaacgtccac agcctgccgc acggccacca 1620
ccagccttcc tataacgtgc tgggtggagga cggctcctgt cgatacgag cccaagaaaa 1680
cttggaatat aacgtggagc ctcaagaaat ctcacaacct gacgtgggac gctatttctc 1740
agagtttact ggcactcact acatcccaaa cgcagagctg gagatccggt atccagaaga 1800
tctggagttt gtctatgaaa cgggtgcagaa tatttacagt gcaaagaaag agaacataga 1860
tgagtaaaat ctagagagga cattgcaact ttgctgctgc tgctatcttc caagagaacg 1920
ggactccgga agaagacgtc tccacggagc cctcgggacc tgctgcacca ggaaagccac 1980
tccaccagta gtctgggtg cctcctacta agtttaaata cgtgtgtctc ttcccccagt 2040
gcaaagacaa tgtgtctctc cgcctacact agtgaattaa tctgaaaggc actgtgtcag 2100
tggcatggct tgtatgcttg tccgtgtgtg acagtttgtg acattctgtc ttcatgaggt 2160
ctcacagtcg acgctcctgt aatcattctt tgtattcact ccattccctt gtctgtctgc 2220
atttgtctca gaacatttcc ttgctgggac agatgggggt atgcatttgc aataatttcc 2280
ttctgatttc tctgtggaac gtgttcggtc ccgagtgagg actgtgtgtc tttttaccct 2340
gaagttagtt gcataattcag aggtlaaagt gtgtgtctac ttggcagcat cttagagatg 2400
gagacattaa caagctaatt gtaattagaa tcatttgaat ttattttttt ctaatatgtg 2460

```

23/57

```

aaacacagat ttcaagtgtt ttatcttttt tttttaaatt taaatgggaa tataacacag 2520
ttttcccttc catattccctc tottgagttt atgcacatct ctataaatca ttagttttct 2580
attttattac ataaaaattct tttagnaaat gcaaatagtg aactttgtga atggattttt 2640
ccatactcat ctacaattcc tocattttaa atgactactt ttatttttta atttaaaaaa 2700
tctacttcag tatcatgagt aggtctttaca tcagtgatgg gttctttttg tagtgagaca 2760
tacaaatctg atgttaalgt ttgtctttag aagtcatact ccatgggtctt caaagaccaa 2820
aaaatgaggt tttgcttttg taatcaggaa aaaaaaaaat taatgaacct taaaaaaaaa 2880
aaaaaagggt ttgaagggaa aaaaagtggg ttcacacctc ttgttattcc ttagagtcac 2940
ttcaaggcct gtttgaatgt ggcaggttag aaagagagag aatgtctttc atttgaagag 3000
tgttggactt gtgtgaaagg agatgtgcgt gttggaatct gcttttccaa gcgcgcaggg 3060
tcttgacggc agcaggacga agcctgttgt ggcgtcttct gggaaagcct gaccgtgtgt 3120
tcggacggca ctggctcctt tccgaagtto tcagttaact agcccagagt aactgcacgc 3180
ctttgtgcag ctctggagct ccaccaactc tcggcctgcc agttctcaag cgagctaata 3240
ttgttcataa tcgtagaagg ctaacttccg aagttaggac ctagttaact tgctctcaac 3300
atttaaaata atgcagttgc totagtgaat ggggcgttag gggcctgtct ctgcacctgt 3360
ctgtccatct gcatgcagta ttctcaccac tgttgaatgc ctgctgcttg tttacctttt 3420
ggaaaccctg ggtgaccaa ggtttggaaa gccacctgag accacttcat agcaagggaa 3480
ggctttaagc agttactaga aagagatggg gatttggccc cnggctcctc cagcctgaat 3540
gagctattta atccactgtc catgttccct atcagtcaaa tccaaagtca aaggatttga 3600
acctgcctct ggaaacgtaa ccactcacag cacttgccc gccaaaggtg ggaggattgt 3660
acactacttt catttaaagg ggaaagtttg ataatacgga attaatlaat atgaatgaga 3720
tgcattaata agaacctgag catgctgaga gttgcaattg ttggttttct ggtttgattg 3780
atttcccttt ttcttagaca catcaaagtc aagaaagatg gttttacctt tactgaccca 3840
gctgtacata tgtatctaga ctgtttttaa atgtctttct tcatgaatgc ttcattgggc 3900
tccaggaagc ctgtatcacc tgtgtaagtt ggtatttggg cactttalat ttttctaaaa 3960
acgtgttttg gatcctgtac tctaataaat cataagtttc tttttaaaaa ttttccaaaa 4020
cttttctcca ttttaaaaag cctgtttata aacgttgaac ttccacaatg ttaaaatgtt 4080
aaatatttgg atatagcaac ttcttttctc tlcaaatagaa tgccaagatt tttttgtaca 4140
atgattaata aatggaactt atccagag                                     4168

```

<210> 28
 <211> 621
 <212> PRT
 <213> Homo sapiens

<400> 28
 Met Ala Ala Ala Val Asp Ser Ala Met Glu Val Val Pro Ala Leu
 1 5 10 15
 Ala Glu Glu Ala Ala Pro Glu Val Ala Gly Leu Ser Cys Leu Val Asn
 20 25 30
 Leu Pro Gly Glu Val Leu Glu Tyr Ile Leu Cys Cys Gly Ser Leu Thr
 35 40 45
 Ala Ala Asp Ile Gly Arg Val Ser Ser Thr Cys Arg Arg Leu Arg Glu
 50 55 60
 Leu Cys Gln Ser Ser Gly Lys Val Trp Lys Glu Gln Phe Arg Val Arg
 65 70 75 80
 Trp Pro Ser Leu Met Lys His Tyr Ser Pro Thr Asp Tyr Val Asn Trp
 85 90 95
 Leu Glu Glu Tyr Lys Val Arg Gln Lys Ala Gly Leu Glu Ala Arg Lys
 100 105 110
 Ile Val Ala Ser Phe Ser Lys Arg Phe Phe Ser Glu His Val Pro Cys
 115 120 125
 Asn Gly Phe Ser Asp Ile Glu Asn Leu Glu Gly Pro Glu Ile Phe Phe
 130 135 140

24/57

Glu Asp Glu Leu Val Cys Ile Leu Asn Met Glu Gly Arg Lys Ala Leu
 145 150 155 160
 Thr Trp Lys Tyr Tyr Ala Lys Lys Ile Leu Tyr Tyr Leu Arg Gln Gln
 165 170 175
 Lys Ile Leu Asn Asn Leu Lys Ala Phe Leu Gln Gln Pro Asp Asp Tyr
 180 185 190
 Glu Ser Tyr Leu Glu Gly Ala Val Tyr Ile Asp Gln Tyr Cys Asn Pro
 195 200 205
 Leu Ser Asp Ile Ser Leu Lys Asp Ile Gln Ala Gln Ile Asp Ser Ile
 210 215 220
 Val Glu Leu Val Cys Lys Thr Leu Arg Gly Ile Asn Ser Arg His Pro
 225 230 235 240
 Ser Leu Ala Phe Lys Ala Gly Glu Ser Ser Met Ile Met Glu Ile Glu
 245 250 255
 Leu Gln Ser Gln Val Leu Asp Ala Met Asn Tyr Val Leu Tyr Asp Gln
 260 265 270
 Leu Lys Phe Lys Gly Asn Arg Met Asp Tyr Tyr Asn Ala Leu Asn Leu
 275 280 285
 Tyr Met His Gln Val Leu Ile Arg Arg Thr Gly Ile Pro Ile Ser Met
 290 295 300
 Ser Leu Leu Tyr Leu Thr Ile Ala Arg Gln Leu Gly Val Pro Leu Glu
 305 310 315 320
 Pro Val Asn Phe Pro Ser His Phe Leu Leu Arg Trp Cys Gln Gly Ala
 325 330 335
 Glu Gly Ala Thr Leu Asp Ile Phe Asp Tyr Ile Tyr Ile Asp Ala Phe
 340 345 350
 Gly Lys Gly Lys Gln Leu Thr Val Lys Glu Cys Glu Tyr Leu Ile Gly
 355 360 365
 Gln His Val Thr Ala Ala Leu Tyr Gly Val Val Asn Val Lys Lys Val
 370 375 380
 Leu Gln Arg Met Val Gly Asn Leu Leu Ser Leu Gly Lys Arg Glu Gly
 385 390 395 400
 Ile Asp Gln Ser Tyr Gln Leu Leu Arg Asp Ser Leu Asp Leu Tyr Leu
 405 410 415
 Ala Met Tyr Pro Asp Gln Val Gln Leu Leu Leu Leu Gln Ala Arg Leu
 420 425 430
 Tyr Phe His Leu Gly Ile Trp Pro Glu Lys Val Leu Asp Ile Leu Gln
 435 440 445
 His Ile Gln Thr Leu Asp Pro Gly Gln His Gly Ala Val Gly Tyr Leu
 450 455 460
 Val Gln His Thr Leu Glu His Ile Glu Arg Lys Lys Glu Glu Val Gly
 465 470 475 480

25/57

Val Glu Val Lys Leu Arg Ser Asp Glu Lys His Arg Asp Val Cys Tyr
 485 490 495

Ser Ile Gly Leu Ile Met Lys His Lys Arg Tyr Gly Tyr Asn Cys Val
 500 505 510

Ile Tyr Gly Trp Asp Pro Thr Cys Met Met Gly His Glu Trp Ile Arg
 515 520 525

Asn Met Asn Val His Ser Leu Pro His Gly His His Gln Pro Phe Tyr
 530 535 540

Asn Val Leu Val Glu Asp Gly Ser Cys Arg Tyr Ala Ala Gln Glu Asn
 545 550 555 560

Leu Glu Tyr Asn Val Glu Pro Gln Glu Ile Ser His Pro Asp Val Gly
 565 570 575

Arg Tyr Phe Ser Glu Phe Thr Gly Thr His Tyr Ile Pro Asn Ala Glu
 580 585 590

Leu Glu Ile Arg Tyr Pro Glu Asp Leu Glu Phe Val Tyr Glu Thr Val
 595 600 605

Gln Asn Ile Tyr Ser Ala Lys Lys Glu Asn Ile Asp Glu
 610 615 620

<210> 29
 <211> 278
 <212> DNA
 <213> Homo sapiens

<220>
 <221> modified_base
 <222> all n positions
 <223> n=a, c, g or t

<400> 29
 ccgtagtact ggnttcgggc gggtggtga ggaatggagc cggtagntgc ttgcggcgag 60
 tcccgggntc ctccgtagac ccgcgganac ctctgtgttg agtaacctgg cggaggtgt 120
 ggagcgtgtg ctcaccttc tgcgcgccaa ggcgttgttg cgggtggcct gcgtgtgccg 180
 cttatggagg gagtgtgtgc gcagagtatt gcggacccat cggagcgtaa cctggatctc 240
 cgcaggcctg gcggaggccg gccacctggn ggggcatt 278

<210> 30
 <211> 91
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> all Xaa positions
 <223> Xaa-unknown amino acid residue

<400> 30
 Arg Ser Thr Gly Phe Arg Arg Ala Gly Glu Glu Trp Ser Arg Xaa Leu
 1 5 10 15

Ala Ala Ser Pro Gly Xaa Leu Arg Arg Pro Ala Xaa Thr Phe Val Leu
 20 25 30

Ser Asn Leu Ala Glu Val Val Glu Arg Val Leu Thr Phe Leu Pro Ala

26/57

35 40 45

Lys Ala Leu Leu Arg Val Ala Cys Val Cys Arg Leu Trp Arg Glu Cys
50 55 60

Val Arg Arg Val Leu Arg Thr His Arg Ser Val Thr Trp Ile Ser Ala
65 70 75 80

Gly Leu Ala Glu Ala Gly His Leu Xaa Gly His
85 90

<210> 31
<211> 592
<212> DNA
<213> Homo sapiens

<400> 31

gcggccgcgc cccgtgcagc aacagcagca gcagccccc cagcagccgc cgcgcagacc 60
gccccagcag cagccgcccc agcagcagcc tcgcgcgcgc cgcgcagcagc agcagcagca 120
gcagccctccg ccgcgcgccac cgcgcgcctcc gccgtgcct caggagcggg acaacgtcgg 180
cgagcgggat gatgatgtgc ctgcagatat gttgcagaa gaatcaggtc ctggtgcaca 240
aaatagtcca taccaacttc gtagaaaaac tcttttgccg aaaagaacag cgtgtccacc 300
aaagaacagt atggaggcgc cctcaacttc aactacagaa aactttggtc atcgtgcaaa 360
acgtgcaaga gtgtctggaa aatcacaaga tctatcagca gcacctgctg aacagtatct 420
tcaggagaaa ctgccagatg aagtggttct aaaaatcttc tcttacttgc tggacagga 480
tctttgtaga gcagcttggtg tatgtaaacg cttcagtgaa cttgctaatt atcccaattt 540
gtggaaaacga ttatatatgg aagtatttga atatactcgc cctatgatgc at 592

<210> 32
<211> 197
<212> PRT
<213> Homo sapiens

<400> 32

Arg Pro Arg Pro Val Gln Gln Gln Gln Gln Gln Pro Pro Gln Gln Pro
1 5 10 15

Pro Pro Gln Pro Pro Gln Gln Gln Pro Pro Gln Gln Gln Pro Pro Pro
20 25 30

Pro Pro Gln Gln Gln Gln Gln Gln Gln Pro Pro Pro Pro Pro Pro Pro
35 40 45

Pro Pro Pro Leu Pro Gln Glu Arg Asn Asn Val Gly Glu Arg Asp Asp
50 55 60

Asp Val Pro Ala Asp Met Val Ala Glu Glu Ser Gly Pro Gly Ala Gln
65 70 75 80

Asn Ser Pro Tyr Gln Leu Arg Arg Lys Thr Leu Leu Pro Lys Arg Thr
85 90 95

Ala Cys Pro Thr Lys Asn Ser Met Glu Gly Ala Ser Thr Ser Thr Thr
100 105 110

Glu Asn Phe Gly His Arg Ala Lys Arg Ala Arg Val Ser Gly Lys Ser
115 120 125

Gln Asp Leu Ser Ala Ala Pro Ala Glu Gln Tyr Leu Gln Glu Lys Leu
130 135 140

Pro Asp Glu Val Val Leu Lys Ile Phe Ser Tyr Leu Leu Glu Gln Asp

PCT/US02/00311

[illegible]

```
<210> 33
<211> 537
<212> DNA
<213> Homo sapiens
```

<400>	33						
gcggcgcgcg	cccggaactcc	gcggtgggcy	agcgccctgt	gaggtgacca	tggaggctcg	60	
tggcctcccc	ttggagctgt	ggcgcatgat	cttagcctac	ttgcaccttc	cgcacctggg	120	
ccgctgcgac	ctgggtatgca	gggcctggta	tgaactgatc	ctcagttctg	acagaccggc	180	
ctggcgggcag	clgtgtctgg	gttgccaccga	gtgccgccat	cccaattggc	ccaaccagcc	240	
agatgtggag	cctgagtcct	ggagagaagc	cttcaagcag	cattaccttg	catccaagac	300	
atggaccaag	aatgccttgg	acttggagtc	tcccatctgc	ttttctctat	tccgccggag	360	
gaggggaacga	cgtaccctga	gtgtttgggc	aggccgtgag	tttgacagcc	tgggcagtcg	420	
cttggccatg	gccagcctgt	atgaccgaat	tgtgtctctc	ccaggtgtgt	acgaagagca	480	
aqctgaattc	atcttgaagc	tgccttgga	gattgtaggg	caggggaagt	tgggtga	537	

```
<210> 34
<211> 178
<212> PRT
<213> Homo sapiens
```

```

<400> 34
Arg Pro Arg Pro Gly Leu Arg Gly Gly Arg Ala Pro Cys Glu Val Thr
  1          5          10          15

Met Glu Ala Gly Gly Leu Pro Leu Glu Leu Trp Arg Met Ile Leu Ala
      20          25          30

Tyr Leu His Leu Pro Asp Leu Gly Arg Cys Ser Leu Val Cys Arg Ala
      35          40          45

Trp Tyr Glu Leu Ile Leu Ser Leu Asp Ser Thr Arg Trp Arg Gln Leu
      50          55          60

Cys Leu Gly Cys Thr Glu Cys Arg His Pro Asn Trp Pro Asn Gln Pro
      65          70          75          80

Asp Val Glu Pro Glu Ser Trp Arg Glu Ala Phe Lys Gln His Tyr Leu
      85          90          95

Ala Ser Lys Thr Trp Thr Lys Asn Ala Leu Asp Leu Glu Ser Ser Ile
      100          105          110

Cys Phe Ser Leu Phe Arg Arg Arg Glu Arg Arg Thr Leu Ser Val
      115          120          125

Gly Pro Gly Arg Glu Phe Asp Ser Leu Gly Ser Ala Leu Ala Met Ala
      130          135          140

Ser Leu Tyr Asp Arg Ile Val Leu Phe Pro Gly Val Tyr Glu Glu Gln
      145          150          155          160

```

28/57

Gly Glu Ile Ile Leu Lys Val Pro Val Glu Ile Val Gly Gln Gly Lys
 165 170 175

Leu Gly

<210> 35
 <211> 751
 <212> DNA
 <213> Homo sapiens

<400> 35
 gagaccgaga cggcgccgct gaccctagag tcgctgccca ccgatccct gctctccatc 60
 ttatcccttt tggactatcg ggatctaata aactgttggt atgtcagtcg aagattaagc 120
 cagctatcaa gtcctgatcc gctgtggaga agacattgca aaaaatactg gctgatattc 180
 gaggaagaga aaacacagaa gaatcagtgt tggaaatctc tcttcataga tacttactct 240
 gatgtaggaa gatacattga ccattatgct gctattaaaa aggcctcggg aatgatctca 300
 agaaatattt ggagcccagg tgcctcggga tgggttttat ctctgaaaga ggggtgctcg 360
 agaggaagac ctccatgctg tggaaagcga gattgggctg caagtttctt ggaagattat 420
 cgatgttcat accgaattca caatggacag aagttagttg gttcctgggg ttattgggaa 480
 gcatggcaat gtctaatac tatcgttctg aagatttggt agacgtcgat acagctgccg 540
 gagattccag cagagacagg gactgaaata ctgtctccct ttaacttttg catacatact 600
 gggttgagtc agtacatagc agtggaagct gcagagggtt gaaacaaaaa tgaagttttc 660
 taccaatgtc agacagtaga acgtgtgttt aaatatggca ttaagatgtg ttctgatggt 720
 tgtataaatg gcatgcatta ggtattttca g 751

<210> 36
 <211> 247
 <212> PRT
 <213> Homo sapiens

<400> 36
 Glu Thr Glu Thr Ala Pro Leu Thr Leu Glu Ser Leu Pro Thr Asp Pro
 1 5 10 15
 Leu Leu Leu Ile Leu Ser Phe Leu Asp Tyr Arg Asp Leu Ile Asn Cys
 20 25 30
 Cys Tyr Val Ser Arg Arg Leu Ser Gln Leu Ser Ser His Asp Pro Leu
 35 40 45
 Trp Arg Arg His Cys Lys Lys Tyr Trp Leu Ile Ser Glu Glu Glu Lys
 50 55 60
 Thr Gln Lys Asn Gln Cys Trp Lys Ser Leu Phe Ile Asp Thr Tyr Ser
 65 70 75 80
 Asp Val Gly Arg Tyr Ile Asp His Tyr Ala Ala Ile Lys Lys Ala Ser
 85 90 95
 Gly Met Ile Ser Arg Asn Ile Trp Ser Pro Gly Val Leu Gly Trp Val
 100 105 110
 Leu Ser Leu Lys Glu Gly Cys Ser Arg Gly Arg Pro Arg Cys Cys Gly
 115 120 125
 Ser Ala Asp Trp Ala Ala Ser Phe Leu Asp Asp Tyr Arg Cys Ser Tyr
 130 135 140
 Arg Ile His Asn Gly Gln Lys Leu Val Gly Ser Trp Gly Tyr Trp Glu
 145 150 155 160

29/57

Ala Trp His Cys Leu Ile Thr Ile Val Leu Lys Ile Cys Thr Ser Ile
 165 170 175

Gln Leu Pro Glu Ile Pro Ala Glu Thr Gly Thr Glu Ile Leu Ser Pro
 180 185 190

Phe Asn Phe Cys Ile His Thr Gly Leu Ser Gln Tyr Ile Ala Val Glu
 195 200 205

Ala Ala Glu Gly Asn Lys Asn Glu Val Phe Tyr Gln Cys Gln Thr Val
 210 215 220

Glu Arg Val Phe Lys Tyr Gly Ile Lys Met Cys Ser Asp Gly Cys Ile
 225 230 235 240

Asn Gly Met His Val Phe Ser
 245

<210> 37
 <211> 368
 <212> DNA
 <213> Homo sapiens

<220>
 <221> modified_base
 <222> all n positions
 <223> n=a, c, g or t

<400> 37
 ggctccgggtt tccggggccgg cgggtggcgg ctcaccatgc ccggnaaagca ccagcatttc 60
 caggaacctg aggtcggctg ctgcgggaaa tacttccgtg ttggcttcaa cattgtcttc 120
 tgggtgcteg gagccctgtt cctggetatc ggccctctgg cctgggggtga gaagggcgtt 180
 ctctcgaaca tctcagcgcg gacagatctg ggaggccttg acccctgtg gcttggttgt 240
 ggtagttgga ggcgtcatgt cgggtgctggg ctttgcctgg ctgcaattgg ggccctccgg 300
 gagaacacct tctgctcaa gttttctcnc gngttcctcg gtctcatctt ctccctggag 360
 ctggcaac 368

<210> 38
 <211> 122
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> all Xaa positions
 <223> Xaa=unknown amino acid residue

<400> 38
 Gly Ser Gly Phe Arg Ala Gly Gly Trp Pro Leu Thr Met Pro Gly Lys
 1 5 10 15

His Gln His Phe Gln Glu Pro Glu Val Gly Cys Cys Gly Lys Tyr Phe
 20 25 30

Leu Phe Gly Phe Asn Ile Val Phe Trp Val Leu Gly Ala Leu Phe Leu
 35 40 45

Ala Ile Gly Leu Trp Ala Trp Gly Glu Lys Gly Val Leu Ser Asn Ile
 50 55 60

Ser Ala Leu Thr Asp Leu Gly Gly Leu Asp Pro Val Trp Leu Val Cys
 65 70 75 80

30/57

Gly Ser Trp Arg Arg His Val Gly Ala Gly Leu Cys Trp Ala Ala Ile
85 90 95

Gly Ala Leu Arg Glu Asn Thr Phe Leu Leu Lys Phe Phe Xaa Xaa Phe
100 105 110

Leu Gly Leu Ile Phe Phe Leu Glu Leu Ala
115 120

<210> 39
<211> 774
<212> DNA
<213> Homo sapiens

<400> 39
gcgccggcgc cgcgcgcgta cctggacgag ctgcccgagc cgctgctgct gcgcgtgctg 60
gccgcactgc cggccgcgca gctgggtgcag gcctgcccgc tgggtgtgct gcgctggaag 120
gagctggttg acggcgcccc gctgtggctg ctcaagtgc agcaggaggg gctgggtgcc 180
gagggcgggc tggaggagga gcgcgaccac tggcagcagt tctacttct gagcaagcgg 240
cgccgcaacc ttctgcgtaa cccgtgtggg gaagaggact tggaaaggct gtgtgacgtg 300
gagcatggtg gggacggctg gaggtgtggag gagctgctg gagacagtgg ggtggagttc 360
accacagatg agagcgtaaa gaagtacttc gcctcctct ttgagtgggt tgcgaaaagca 420
caggtcattg acctgcaggc tgagggtac tgggaggagc tgctggacac gactcagccg 480
gccatcggtg tgaaggactg gtactcgggc cgcagcgacg ctggttgct ctacgagctc 540
accgttaagc tactgtccga gcacgagaac gtgctggctg agttcagcag cgggcaggtg 600
gcagtgcgcc aagacagtga cggcgggggc tggatggaga tctccacac ctccaccgac 660
tacgggcggg gcgtccgctt cgtccgcttc gagcacgggg ggcagggctc cgtctactlg 720
aagggtggt tgcgggcccg ggtgaccaac agcagcgtgt gggtagaacc ctga 774

<210> 40
<211> 257
<212> PRT
<213> Homo sapiens

<400> 40
Ala Ala Ala Ala Ala Tyr Leu Asp Glu Leu Pro Glu Pro Leu Leu
1 5 10 15
Leu Arg Val Leu Ala Ala Leu Pro Ala Ala Glu Leu Val Gln Ala Cys
20 25 30
Arg Leu Val Cys Leu Arg Trp Lys Glu Leu Val Asp Gly Ala Pro Leu
35 40 45
Trp Leu Leu Lys Cys Gln Gln Glu Gly Leu Val Pro Glu Gly Gly Val
50 55 60
Glu Glu Glu Arg Asp His Trp Gln Gln Phe Tyr Phe Leu Ser Lys Arg
65 70 75 80
Arg Arg Asn Leu Leu Arg Asn Pro Cys Gly Glu Glu Asp Leu Glu Gly
85 90 95
Trp Cys Asp Val Glu His Gly Gly Asp Gly Trp Arg Val Glu Glu Leu
100 105 110
Pro Gly Asp Ser Gly Val Glu Phe Thr His Asp Glu Ser Val Lys Lys
115 120 125
Tyr Phe Ala Ser Ser Phe Glu Trp Cys Arg Lys Ala Gln Val Ile Asp
130 135 140

31/57

Leu Gln Ala Glu Gly Tyr Trp Glu Glu Leu Leu Asp Thr Thr Gln Pro
 145 150 155 160
 Ala Ile Val Val Lys Asp Trp Tyr Ser Gly Arg Ser Asp Ala Gly Cys
 165 170 175
 Leu Tyr Glu Leu Thr Val Lys Leu Leu Ser Glu His Glu Asn Val Leu
 180 185 190
 Ala Glu Phe Ser Ser Gly Gln Val Ala Val Pro Gln Asp Ser Asp Gly
 195 200 205
 Gly Gly Trp Met Glu Ile Ser His Thr Phe Thr Asp Tyr Gly Pro Gly
 210 215 220
 Val Arg Phe Val Arg Phe Glu His Gly Gly Gln Gly Ser Val Tyr Trp
 225 230 235 240
 Lys Gly Trp Phe Gly Ala Arg Val Thr Asn Ser Ser Val Trp Val Glu
 245 250 255

Pro

<210> 41
 <211> 957
 <212> DNA
 <213> Homo sapiens

<400> 41
 atgggagcaga aggcgggtccc ttgtgctaagg agggaggcggg tgaagagaag ctgcccttct 60
 tgtggctcgg agcttgggggt tgaagagaag aggggggaaag gaaatccgat ttccatccag 120
 ttgttccccc cagagctgggt ggagcatatc atctcattcc tccagtcag agaccttgtt 180
 gccctcggcc agacctgccc ctacttccac gaagtgtgag atggggaagg cgtgtggaga 240
 cgcattctgt gcagactcag tccgcgcctc caagatcagg acacgaaggg cctgtatttc 300
 caggcatttg gaggccgccc ccgatgtctc agcaagagcg tggccccctt gctagccccc 360
 ggctaccgcc gcttcttgcc caccaaggat cactgtctca ttcttgacta cgtgggggacc 420
 ctcttcttcc tcaaaaatgc cctgggtctcc accctcggcc agatgcagtg gaagcgggcc 480
 ttgtcgtatg ttgtgttgtg tcgtggagcc aaggattttg cctcggaccc aaggtgtgac 540
 acagtttacc gtaaatacct ctaogtcttg gccactcggg agccgcagga agtgggtggg 600
 accaccagca gccggggcctg tgaetgtgtt gaggtctatc tgcagtctag tgggcagcgg 660
 gtcttcaaga tgacattcca ccaactcaatg accttcaagc agatcgtgct ggttggtcag 720
 gagaccagc gggctctact gctcctcaca taggaaggaa agatctactc ttgtgtagtg 780
 aatgagaccc agcttgacca gccacgctcc tacacggttc agctggccct gaggaagggt 840
 tcccactacc tgcctcaact gcgctgggcc tgcattgactt ccaaccagag cagcaccttc 900
 taogtcacag atcctattct gtgtctttgg ctacaaccac cttggcctgg tggatga 957

<210> 42
 <211> 318
 <212> PRT
 <213> Homo sapiens

<400> 42
 Met Gly Glu Lys Ala Val Pro Leu Leu Arg Arg Arg Arg Val Lys Arg
 1 5 10 15
 Ser Cys Pro Ser Cys Gly Ser Glu Leu Gly Val Glu Glu Lys Arg Gly
 20 25 30
 Lys Gly Asn Pro Ile Ser Ile Gln Leu Phe Pro Pro Glu Leu Val Glu
 35 40 45

His Ile Ile Ser Phe Leu Pro Val Arg Asp Leu Val Ala Leu Gly Gln
 30 55 60
 Thr Cys Arg Tyr Phe His Glu Val Cys Asp Gly Glu Gly Val Trp Arg
 65 70 75 80
 Arg Ile Cys Arg Arg Leu Ser Pro Arg Leu Gln Asp Gln Asp Thr Lys
 85 90 95
 Gly Leu Tyr Phe Gln Ala Phe Gly Gly Arg Arg Arg Cys Leu Ser Lys
 100 105 110
 Ser Val Ala Pro Leu Leu Ala His Gly Tyr Arg Arg Phe Leu Pro Thr
 115 120 125
 Lys Asp His Val Phe Ile Leu Asp Tyr Val Gly Thr Leu Phe Phe Leu
 130 135 140
 Lys Asn Ala Leu Val Ser Thr Leu Gly Gln Met Gln Trp Lys Arg Ala
 145 150 155 160
 Cys Arg Tyr Val Val Leu Cys Arg Gly Ala Lys Asp Phe Ala Ser Asp
 165 170 175
 Pro Arg Cys Asp Thr Val Tyr Arg Lys Tyr Leu Tyr Val Leu Ala Thr
 180 185 190
 Arg Glu Pro Gln Glu Val Val Gly Thr Thr Ser Ser Arg Ala Cys Asp
 195 200 205
 Cys Val Glu Val Tyr Leu Gln Ser Ser Gly Gln Arg Val Phe Lys Met
 210 215 220
 Thr Phe His His Ser Met Thr Phe Lys Gln Ile Val Leu Val Gly Gln
 225 230 235 240
 Glu Thr Gln Arg Ala Leu Leu Leu Leu Thr Glu Glu Gly Lys Ile Tyr
 245 250 255
 Ser Leu Val Val Asn Glu Thr Gln Leu Asp Gln Pro Arg Ser Tyr Thr
 260 265 270
 Val Gln Leu Ala Leu Arg Lys Val Ser His Tyr Leu Pro His Leu Arg
 275 280 285
 Val Ala Cys Met Thr Ser Asn Gln Ser Ser Thr Leu Tyr Val Thr Asp
 290 295 300
 Pro Ile Leu Cys Ser Trp Leu Gln Pro Pro Trp Pro Gly Gly
 305 310 315

<210> 43
 <211> 1590
 <212> DNA
 <213> Homo sapiens

<400> 43
 cgaggggggaa gcgaaggaag gggaagagga agggaaaagc gagcgagagg ggcaaggcgg 60
 aagaggaagc agggcggaag ggaagcccg gcgcagagcg gcgaaggagg cagcggggccg 120
 ggggctgagg cgggagcgag gacacgcca agagaggaag cagagggagg cggaagcgtg 180
 gaggaagggg cgagagggcat catcaaagga gatgagggga gcgtaggggc cgggaaagag 240
 gcacaaggaa gaaagtatgg gaaggaggaa tggaggttca gggctaggcg gcgggagggc 300

33/57

```

gccaggccgg gaagagtaca aggacaagga ggtcaggttt gggcctacat cccggggaca 360
ggggcggcca tggcggcgcc agccaggag gaggaggagg aggcggctcg qgagtcagcc 420
gcctgcccg ctgcggggcc agcgcctctg cgcctgccgg aagtgtctgt gctgcacatg 480
tgctcctacc tcgacatgcg ggccctcggc cgcctggccc aggtgtaccg ctggctgtgg 540
cacttcacca actgcgacct gctccggcgc cagatagcct gggcctcgct caactccggc 600
ttcacggcgc tcggcaccaa cctgatgacc agtgtccag tgaagggtgc tcagaactgg 660
atagtggggt gctgcgagag ggggattctg ctgaagtggg gatgcagtca gatgccctgg 720
atgcagctag aggatgatgc tttgtacata tcccaggcta atttcatcct ggccctaccag 780
ttccgtccag atggtgcccag cttgaaccgt cagcctctgg gagtctctgc tgggcatgat 840
gaggacgttt gccactttgt gctggccacc tcgcataatg tcagtgcagg aggagatggg 900
aagattggcc ttgtaagat tcacagcacc ttcgctgcca agtactgggc tcatgaacag 960
gaggtgaact gtgtggattg caaagggggc atcatatcat ttggctccag ggacaggagc 1020
gccaaagtgt ggcccttggc ctcaggccag ctggggcagt gtttatacac catccagact 1080
gaagacccaa tctggctctgt tgctatcagg ccattactca gctcttttgt gacagggacg 1140
gcttgtttgt ggcactttct acccctgaaa atctgggacc tcaacagtgg gcagctgatg 1200
acacacttgg acagagaact tcccccaagg gctggggtgc tggatgtcat atatgagtc 1260
cctttcgcac tgctctcctg tgctatgac acctatgttc gctactggga ctgccgcacc 1320
agtgtccgga aatgtgtcat ggagtgggag gagcccccaca acagcaacct gtactgctg 1380
cagacagatg gcaaccaact gcttggccaca ggctcctcct tctatagcgt tgtacggctg 1440
tgggaccggc accaaagggc ctgccgcac accttccgc tgacgtcgac ccgcctcggc 1500
agccctgtgt actgcctgca tctcaccacc aagcatctct atgctgcgct gtcttacaac 1560
ctccacgtcc tggatattca aaaccctga 1590

```

<210> 44
 <211> 529
 <212> PRT
 <213> Homo sapiens

<400> 44
 Arg Gly Gly Ser Glu Gly Arg Gly Arg Gly Arg Glu Lys Arg Ala Arg
 1 5 10 15
 Gly Ala Arg Arg Lys Arg Lys Gln Gly Gly Arg Glu Ala Arg Ala Ala
 20 25 30
 Asp Gly Glu Gly Gly Ser Gly Pro Gly Ala Glu Ala Gly Ala Arg Thr
 35 40 45
 Arg Pro Arg Glu Glu Ala Glu Gly Gly Gly Ser Val Glu Glu Gly Ala
 50 55 60
 Arg Gly Ile Ile Lys Gly Asp Glu Gly Ser Val Gly Ala Gly Lys Glu
 65 70 75 80
 Ala Gln Gly Arg Lys Tyr Gly Lys Glu Glu Trp Arg Val Arg Ala Arg
 85 90 95
 Arg Arg Glu Gly Ala Arg Pro Gly Arg Val Gln Gly Gln Gly Gln
 100 105 110
 Val Trp Ala Tyr Ile Pro Gly Thr Gly Ala Ala Met Ala Ala Ala Ala
 115 120 125
 Arg Glu Glu Glu Glu Glu Ala Ala Arg Glu Ser Ala Ala Cys Pro Ala
 130 135 140
 Ala Gly Pro Ala Leu Trp Arg Leu Pro Glu Val Leu Leu Leu His Met
 145 150 155 160
 Cys Ser Tyr Leu Asp Met Arg Ala Leu Gly Arg Leu Ala Gln Val Tyr
 165 170 175
 Arg Trp Leu Trp His Phe Thr Asn Cys Asp Leu Leu Arg Arg Gln Ile

180										185										190													
Ala	Trp	Ala	Ser	Leu	Asn	Ser	Gly	Phe	Thr	Arg	Leu	Gly	Thr	Asn	Leu																		
		195					200					205																					
Met	Thr	Ser	Val	Pro	Val	Lys	Val	Ser	Gln	Asn	Trp	Ile	Val	Gly	Cys																		
		210					215					220																					
Cys	Arg	Glu	Gly	Ile	Leu	Leu	Lys	Trp	Arg	Cys	Ser	Gln	Met	Pro	Trp																		
		225				230					235				240																		
Met	Gln	Leu	Glu	Asp	Asp	Ala	Leu	Tyr	Ile	Ser	Gln	Ala	Asn	Phe	Ile																		
				245					250					255																			
Leu	Ala	Tyr	Gln	Phe	Arg	Pro	Asp	Gly	Ala	Ser	Leu	Asn	Arg	Gln	Pro																		
			260					265					270																				
Leu	Gly	Val	Ser	Ala	Gly	His	Asp	Glu	Asp	Val	Cys	His	Phe	Val	Leu																		
		275					280					285																					
Ala	Thr	Ser	His	Ile	Val	Ser	Ala	Gly	Gly	Asp	Gly	Lys	Ile	Gly	Leu																		
		290					295				300																						
Gly	Lys	Ile	His	Ser	Thr	Phe	Ala	Ala	Lys	Tyr	Trp	Ala	His	Glu	Gln																		
		305				310				315				320																			
Glu	Val	Asn	Cys	Val	Asp	Cys	Lys	Gly	Gly	Ile	Ile	Ser	Phe	Gly	Ser																		
			325					330						335																			
Arg	Asp	Arg	Thr	Ala	Lys	Val	Trp	Pro	Leu	Ala	Ser	Gly	Gln	Leu	Gly																		
			340					345					350																				
Gln	Cys	Leu	Tyr	Thr	Ile	Gln	Thr	Glu	Asp	Gln	Ile	Trp	Ser	Val	Ala																		
		355					360					365																					
Ile	Arg	Pro	Leu	Leu	Ser	Ser	Phe	Val	Thr	Gly	Thr	Ala	Cys	Cys	Gly																		
		370					375				380																						
His	Phe	Ser	Pro	Leu	Lys	Ile	Trp	Asp	Leu	Asn	Ser	Gly	Gln	Leu	Met																		
		385				390				395				400																			
Thr	His	Leu	Asp	Arg	Asp	Phe	Pro	Pro	Arg	Ala	Gly	Val	Leu	Asp	Val																		
			405					410					415																				
Ile	Tyr	Glu	Ser	Pro	Phe	Ala	Leu	Leu	Ser	Cys	Gly	Tyr	Asp	Thr	Tyr																		
		420					425					430																					
Val	Arg	Tyr	Trp	Asp	Cys	Arg	Thr	Ser	Val	Arg	Lys	Cys	Val	Met	Glu																		
		435					440					445																					
Trp	Glu	Glu	Pro	His	Asn	Ser	Thr	Leu	Tyr	Cys	Leu	Gln	Thr	Asp	Gly																		
		450				455					460																						
Asn	His	Leu	Leu	Ala	Thr	Gly	Ser	Ser	Phe	Tyr	Ser	Val	Val	Arg	Leu																		
		465				470				475				480																			
Trp	Asp	Arg	His	Gln	Arg	Ala	Cys	Pro	His	Thr	Phe	Pro	Leu	Thr	Ser																		
			485					490					495																				
Thr	Arg	Leu	Gly	Ser	Pro	Val	Tyr	Cys	Leu	His	Leu	Thr	Thr	Lys	His																		
		500						505					510																				
Leu	Tyr	Ala	Ala	Leu	Ser	Tyr	Asn	Leu	His	Val	Leu	Asp	Ile	Gln	Asn																		

515

520

525

Pro

<210> 45

<211> 1214

<212> DNA

<213> Homo sapiens

<400> 45

```

gcattgctat aattttacta tactctcacc taaatctaaa atcagctcttc aaaataaaaa 60
caaatgtgcc ttggccaaaa atttttttta tcgcacaatt aattgacatt aactgccaat 120
tctttttggc taattgacta attttaactt ctgtgttgct ttccagagg catggctatt 180
gcacottggg agaagccttt aatcggttag acttctcaag tgcaattcaa gatataccga 240
cgttcaatta tgtggtcaaa ctgttgccgc taattgcaaa atccagatta acttcattga 300
gtggcgtggc acagaagaat tacttcaaca ttttgataa aatcgttcaa aaggttcttg 360
atgaccacca caatcctcgc ttaatcaaa atcttctgca agacctaaag tctacctctt 420
gcattcttat tagaggagta ggaagctctg tattagtggg aaacatcaat atttggattt 480
gccgattaga aactattctc gcctggcaac aacagctaca ggaatctcag atgactaagc 540
aagtgaacaa tggcctcacc ctgagtgcac ttcctctgca catgctgaac aacatcctat 600
accggttctc agacggatgg gacatcatca cctlaggcca ggtgaccccc acgttgatata 660
tgcttagtga agacagacag ctgtggaaga agctttgtca gtaccatttt gctgaaaagc 720
agtattgtag acatttgatc ctttcagaaa aaggtcatat tgaatggaag ttgatgtact 780
ttgcacttca gaaacattac ccagcgaagg agcagtacgg agacacactg catttctgtc 840
ggcactgcag cattctcttt lggaaaggact caggacaccc ctgcacggcg gccgacctg 900
acagctgctt cagcctgtg tctccgcagc acttcacoga cctcttcaag ttttaaggcg 960
tgccctgccc atccctattg gagattgtga atcctgctgt ctgtgcaggg ctcatagtga 1020
gtgttctgtg aggtgggtgg agactcctcg gaagccctcg cttccagaaa gccgtgggaag 1080
aactgccctt ctgcaaaagg gggactgcac ggttgcatth tcatcactga aagtcagagg 1140
ccaaggaaat catttctact tctttaaaaa ctcttcttaa gcatattaaa atgtgaaatt 1200
ttgcgtactc tctc                                     1214

```

<210> 46

<211> 272

<212> PRT

<213> Homo sapiens

<400> 46

```

Leu Ile Leu Thr Ser Val Leu Leu Phe Gln Arg His Gly Tyr Cys Thr
 1             5             10             15

Leu Gly Glu Ala Phe Asn Arg Leu Phe Ser Ser Ala Ile Gln Asp
      20             25             30

Ile Arg Thr Phe Asn Tyr Val Val Lys Leu Leu Gln Leu Ile Ala Lys
 35             40             45

Ser Gln Leu Thr Ser Leu Ser Gly Val Ala Gln Lys Asn Tyr Phe Asn
 50             55             60

Ile Leu Asp Lys Ile Val Gln Lys Val Leu Asp Asp His His Asn Pro
 65             70             75             80

Arg Leu Ile Lys Asp Leu Leu Gln Asp Leu Ser Ser Thr Leu Cys Ile
      85             90             95

Leu Ile Arg Gly Val Gly Lys Ser Val Leu Val Gly Asn Ile Asn Ile
 100             105             110

Trp Ile Cys Arg Leu Glu Thr Ile Leu Ala Trp Gln Gln Gln Leu Gln
 115             120             125

```

36/57

Asp Leu Gln Met Thr Lys Gln Val Asn Asn Gly Leu Thr Leu Ser Asp
 130 135 140
 Leu Pro Leu His Met Leu Asn Asn Ile Leu Tyr Arg Phe Ser Asp Gly
 145 150 155 160
 Trp Asp Ile Ile Thr Leu Gly Gln Val Thr Pro Thr Leu Tyr Met Leu
 165 170 175
 Ser Glu Asp Arg Gln Leu Trp Lys Lys Leu Cys Gln Tyr His Phe Ala
 180 185 190
 Glu Lys Gln Phe Cys Arg His Leu Ile Leu Ser Glu Lys Gly His Ile
 195 200 205
 Glu Trp Lys Leu Met Tyr Phe Ala Leu Gln Lys His Tyr Pro Ala Lys
 210 215 220
 Glu Gln Tyr Gly Asp Thr Leu His Phe Cys Arg His Cys Ser Ile Leu
 225 230 235 240
 Phe Trp Lys Asp Ser Gly His Pro Cys Thr Ala Ala Asp Pro Asp Ser
 245 250 255
 Cys Phe Thr Pro Val Ser Pro Gln His Phe Ile Asp Leu Phe Lys Phe
 260 265 270

<210> 47
 <211> 4059
 <212> DNA
 <213> Homo sapiens

<400> 47
 agtacggcag tgaggggcaaa ggcagctcga gcatctcacc tgacgtgagt tcaagtacag 60
 atcacacgcc cactaaagcc cagaagaatg tggctaccag cgaagactcc gacctgagca 120
 tgcgcacact gagcagccc agcccagccc tgatatgtcc accgaatctc ccaggatttc 180
 agaatggaag gggctcgtcc acctctcgt cctccatcac cggggagacg gtggccatgy 240
 tgcactcccc gcccccgacc cgctcacac acccgctcat cgggctcgcc tccagacccc 300
 agaaggagca ggccagcata gaccggctcc cggaccactc catggtgcag atcttctcct 360
 tctgcccac caaccagctg tgcgctgctg cgcgagtgtg ccgcccgtgg tacaacctgg 420
 cctgggaccc gggctctgag aggactatcc gctgacggg cgagaccatc aacgtggacc 480
 gcgcctcaa ggtgctgacc cgcagactct gccaggacac ccccaacgtg tgtctcatgc 540
 tggaaaacct aactgtcagt ggtgcagcc ggctcacaga ccgagggctg tacaccatcg 600
 cccagtgtg ccccgaaactg aggcgactgg aagtctcagg ctgttacaat atctccaacg 660
 aggcgctctt tgatgtggtg tccctctgcc ctaatctgga gcacctggat gtgtcaggat 720
 gctccaaaag gacctgcac agcttgaccc gggaggcctc cattaaactg tcaaccttgc 780
 atggcaaaaca gatttccatc cgctacctgg acatgacgga ctgcttctgt ctggaggacg 840
 aaggcctgca caccatcgcg gcgcactgca cgcagctcac ccacctctac ctgcgccgtg 900
 gcgtccgct gaccgacgaa ggccctgcgt acctggtgat ctactgcgcc tccatcaagg 960
 agctgagcgt cagcgactgc cgcttctgca gcgacttogg cctgcccggag atcgccaagc 1020
 ttgagtcocg cctgcggtag ctgagcatcg cgcactgcgg ccgggtcacc gacgtgggca 1080
 tccgctacgt ggccaagtac tgcagcaagc tgcgtacct caacgcgagg ggctgcgagg 1140
 gcatcacgga ccacggtgtg gagtacctcg ccaagaactg caccaaactc aaatccctgg 1200
 atatcggaat atgocctttg gtatccgaca cgggcctgga gtgcctggcc ctgaactgct 1260
 tcaacctcaa gcggtcagc ctcaagtctt gcgagagcat caccggccag ggcttgca 1320
 togtggccgc caactgcttt gacctcaga cgtgaaatgt ccaggactgc gaggtctccg 1380
 tggaggccct gcgctttgtc aaacgcact gcaagcgtg cgtcatcgag cacaccaacc 1440
 cggctttctt ctgaagggac agagttcatc cggcgttgta ttcacacaaa cctgaacaaa 1500
 gcaaatTTTT ttaaaagcag cgtatgtaag caccgacacc cactcaaaac agctctttct 1560

```

tccgggaagg ttattaggaa tctggccttt atttttcctc atttctcatg ggcaacagag 1620
gccaaagaaa cgaagcaaga caaacagcaa acaggcattt tggtcaggctc atttctagcc 1680
agtttctctt ctcacaaaag atgtacttaa gcaggctgat cgctgttctt tgagcaaggc 1740
gcttactctc ctccgctcag gccccaaagg ccgccccttc cctcgacacac aggcccccacc 1800
cccacagttc cacgcccccc ccccaaggcc acaccctccc tccctagagc agcagcgagg 1860
atccatcctc agaatacacag tgcctctccag aactcctctc taaactgctt cattgacctt 1920
agtcactctc ttcaatccca caccatgga cattcttgtc aactcaatac catagcactt 1980
tgcataggca aaatactttt caggcctttt taaaaaatte attacagcaa acagctgggg 2040
aaggacatgc agtctctccc cagctctgtc aatgactatg acctgggcca aagcacttca 2100
ctgctctggg ctgcagcttc cagcactgaa tcagaggcca cacagcccaa agattagctt 2160
catgtccatt atagcattga gggagcagag ataccatac acagaagcac cltggcatag 2220
agcaccagc catcgacctc ttccaggaga actgattctg tggatggatg tgatttcagg 2280
agatttgtga gtgccagcat cagtgcataa aggtctctgt atgtcctttg gctgcaaatc 2340
acccacttcc ctgtgtttca gtgggagaat ttctctctcc acctctcac atctctttt 2400
gccaggctgg atgtctgtgt ctctgtacac aaatacttct tgcattcccc cctccacacc 2460
atctatcag ggcaccagca cactaatca cagcaaagcc cagatcccc catcagttgc 2520
ttttactcag ttttttcaaa taggagtaaa ggcctctgca atttttaatt aacaagcaag 2580
gccccaggga acacatgtcc tcaaaaagtt ttctgatccc tgccttgca cacctggcat 2640
gcacagga catctgtctt acagctggca gagacagatg cctcggttct ttgtcattca 2700
gattgcattt gacctcttct catctattta ttctttata catccagact tcatcacatg 2760
aagcctattg gggtaagtt tgtaagtgtt taattgtgca aattgccacc ctgtgtact 2820
cctccatgtc tgtctgctgt ttttccacca aagaatgcaa agcagacttc cagggtgtta 2880
aattctgttc actcaacaat gccagatgaa tgggaagagg aacacactga gatgacttag 2940
actctgttcc accaaccaga cccttggaat ggaatactaa aatcattaca aggtatggat 3000
tttaaatgga tgaaacttca aattatctta ttggataga agtctatatt ctagcctcat 3060
ttgcataag tcagatagcc agaagaaatt ccattgctgg ttttcaagaa attcacttgt 3120
cttttgctaa taaacacatg gccctttccc agattatct ctagccaagc cccaccttg 3180
ttacgttgaa atccctcatt tattttcttc tcaaaatgcc cattatccaa atgcagaacc 3240
tctgcacttc caagccagtt atgtgaatt tgtcaactt agacaccctt gacaactgca 3300
ctcctactgt aggtcctgt gcatactgtc gtcttctgtg ggggatggag aggttagtgt 3360
gatgaggtgg tgtctgccc gagggttct ttcaaacatc atggcctccc atccaatcaa 3420
catcatcaaa ttacatgtgt aatcaaggct ctgtgccatg ggggaaatga atcatttagc 3480
taaggccagga tctagtgaat gccacagagt ttaaaaccat gaaagaagtt gaaggcagca 3540
ttcctcagct ctgtgacttg tgacctatt tgaagtttca ggatttgggt gtcacaaaagg 3600
attgtcccta atccttgccc ctggggtctt ccgagtgaag tggtttaata ctctgagaat 3660
gagcagggag atccagagaa tgaatccctg accgcatcac ctaaaactgtc ttccaaacat 3720
gagacaaaag tgactgttca cactgattgc ccagcacata ccgtcttgcc agtttcttct 3780
tttctcccag tctctgttcc atccattctg ttctcccttg ggttggaat ctatgatgga 3840
ggttactggg gaaacagctc agcagatttt tggagacca accaaagggtc tcaactaggaa 3900
atctatctgt ttttaaacat tgcttcttcc ctggctctgc taaattgaat gctcattgtt 3960
tggtgtgtgt gttttttaat tctaattgtt aaatcactgc gtgctgtatg aatctagaaa 4020
gccttaattt actaccaaga aataaagcaa tatgttctg 4059

```

<210> 48
 <211> 483
 <212> PRT
 <213> Homo sapiens

<400> 48
 Tyr Gly Ser Glu Gly Lys Gly Ser Ser Ser Ile Ser Ser Asp Val Ser
 1 5 10 15
 Ser Ser Thr Asp His Thr Pro Thr Lys Ala Gln Lys Asn Val Ala Thr
 20 25 30
 Ser Glu Asp Ser Asp Leu Ser Met Arg Thr Leu Ser Thr Pro Ser Pro
 35 40 45
 Ala Leu Ile Cys Pro Pro Asn Leu Pro Gly Phe Gln Asn Gly Arg Gly
 50 55 60
 Ser Ser Thr Ser Ser Ser Ser Ile Thr Gly Glu Thr Val Ala Met Val
 65 70 75 80

His Ser Pro Pro Pro Thr Arg Leu Thr His Pro Leu Ile Arg Leu Ala
 85 90 95
 Ser Arg Pro Gln Lys Glu Gln Ala Ser Ile Asp Arg Leu Pro Asp His
 100 105 110
 Ser Met Val Gln Ile Phe Ser Phe Leu Pro Thr Asn Gln Leu Cys Arg
 115 120 125
 Cys Ala Arg Val Cys Arg Arg Trp Tyr Asn Leu Ala Trp Asp Pro Arg
 130 135 140
 Leu Trp Arg Thr Ile Arg Leu Thr Gly Glu Thr Ile Asn Val Asp Arg
 145 150 155 160
 Ala Leu Lys Val Leu Thr Arg Arg Leu Cys Gln Asp Thr Pro Asn Val
 165 170 175
 Cys Leu Met Leu Glu Thr Val Thr Val Ser Gly Cys Arg Arg Leu Thr
 180 185 190
 Asp Arg Gly Leu Tyr Thr Ile Ala Gln Cys Cys Pro Glu Leu Arg Arg
 195 200 205
 Leu Glu Val Ser Gly Cys Tyr Asn Ile Ser Asn Glu Ala Val Phe Asp
 210 215 220
 Val Val Ser Leu Cys Pro Asn Leu Glu His Leu Asp Val Ser Gly Cys
 225 230 235 240
 Ser Lys Val Thr Cys Ile Ser Leu Thr Arg Glu Ala Ser Ile Lys Leu
 245 250 255
 Ser Pro Leu His Gly Lys Gln Ile Ser Ile Arg Tyr Leu Asp Met Thr
 260 265 270
 Asp Cys Phe Val Leu Glu Asp Glu Gly Leu His Thr Ile Ala Ala His
 275 280 285
 Cys Thr Gln Leu Thr His Leu Tyr Leu Arg Arg Cys Val Arg Leu Thr
 290 295 300
 Asp Glu Gly Leu Arg Tyr Leu Val Ile Tyr Cys Ala Ser Ile Lys Glu
 305 310 315 320
 Leu Ser Val Ser Asp Cys Arg Phe Val Ser Asp Phe Gly Leu Arg Glu
 325 330 335
 Ile Ala Lys Leu Glu Ser Arg Leu Arg Tyr Leu Ser Ile Ala His Cys
 340 345 350
 Gly Arg Val Thr Asp Val Gly Ile Arg Tyr Val Ala Lys Tyr Cys Ser
 355 360 365
 Lys Leu Arg Tyr Leu Asn Ala Arg Gly Cys Glu Gly Ile Thr Asp His
 370 375 380
 Gly Val Glu Tyr Leu Ala Lys Asn Cys Thr Lys Leu Lys Ser Leu Asp
 385 390 395 400
 Ile Gly Lys Cys Pro Leu Val Ser Asp Thr Gly Leu Glu Cys Leu Ala
 405 410 415

39/57

Leu Asn Cys Phe Asn Leu Lys Arg Leu Ser Leu Lys Ser Cys Glu Ser
 420 425 430

Ile Thr Gly Gln Gly Leu Gln Ile Val Ala Ala Asn Cys Phe Asp Leu
 435 440 445

Gln Thr Leu Asn Val Gln Asp Cys Glu Val Ser Val Glu Ala Leu Arg
 450 455 460

Phe Val Lys Arg His Cys Lys Arg Cys Val Ile Glu His Thr Asn Pro
 465 470 475 480

Ala Phe Phe

<210> 49
 <211> 850
 <212> DNA
 <213> Homo sapiens

<400> 49
 tgccggccgcg ccgcgaccgc caccggcacc caccgccacg ccgaggaag ggcccgacgc 60
 gggctgggga gaccgcattc ccttggaat cctgggcag attttcgggt tgttggtggc 120
 ggccgaacgc cccatgccct tcctgggcag ggctgcgcgc gtgtgccgc gctggcagga 180
 ggccgcttcc caaccgcgc totggcacac cgtgaccctg tcgtccccc tggtcggcgc 240
 gcctgccaag ggccgggtca agccggagaa gaagctcctt gcttccttg agtggcttat 300
 gcccattcgg ttttcacagc tccagaggct gacctcacc cactggaagt ctcagggtaca 360
 ccccggttg aagctggtag gtgagtgctg tcctcggctc actttcctca agctctccgc 420
 ctgccacggt gtgactgctg acgctctggt catgctaacc aaagcctgct gccagctcca 480
 tagcctggac ctacagcact ccattggtga gtccacagct gtgggtgagct tcttgagga 540
 ggccagggtcc cgaatgcgca agttgtggct gacctacagc tcccagacga cagccatcct 600
 ggccgcattg ctgggcagct gctgccccca gctccaggtc ctggagggtga gcaccggcat 660
 caaccgtaat agcattcccc ttcagctgcc tgcagaggct ctgcagaaag gctgccctca 720
 gctccagggt ctgcggctgt tgaacctgat gtggtgccc aagcctccg gacgagggt 780
 ggctcccgga ccaggcttcc ctagcctaga ggagctctgc ctggcgagct caacctgcaa 840
 ctttctgagc 850

<210> 50
 <211> 283
 <212> PRT
 <213> Homo sapiens

<400> 50
 Ala Ala Ala Pro Ala Pro Ala Pro Ala Pro Thr Pro Thr Pro Glu Glu
 1 5 10 15

Gly Pro Asp Ala Gly Trp Gly Asp Arg Ile Pro Leu Glu Ile Leu Val
 20 25 30

Gln Ile Phe Gly Leu Leu Val Ala Ala Asp Gly Pro Met Pro Phe Leu
 35 40 45

Gly Arg Ala Ala Arg Val Cys Arg Arg Trp Gln Glu Ala Ala Ser Gln
 50 55 60

Pro Ala Leu Trp His Thr Val Thr Leu Ser Ser Pro Leu Val Gly Arg
 65 70 75 80

Pro Ala Lys Gly Gly Val Lys Ala Glu Lys Lys Leu Leu Ala Ser Leu
 85 90 95

Glu Trp Leu Met Pro Asn Arg Phe Ser Gln Leu Gln Arg Leu Thr Leu

40/57

100	105	110
Ile His Trp Lys Ser Gln Val His	Pro Val Leu Lys Leu Val Gly Glu	
115	120	125
Cys Cys Pro Arg Leu Thr Phe Leu Lys Leu Ser Gly Cys His Gly Val		
130	135	140
Thr Ala Asp Ala Leu Val Met Leu Ala Lys Ala Cys Cys Gln Leu His		
145	150	155
Ser Leu Asp Leu Gln His Ser Met Val Glu Ser Thr Ala Val Val Ser		
165	170	175
Phe Leu Glu Glu Ala Gly Ser Arg Met Arg Lys Leu Trp Leu Thr Tyr		
180	185	190
Ser Ser Gln Thr Thr Ala Ile Leu Gly Ala Leu Leu Gly Ser Cys Cys		
195	200	205
Pro Gln Leu Gln Val Leu Glu Val Ser Thr Gly Ile Asn Arg Asn Ser		
210	215	220
Ile Pro Leu Gln Leu Pro Val Glu Ala Leu Gln Lys Gly Cys Pro Gln		
225	230	235
Leu Gln Val Leu Arg Leu Leu Asn Leu Met Trp Leu Pro Lys Pro Pro		
245	250	255
Gly Arg Gly Val Ala Pro Gly Pro Gly Phe Pro Ser Leu Glu Glu Leu		
260	265	270
Cys Leu Ala Ser Ser Thr Cys Asn Phe Val Ser		
275	280	

<210> 51
 <211> 1777
 <212> DNA
 <213> Homo sapiens

<220>
 <221> modified_base
 <222> all n positions
 <223> n=a, c, g or t

<400> 51
 acaacactgc tctcagaagg atactgcaga actccttaga ggtcttagcc tatggaatca 60
 tgctgaagag cgacagaart tttttaataa ttccgtggat gaaaagtcag ataaagaagc 120
 agaagtgtca gaacactcca caggtataac ccattcttcc cctgaggtaa tgctgtcaat 180
 tttcagctat cttaatctctc aagagttatg tcgatgcagt caagtaagca tgaaatgggc 240
 tcagctgaca aaaacgggat cgcttttgaa acatctttac cctgttcatt gggccagagg 300
 tgactggtat agtgggtccc caactgaact tgatactgaa cctgatgatg aatgggtgaa 360
 aaataggaaa gatgaaagtc gtgcttttca tgagtgggat gaagatgctg acattgatga 420
 atctgaaagag tctgcgagg aatcaattgc tatcagcatt gcacaaatgg aaaaacggtt 480
 actccatggc ttaattcata acgtttctac atatgttggt acttctgtaa aaacottagt 540
 attagcatac agctctgcag ttccagcaa aatgggttagg cagattttag agctttgtcc 600
 taacctggag catctggatc ttaccagac tgacatttca gattctgcat ttgacagttg 660
 gtcttggtgt ggttgctgac agagtcttgc gcattctgat ctgtctgggt gtgagaaaat 720
 cacagatgtg gccctagaga agatttccag agctotttga attctgacat ctcattcaaa 780
 tggctttttg aaaacatcta caagcaaaat tacttcaact gcgtggaaaa ataaagacat 840
 taccatgcag tccaccaagc agtatgcctg ttgcacgat ttaactaaca agggcattgg 900
 agaagaaata gataatgaac acccctggac taagcctgtt tcttctgaga atttcaactc 960

41/57

```

tccttatgtg tggatgttag atgctgaaga tttggctgat attgaagata ctgtggaatg 1020
gagacataga aatgttgaaa gtctttgtgt aatggaaaca gcatccaact ttagttgttc 1080
cacctctggg tgttttagta aggacattgt tggactaagg actagtgtct gttggcagca 1140
gcattgtgct tctccagcct ttgcgtattg tggcactca tttgtttgta caggaacago 1200
tttaagaact atgtcatcac toccagaatc ttctgcaatg tgtagaaaag cagcaaggac 1260
tagattgcct aggggaaaag acttaattta ctttgggagt gaaaaatctg atcaagagac 1320
tggacgtgta cttctgtttc tcagtttato tggatgttat cagatcacag accatggtct 1380
cagggttttg actctgggag gagggtgccc ttatttggag caccttaatc tctctggttg 1440
tcttactata actggtgcag gctgcagga tttggtttca gcatgtcctt ctctgaatga 1500
tgaatacttt tactactgtg acaacattaa cggctcctcat gctgataccg ccagtggatg 1560
ccagaatttg cagtgtggtt ttcgagcctg ctgccgtctt ggogaatgac ccttgacttc 1620
tgatctttgt ctacttcatt tagctgagca ggctttcttt catgcacttt actcatagca 1680
catttcttgt gtttaaccatc cctttttgag cgtgacttgt tttgggccc a ttnyttacaa 1740
cttcagaaat ctttaattacc agtgrattgt aatgttg 1777

```

<210> 52

<211> 590

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> all Xaa positions

<223> Xaa=unknown amino acid residue

<400> 52

```

Gln His Cys Ser Gln Lys Asp Thr Ala Glu Leu Leu Arg Gly Leu Ser
  1             5             10             15

```

```

Leu Trp Asn His Ala Glu Glu Arg Gln Lys Phe Phe Lys Tyr Ser Val
      20             25             30

```

```

Asp Glu Lys Ser Asp Lys Glu Ala Glu Val Ser Glu His Ser Thr Gly
      35             40             45

```

```

Ile Thr His Leu Pro Pro Glu Val Met Leu Ser Ile Phe Ser Tyr Leu
      50             55             60

```

```

Asn Pro Gln Glu Leu Cys Arg Cys Ser Gln Val Ser Met Lys Trp Ser
      65             70             75             80

```

```

Gln Leu Thr Lys Thr Gly Ser Leu Trp Lys His Leu Tyr Pro Val His
      85             90             95

```

```

Trp Ala Arg Gly Asp Trp Tyr Ser Gly Pro Ala Thr Glu Leu Asp Thr
      100            105            110

```

```

Glu Pro Asp Asp Glu Trp Val Lys Asn Arg Lys Asp Glu Ser Arg Ala
      115            120            125

```

```

Phe His Glu Trp Asp Glu Asp Ala Asp Ile Asp Glu Ser Glu Glu Ser
      130            135            140

```

```

Ala Glu Glu Ser Ile Ala Ile Ser Ile Ala Gln Met Glu Lys Arg Leu
      145            150            155            160

```

```

Leu His Gly Leu Ile His Asn Val Leu Pro Tyr Val Gly Thr Ser Val
      165            170            175

```

```

Lys Thr Leu Val Leu Ala Tyr Ser Ser Ala Val Ser Ser Lys Met Val
      180            185            190

```

```

Arg Gln Ile Leu Glu Leu Cys Pro Asn Leu Glu His Leu Asp Leu Thr

```

195	200	205
Gln Thr Asp Ile Ser Asp Ser Ala Phe Asp Ser Trp Ser Trp Leu Gly		
210	215	220
Cys Cys Gln Ser Leu Arg His Leu Asp Leu Ser Gly Cys Glu Lys Ile		
225	230	235 240
Thr Asp Val Ala Leu Glu Lys Ile Ser Arg Ala Leu Gly Ile Leu Thr		
	245 250	255
Ser His Gln Ser Gly Phe Leu Lys Thr Ser Thr Ser Lys Ile Thr Ser		
	260 265	270
Thr Ala Trp Lys Asn Lys Asp Ile Thr Met Gln Ser Thr Lys Gln Tyr		
	275 280	285
Ala Cys Leu His Asp Leu Thr Asn Lys Gly Ile Gly Glu Glu Ile Asp		
	290 295	300
Asn Glu His Pro Trp Thr Lys Pro Val Ser Ser Glu Asn Phe Thr Ser		
305	310 315	320
Pro Tyr Val Trp Met Leu Asp Ala Glu Asp Leu Ala Asp Ile Glu Asp		
	325 330	335
Thr Val Glu Trp Arg His Arg Asn Val Glu Ser Leu Cys Val Met Glu		
	340 345	350
Thr Ala Ser Asn Phe Ser Cys Ser Thr Ser Gly Cys Phe Ser Lys Asp		
	355 360	365
Ile Val Gly Leu Arg Thr Ser Val Cys Trp Gln Gln His Cys Ala Ser		
	370 375	380
Pro Ala Phe Ala Tyr Cys Gly His Ser Phe Cys Cys Thr Gly Thr Ala		
385	390 395	400
Leu Arg Thr Met Ser Ser Leu Pro Glu Ser Ser Ala Met Cys Arg Lys		
	405 410	415
Ala Ala Arg Thr Arg Leu Pro Arg Gly Lys Asp Leu Ile Tyr Phe Gly		
	420 425	430
Ser Glu Lys Ser Asp Gln Glu Thr Gly Arg Val Leu Leu Phe Leu Ser		
	435 440	445
Leu Ser Gly Cys Tyr Gln Ile Thr Asp His Gly Leu Arg Val Leu Thr		
	450 455	460
Leu Gly Gly Gly Leu Pro Tyr Leu Glu His Leu Asn Leu Ser Gly Cys		
465	470 475	480
Leu Thr Ile Thr Gly Ala Gly Leu Gln Asp Leu Val Ser Ala Cys Pro		
	485 490	495
Ser Leu Asn Asp Glu Tyr Phe Tyr Tyr Cys Asp Asn Ile Asn Gly Pro		
	500 505	510
His Ala Asp Thr Ala Ser Gly Cys Gln Asn Leu Gln Cys Gly Phe Arg		
	515 520	525
Ala Cys Cys Arg Ser Gly Glu Pro Leu Thr Ser Asp Leu Cys Leu Leu		

43/57

530	535	540
His Leu Ala Glu Gln Ala Phe Phe His Ala Leu Tyr Ser His Ile Ser		
545	550	555
Cys Val Asn His Pro Phe Leu Ser Val Thr Cys Phe Gly Pro Ile Xaa		
	565	570
		575
Tyr Asn Phe Arg Asn Leu Asn Tyr Gln Xaa Ile Val Met Leu		
	580	585
		590

<210> 53
 <211> 1681
 <212> DNA
 <213> Homo sapiens

<220>
 <221> modified_base
 <222> all n positions
 <223> n=a, c, g or t

<400> 53
 ttttactgta cacagttgat gtattttgat gctgggcctg tctgggtctgt cttgaggatt 60
 attaaccttt agaggtatca gagaagcaaa tgggtactgg tgaggctgct cattagggaa 120
 gagggcaaaa ggagcactag ctaggtcaga gccatgtttc aggtcacaat gtgatgtcag 180
 atgttgctta taaatccttt cttgtcttcg ccattcttaa atcttgatag gtgcctgttg 240
 ggaaaactgta aatgcctttc ccaatggaga atcaacagat tggglgatgg tggagtcggg 300
 caggaagact caggtcttct agaggaaagg atgcctcacc accccttngg occaggcagc 360
 tgctgtcaga gaatgacaca gcacctgcac agtcgctgtc cacttctctc cactgctgtc 420
 ggtgggggtga cgggagcaaa gtaggcgtgg actttgacat gagggagctg agcccgcatc 480
 cgcttgatgc ctgcacgggt aacctgctgg cagtcgtaca gctcgaggcg ctccaggcct 540
 cggcagttct ctagggtgtc cagggccaca tcagtgtatg ggaggcagtt gtccaactcc 600
 agtaccogca gcctctcatg gccacaggta ctgttgctca ggtgcaggat cccatcatct 660
 gkgatgagtt cacagtggga caggctcagg gcttgagtt taggacagtg aatggagagc 720
 tggatgagtg tgctgtcggg tatcaggatg cawtcttcaa gatccatctt ctccaattcg 780
 tggcaatttc gagctaaaag tgtaaaacct gcgtcagtc aatgggagca tcgggcagcc 840
 tccaaaattt gcagtcgagg acagttcaaa cccagggtcg taagagaggg atctgtgagg 900
 ttgctgcaac ccgaaaggca gagagcctgt agccggtgac agccctgca tatctgcaac 960
 acacctcat ccgtgatagc tgagcaggac tgcaagtga ggctcacaag ctcatggcag 1020
 taatttgaa tgtgtttcag agcttcatct tctaactgtg tgcagccctc caggagcagg 1080
 gctttcaggc ctgcacaacc tcgcaccagt gcctcgatgc catccttcgt gatctgatca 1140
 caccaagaga ggttcaggta ctccaggttt cggcagccct cactgatccc cttcaaggag 1200
 ctgtttgtaa tagacacaca ggaggtcaga wccagatgtt tcagcttggg acagaatctg 1260
 ctaaggctat aacacgtgct gtcagtgatt tttgtgcacc cattgaggtt caaatgttca 1320
 atgtttcggc agttctgtgc aaaggtcttc aaggagggaat ccccaacacc aatgcagcct 1380
 cgcaagctga gcttctcag gaatccaacg categcttcg agatattttc caccactoga 1440
 ccctctacat ctatttgaaa gttaaaaaga tctattcttt gccagtgtct tccatccagg 1500
 gctaagatgt tccaagcctt ggaaatctgt gcacatcggc acaaagttac tatatccaag 1560
 aaggaaaata ttcttaacag aagtctcttg ggtaactttt tgttaataag gccttcatca 1620
 ttgtttgaga aaaccatggc cgaagagccg cgagcgagcc cacagcccga agtcacacgg 1680
 c 1681

<210> 54
 <211> 437
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> all Xaa positions
 <223> Xaa=unknown amino acid residue

<400> 54
 Arg Val Thr Ser Gly Cys Gly Leu Ala Arg Gly Ser Ser Ala Met Val
 1 5 10 15
 Phe Ser Asn Asn Asp Glu Gly Leu Ile Asn Lys Lys Leu Pro Lys Glu
 20 25 30
 Leu Leu Leu Arg Ile Phe Ser Phe Leu Asp Ile Val Thr Leu Cys Arg
 35 40 45
 Cys Ala Gln Ile Ser Lys Ala Trp Asn Ile Leu Ala Leu Asp Gly Ser
 50 55 60
 Asn Trp Gln Arg Ile Asp Leu Phe Asn Phe Gln Ile Asp Val Glu Gly
 65 70 75 80
 Arg Val Val Glu Asn Ile Ser Lys Arg Cys Val Gly Phe Leu Arg Lys
 85 90 95
 Leu Ser Leu Arg Gly Cys Ile Gly Val Gly Asp Ser Ser Leu Lys Thr
 100 105 110
 Phe Ala Gln Asn Cys Arg Asn Ile Glu His Leu Asn Leu Asn Gly Cys
 115 120 125
 Thr Lys Ile Thr Asp Ser Thr Cys Tyr Ser Leu Ser Arg Phe Cys Ser
 130 135 140
 Lys Leu Lys His Leu Xaa Leu Thr Ser Cys Val Ser Ile Thr Asn Ser
 145 150 155 160
 Ser Leu Lys Gly Ile Ser Glu Gly Cys Arg Asn Leu Glu Tyr Leu Asn
 165 170 175
 Leu Ser Trp Cys Asp Gln Ile Thr Lys Asp Gly Ile Glu Ala Leu Val
 180 185 190
 Arg Gly Cys Arg Gly Leu Lys Ala Leu Leu Leu Arg Gly Cys Thr Gln
 195 200 205
 Leu Glu Asp Glu Ala Leu Lys His Ile Gln Asn Tyr Cys His Glu Leu
 210 215 220
 Val Ser Leu Asn Leu Gln Ser Cys Ser Arg Ile Thr Asp Glu Gly Val
 225 230 235 240
 Val Gln Ile Cys Arg Gly Cys His Arg Leu Gln Ala Leu Cys Leu Ser
 245 250 255
 Gly Cys Ser Asn Leu Thr Asp Ala Ser Leu Thr Ala Leu Gly Leu Asn
 260 265 270
 Cys Pro Arg Leu Gln Ile Leu Glu Ala Ala Arg Cys Ser His Leu Thr
 275 280 285
 Asp Ala Gly Phe Thr Leu Leu Ala Arg Asn Cys His Glu Leu Glu Lys
 290 295 300
 Met Asp Leu Glu Xaa Cys Ile Leu Ile Thr Asp Ser Thr Leu Ile Gln
 305 310 315 320
 Leu Ser Ile His Cys Pro Lys Leu Gln Ala Leu Ser Leu Ser His Cys
 325 330 335

45/57

Glu Leu Ile Xaa Asp Asp Gly Ile Leu His Leu Ser Asn Ser Thr Cys
 340 345 350

Gly His Glu Arg Leu Arg Val Leu Glu Leu Asp Asn Cys Leu Leu Ile
 355 360 365

Thr Asp Val Ala Leu Xaa His Leu Glu Asn Cys Arg Gly Leu Glu Arg -
 370 375 380

Leu Glu Leu Tyr Asp Cys Gln Gln Val Thr Arg Ala Gly Ile Lys Arg
 385 390 395 400

Met Arg Ala Gln Leu Pro His Val Lys Val His Ala Tyr Phe Ala Pro
 405 410 415

Val Thr Pro Pro Thr Ala Val Ala Gly Ser Gly Gln Arg Leu Cys Arg
 420 425 430

Cys Cys Val Ile Leu
 435

<210> 55
 <211> 1866
 <212> DNA
 <213> Homo sapiens

<400> 55
 atgtcaccgg tctttcccat gttacacggt ctgaccatgt tttattatat atgccttcgg 60
 cgccgagcca ggacagctac aagaggagaa atgatgaaca cccatagagc tatagaatca 120
 aacagccaga cttccctctt caatgcagag gtagtcacgt atgccaaaga agtagtggat 180
 ttcagttccc attatggaag tgagaatagt atgtccctata ctatgtggaa ttggcctgg 240
 gtaccaaatag tattcccaag ttctgggtgac ttactcaga cagctgtgtt tcgaacttat 300
 gggacatggt gggatcagtg tcttagtget tcttgccat tcaaggaggac gccacctaat 360
 ttccagagcc aggactagt ggaacttact ttgaacaac aggtgtatcc tacagctgta 420
 catgttctag aaacctatca tcccgagaca gtcattagaa ttctcgcttg ttctgcaaat 480
 ccttattccc caaatccacc agctgaagta agatgggaga ttctttggtc agagagacct 540
 acgaaggtga atgttccca agctgcacag tttaaacctt gtattaagca gataaatttc 600
 cccacaaatc ttatacgact ggaagtaa atgttctctt tggaaatatta cactgaatta 660
 gatgcagttg tgctacatgg tgtgaaggac aagccagtc tttctctcaa gacttcactt 720
 attgacatga atgatataga agatgatgcc tatgcagaaa aggatgggtg tggaaatggc 780
 agtcttaaca aaaagttag cagtgtctgc ctgggggaag ggccaaataa tgggtatttt 840
 galaaactac cttatgagct tttcagctg attctgaatc atcttacct accagacctg 900
 tgtagattag cacagacttg caaactactg agccagcatt gctgtgatcc tctgcaatac 960
 atccacctca atctgcaacc ataactggga aaactagatg acacttctct ggaatttcta 1020
 cagtctcgct gcactcttgt ccagtggctt aatttatctt ggactggcaa tagaggcttc 1080
 atctctgttg caggatttag caggtttctg aaggtttgtg gatccgaatt agtacgcctt 1140
 gaattgtctt gcagccactt tcttaatgaa acttgcttag aagtatttct tgagatgtgt 1200
 ccaaattctac aggccttaaa tctctctctc tgtgataagc taccacctca agctttcaac 1260
 cacattgcca agttatgcag ccttaaacga cttgttctct atcgaaacaa agtagagcaa 1320
 acagcactgc tcagcatttt gaacttctgt tcagagcttc agcacctcag tttaggcagt 1380
 tgtgtcatga ttgaagacta tgatgtgata gctagcatga taggagccaa gtgtaaaaaa 1440
 ctccggaccc tggatctgtg gagatgtaag aatattactg agaattggaat agcagaactg 1500
 gcttctgggt gtccactact ggaggagctt gaccttgggt ggtgcccaac tctgcagagc 1560
 agcaccgggt gcttcaccag actggcacac cagctcccaa acttgcaaaa actctttctt 1620
 acagctaata gatctgtgtg tgacacagac attgatgaat tggcatgtaa ttgtaccagg 1680
 ttacagcagc tggacatatt aggaacaaga atggtaaagtc cggcatcctt aagaaaactc 1740
 ctggaatctt gtaaagatct ttctttactt gatgtgtcct tctgttcgca gattgalaac 1800
 agagctgtgc tagaactgaa tgcaagcttt ccaaaagtgt tcataaaaaa gagctttact 1860
 cagtga 1866

<210> 56
 <211> 621

46/57

<212> PRT

<213> Homo sapiens

<400> 56

```

Met Ser Pro Val Phe Pro Met Leu Thr Val Leu Thr Met Phe Tyr Tyr
  1           5           10           15

Ile Cys Leu Arg Arg Arg Ala Arg Thr Ala Thr Arg Gly Glu Met Met
          20           25           30

Asn Thr His Arg Ala Ile Glu Ser Asn Ser Gln Thr Ser Pro Leu Asn
      35           40           45

Ala Glu Val Val Gln Tyr Ala Lys Glu Val Val Asp Phe Ser Ser His
      50           55           60

Tyr Gly Ser Glu Asn Ser Met Ser Tyr Thr Met Trp Asn Leu Ala Gly
      65           70           75           80

Val Pro Asn Val Phe Pro Ser Ser Gly Asp Phe Thr Gln Thr Ala Val
          85           90           95

Phe Arg Thr Tyr Gly Thr Trp Trp Asp Gln Cys Pro Ser Ala Ser Leu
      100           105           110

Pro Phe Lys Arg Thr Pro Pro Asn Phe Gln Ser Gln Asp Tyr Val Glu
      115           120           125

Leu Thr Phe Glu Gln Gln Val Tyr Pro Thr Ala Val His Val Leu Glu
      130           135           140

Thr Tyr His Pro Gly Ala Val Ile Arg Ile Leu Ala Cys Ser Ala Asn
      145           150           155           160

Pro Tyr Ser Pro Asn Pro Pro Ala Glu Val Arg Trp Glu Ile Leu Trp
          165           170           175

Ser Glu Arg Pro Thr Lys Val Asn Ala Ser Gln Ala Arg Gln Phe Lys
          180           185           190

Pro Cys Ile Lys Gln Ile Asn Phe Pro Thr Asn Leu Ile Arg Leu Glu
          195           200           205

Val Asn Ser Ser Leu Leu Glu Tyr Tyr Thr Glu Leu Asp Ala Val Val
      210           215           220

Leu His Gly Val Lys Asp Lys Pro Val Leu Ser Leu Lys Thr Ser Leu
      225           230           235           240

Ile Asp Met Asn Asp Ile Glu Asp Asp Ala Tyr Ala Glu Lys Asp Gly
          245           250           255

Cys Gly Met Asp Ser Leu Asn Lys Lys Phe Ser Ser Ala Val Leu Gly
          260           265           270

Glu Gly Pro Asn Asn Gly Tyr Phe Asp Lys Leu Pro Tyr Glu Leu Ile
          275           280           285

Gln Leu Ile Leu Asn His Leu Thr Leu Pro Asp Leu Cys Arg Leu Ala
      290           295           300

Gln Thr Cys Lys Leu Leu Ser Gln His Cys Cys Asp Pro Leu Gln Tyr
      305           310           315           320

```

47/57

Ile His Leu Asn Leu Gln Pro Tyr Trp Ala Lys Leu Asp Asp Thr Ser
 325 330 335
 Leu Glu Phe Leu Gln Ser Arg Cys Thr Leu Val Gln Trp Leu Asn Leu
 340 345 350
 Ser Trp Thr Gly Asn Arg Gly Phe Ile Ser Val Ala Gly Phe Ser Arg
 355 360 365
 Phe Leu Lys Val Cys Gly Ser Glu Leu Val Arg Leu Glu Leu Ser Cys
 370 375 380
 Ser His Phe Leu Asn Glu Thr Cys Leu Glu Val Ile Ser Glu Met Cys
 385 390 395 400
 Pro Asn Leu Gln Ala Leu Asn Leu Ser Ser Cys Asp Lys Leu Pro Pro
 405 410 415
 Gln Ala Phe Asn His Ile Ala Lys Leu Cys Ser Leu Lys Arg Leu Val
 420 425 430
 Leu Tyr Arg Thr Lys Val Glu Gln Thr Ala Leu Leu Ser Ile Leu Asn
 435 440 445
 Phe Cys Ser Glu Leu Gln His Leu Ser Leu Gly Ser Cys Val Met Ile
 450 455 460
 Glu Asp Tyr Asp Val Ile Ala Ser Met Ile Gly Ala Lys Cys Lys Lys
 465 470 475 480
 Leu Arg Thr Leu Asp Leu Trp Arg Cys Lys Asn Ile Thr Glu Asn Gly
 485 490 495
 Ile Ala Glu Leu Ala Ser Gly Cys Pro Leu Leu Glu Glu Leu Asp Leu
 500 505 510
 Gly Trp Cys Pro Thr Leu Gln Ser Ser Thr Gly Cys Phe Thr Arg Leu
 515 520 525
 Ala His Gln Leu Pro Asn Leu Gln Lys Leu Phe Leu Thr Ala Asn Arg
 530 535 540
 Ser Val Cys Asp Thr Asp Ile Asp Glu Leu Ala Cys Asn Cys Thr Arg
 545 550 555 560
 Leu Gln Gln Leu Asp Ile Leu Gly Thr Arg Met Val Ser Pro Ala Ser
 565 570 575
 Leu Arg Lys Leu Leu Glu Ser Cys Lys Asp Leu Ser Leu Leu Asp Val
 580 585 590
 Ser Phe Cys Ser Gln Ile Asp Asn Arg Ala Val Leu Glu Leu Asn Ala
 595 600 605
 Ser Phe Pro Lys Val Phe Ile Lys Lys Ser Phe Thr Gln
 610 615 620

<210> 57
 <211> 984
 <212> DNA
 <213> Homo sapiens

<400> 57
 atgcaacttg tacctgatat agagttcaag attacttata cccgggtctcc agatgggtgat 60
 ggcgttggaa acagctacat tgaagataat gatgatgaca gcaaaatggc agatctcttg 120
 tccacttcc agcagcaact cacatttcag gagtctgtgc ttaaaactgtg tcagcctgag 180
 cttgagagca gtcagattca catatcagtg ctgccaatgg aggtcctgat gtacatcttc 240
 cgatgggtgg tgtctagtga cttggacctc agatcattgg agcagttgtc gctgggtgtc 300
 agaggattct acatctgtgc cagagacctc gaaatatggc gtctggcctg cttgaaagt 360
 tggggcagaa gctgtattaa acttggtccg tacacgtcct ggagagagat gtttttagaa 420
 cggcctcgtg ttcggtttga tggcgtgtat atcagtaaaa ccacatatat tcgtcaagg 480
 gaacagtctc ttgatggttt ctatagagcc tggcaccaag tggaaatatta caggtacata 540
 agattcttcc ctgatggcca tgtgatgatg ttgacaacct ctgaagagcc tcagtccatt 600
 gttccacgtt taagaactag gaataccagg actgatgcaa ttctactggg tcactatcgc 660
 ttgtcacaag acacagacaa tcagaccaaa gtatttgctg taataactaa gaaaaagaa 720
 gaaaaaccac ttgactataa atacagatat ttctgtctgt tccotgtaca agaagcagat 780
 caqagtttcc atgtggggct acagctatgt tccagtgggc accagaggtt caacaaactc 840
 atctggatac atcattcttg tcacattact tacaatcaa ctgggtgagac tgcagtcagt 900
 gcttttgaga ttgacaagat gtacaccccc ttgttcttcg ccagagtaag gagctacaca 960
 gctttctcag aaaggcctct gtag 984

<210> 58

<211> 327

<212> PRT

<213> Homo sapiens

<400> 58

Met Gln Leu Val Pro Asp Ile Glu Phe Lys Ile Thr Tyr Thr Arg Ser
 1 5 10 15
 Pro Asp Gly Asp Gly Val Gly Asn Ser Tyr Ile Glu Asp Asn Asp Asp
 20 25 30
 Asp Ser Lys Met Ala Asp Leu Leu Ser Tyr Phe Gln Gln Gln Leu Thr
 35 40 45
 Phe Gln Glu Ser Val Leu Lys Leu Cys Gln Pro Glu Leu Glu Ser Ser
 50 55 60
 Gln Ile His Ile Ser Val Leu Pro Met Glu Val Leu Met Tyr Ile Phe
 65 70 75 80
 Arg Trp Val Val Ser Ser Asp Leu Asp Leu Arg Ser Leu Glu Gln Leu
 85 90 95
 Ser Leu Val Cys Arg Gly Phe Tyr Ile Cys Ala Arg Asp Pro Glu Ile
 100 105 110
 Trp Arg Leu Ala Cys Leu Lys Val Trp Gly Arg Ser Cys Ile Lys Leu
 115 120 125
 Val Pro Tyr Thr Ser Trp Arg Glu Met Phe Leu Glu Arg Pro Arg Val
 130 135 140
 Arg Phe Asp Gly Val Tyr Ile Ser Lys Thr Thr Tyr Ile Arg Gln Gly
 145 150 155 160
 Glu Gln Ser Leu Asp Gly Phe Tyr Arg Ala Trp His Gln Val Glu Tyr
 165 170 175
 Tyr Arg Tyr Ile Arg Phe Phe Pro Asp Gly His Val Met Met Leu Thr
 180 185 190
 Thr Pro Glu Glu Pro Gln Ser Ile Val Pro Arg Leu Arg Thr Arg Asn
 195 200 205

49/57

Thr Arg Thr Asp Ala Ile Leu Leu Gly His Tyr Arg Leu Ser Gln Asp
 210 215 220
 Thr Asp Asn Gln Thr Lys Val Phe Ala Val Ile Thr Lys Lys Lys Glu
 225 230 235 240
 Glu Lys Pro Leu Asp Tyr Lys Tyr Arg Tyr Phe Arg Arg Val Pro Val
 245 250 255
 Gln Glu Ala Asp Gln Ser Phe His Val Gly Leu Gln Leu Cys Ser Ser
 260 265 270
 Gly His Gln Arg Phe Asn Lys Leu Ile Trp Ile His His Ser Cys His
 275 280 285
 Ile Thr Tyr Lys Ser Thr Gly Glu Thr Ala Val Ser Ala Phe Glu Ile
 290 295 300
 Asp Lys Met Tyr Thr Pro Leu Phe Phe Ala Arg Val Arg Ser Tyr Thr
 305 310 315 320
 Ala Phe Ser Glu Arg Pro Leu
 325

<210> 59
 <211> 765
 <212> DNA
 <213> Homo sapiens

<220>
 <221> modified_base
 <222> all n positions
 <223> n=a, c, g or t

<400> 59
 gcagccctgg atcttgactt agagaatgat gatttctttg tcagaaagac tggggctttc 60
 catgcaaatc catatgttct ccgagctttt gaagacttta gaaagttctc tgagcaagat 120
 gattctgtag agcgagatat aattttacag tgtagagaag gtgaaactgt acttccggat 180
 ttggaaaaag atgatatgat tgttcgccga atcccagcac agaagaaaqa agtgccgctg 240
 totggggccc cagatagata ccacccagtc ccttttcccg aaccctggac tcttctccca 300
 gaaattcaag caaaatttct ctgtgtactt gaaaggacat gcccatccaa agaaaaaagt 360
 aatagctgta gaattattgt tcttccatat cggcagaaga aagatgacat gctgacacgt 420
 aagattcagt cctggaaact gggaactacc gtgcctccca tcagtttcac ncttggcccc 480
 tgcagtggag ctgacttgaa gagatgggag gccatccggg aggccagcag actcaggcac 540
 aagaaaaaggc tggatggtga gagactcttt caaaaagatt atggtgagaa tgggagtaag 600
 tccatgagtg atgtcagcgc agaagatgtt caaaaacttgc gtcagctgag ttacgaggag 660
 atgcagaaaa taaaatcaca attaaaagaa caagatcaga aatggcagga tgaccttgca 720
 aaatggaag atcgtcgaaa aagttacact tcagatctgc agaag 765

<210> 60
 <211> 255
 <212> PRT
 <213> Homo sapiens

<400> 60
 Ala Ala Leu Asp Pro Asp Leu Glu Asn Asp Asp Phe Phe Val Arg Lys
 1 5 10 15
 Thr Gly Ala Phe His Ala Asn Pro Tyr Val Leu Arg Ala Phe Glu Asp
 20 25 30
 Phe Arg Lys Phe Ser Glu Gln Asp Asp Ser Val Glu Arg Asp Ile Ile

WO 02/055665

PCT/US02/00311

50/57

35 40 45
 Leu Gln Cys Arg Glu Gly Glu Val Leu Pro Asp Leu Glu Lys Asp
 50 55 60
 Asp Met Ile Val Arg Arg Ile Pro Ala Gln Lys Lys Glu Val Pro Leu
 65 70 75 80
 Ser Gly Ala Pro Asp Arg Tyr His Pro Val Pro Phe Pro Glu Pro Trp
 85 90 95
 Thr Leu Pro Pro Glu Ile Gln Ala Lys Phe Leu Cys Val Leu Glu Arg
 100 105 110
 Thr Cys Pro Ser Lys Glu Lys Ser Asn Ser Cys Arg Ile Leu Val Pro
 115 120 125
 Ser Tyr Arg Gln Lys Lys Asp Asp Met Leu Thr Arg Lys Ile Gln Ser
 130 135 140
 Trp Lys Leu Gly Thr Thr Val Pro Pro Ile Ser Phe Thr Pro Gly Pro
 145 150 155 160
 Cys Ser Glu Ala Asp Leu Lys Arg Trp Glu Ala Ile Arg Glu Ala Ser
 165 170 175
 Arg Leu Arg His Lys Lys Arg Leu Met Val Glu Arg Leu Phe Gln Lys
 180 185 190
 Ile Tyr Gly Glu Asn Gly Ser Lys Ser Met Ser Asp Val Ser Ala Glu
 195 200 205
 Asp Val Gln Asn Leu Arg Gln Leu Arg Tyr Glu Glu Met Gln Lys Ile
 210 215 220
 Lys Ser Gln Leu Lys Glu Gln Asp Gln Lys Trp Gln Asp Asp Leu Ala
 225 230 235 240
 Lys Trp Lys Asp Arg Arg Lys Ser Tyr Thr Ser Asp Leu Gln Lys
 245 250 255

 <210> 61
 <211> 36
 <212> PRT
 <213> Homo sapiens

 <400> 61
 Leu Pro Pro Glu Leu Ser Phe Thr Ile Leu Ser Tyr Leu Asn Ala Thr
 1 5 10 15
 Asp Leu Cys Leu Ala Ser Cys Val Trp Gln Asp Leu Ala Asn Asp Glu
 20 25 30
 Leu Leu Trp Gln
 35

<210> 62
 <211> 42
 <212> PRT
 <213> Homo sapiens

<400> 62
Leu Pro Gly Glu Val Leu Glu Tyr Ile Leu Cys Cys Gly Ser Leu Thr
1 5 10 15
Ala Ala Asp Ile Gly Arg Val Ser Ser Thr Cys Arg Arg Leu Arg Glu
20 25 30
Leu Cys Gln Ser Ser Gly Lys Val Trp Lys
35 40

<210> 63
<211> 44
<212> PRT
<213> Homo sapiens

<400> 63
Leu Ala Glu Val Val Glu Arg Val Leu Thr Phe Leu Pro Ala Lys Ala
1 5 10 15
Leu Leu Arg Val Ala Cys Val Cys Arg Leu Trp Arg Glu Cys Val Arg
20 25 30
Arg Val Leu Arg Thr His Arg Ser Val Thr Trp Ile
35 40

<210> 64
<211> 39
<212> PRT
<213> Homo sapiens

<400> 64
Leu Pro Asp Glu Val Val Leu Lys Ile Phe Ser Tyr Leu Leu Glu Gln
1 5 10 15
Asp Leu Cys Arg Ala Ala Cys Val Cys Lys Arg Phe Ser Glu Leu Ala
20 25 30
Asn Asp Pro Asn Leu Trp Lys
35

<210> 65
<211> 41
<212> PRT
<213> Homo sapiens

<400> 65
Leu Pro Leu Glu Leu Trp Arg Met Ile Leu Ala Tyr Leu His Leu Pro
1 5 10 15
Asp Leu Gly Arg Cys Ser Leu Val Cys Arg Ala Trp Tyr Glu Leu Ile
20 25 30
Leu Ser Leu Asp Ser Thr Arg Trp Arg
35 40

<210> 66
<211> 39
<212> PRT
<213> Homo sapiens

<400> 66

Leu Pro Thr Asp Pro Leu Leu Leu Ile Leu Ser Phe Leu Asp Tyr Arg
1 5 10 15

Asp Leu Ile Asn Cys Cys Tyr Val Ser Arg Arg Leu Ser Gln Leu Ser
20 25 30

Ser His Asp Pro Leu Trp Arg
35

<210> 67

<211> 40

<212> PRT

<213> Homo sapiens

<400> 67

Leu Pro Glu Pro Leu Leu Arg Val Leu Ala Ala Leu Pro Ala Ala
1 5 10 15

Glu Leu Val Gln Ala Cys Arg Leu Val Cys Leu Arg Trp Lys Glu Leu
20 25 30

Val Asp Gly Ala Pro Leu Trp Leu
35 40

<210> 68

<211> 40

<212> PRT

<213> Homo sapiens

<400> 68

Leu Phe Pro Pro Glu Leu Val Glu His Ile Ile Ser Phe Leu Pro Val
1 5 10 15

Arg Asp Leu Val Ala Leu Gly Gln Thr Cys Arg Tyr Phe His Glu Val
20 25 30

Cys Asp Gly Glu Gly Val Trp Arg
35 40

<210> 69

<211> 44

<212> PRT

<213> Homo sapiens

<400> 69

Leu Pro Glu Val Leu Leu Leu His Met Cys Ser Tyr Leu Asp Met Arg
1 5 10 15

Ala Leu Gly Arg Leu Ala Gln Val Tyr Arg Trp Leu Trp His Phe Thr
20 25 30

Asn Cys Asp Leu Leu Arg Arg Gln Ile Ala Trp Ala
35 40

<210> 70

<211> 40

<212> PRT

<213> Homo sapiens

WO 02/055665

PCT/US02/00311

53/57

<400> 70

Leu Pro Leu His Met Leu Asn Asn Ile Leu Tyr Arg Phe Ser Asp Gly
1 5 10 15

Trp Asp Ile Ile Thr Leu Gly Gln Val Thr Pro Thr Leu Tyr Met Leu
20 25 30

Ser Glu Asp Arg Gln Leu Trp Lys
35 40

<210> 71

<211> 39

<212> PRT

<213> Homo sapiens

<400> 71

Leu Pro Asp His Ser Met Val Gln Ile Phe Ser Phe Leu Pro Thr Asn
1 5 10 15

Gln Leu Cys Arg Cys Ala Arg Val Cys Arg Arg Trp Tyr Asn Leu Ala
20 25 30

Trp Asp Pro Arg Leu Trp Arg
35

<210> 72

<211> 44

<212> PRT

<213> Homo sapiens

<400> 72

Ile Pro Leu Glu Ile Leu Val Gln Ile Phe Gly Leu Leu Val Ala Ala
1 5 10 15

Asp Gly Pro Met Pro Phe Leu Gly Arg Ala Ala Arg Val Cys Arg Arg
20 25 30

Trp Gln Glu Ala Ala Ser Gln Pro Ala Leu Trp His
35 40

<210> 73

<211> 39

<212> PRT

<213> Homo sapiens

<400> 73

Leu Pro Pro Glu Val Met Leu Ser Ile Phe Ser Tyr Leu Asn Pro Gln
1 5 10 15

Glu Leu Cys Arg Cys Ser Gln Val Ser Met Lys Trp Ser Gln Leu Thr
20 25 30

Lys Thr Gly Ser Leu Trp Lys
35

<210> 74

<211> 39

<212> PRT

<213> Homo sapiens

WO 02/055665

PCT/US02/00311

54/57

<400> 74

Leu Pro Lys Glu Leu Leu Leu Arg Ile Phe Ser Phe Leu Asp Ile Val
1 5 10 15

Thr Leu Cys Arg Cys Ala Gln Ile Ser Lys Ala Trp Asn Ile Leu Ala
20 25 30

Leu Asp Gly Ser Asn Trp Gln
35

<210> 75

<211> 48

<212> PRT

<213> Homo sapiens

<400> 75

Leu Pro Tyr Glu Leu Ile Gln Leu Ile Leu Asn His Leu Thr Leu Pro
1 5 10 15

Asp Leu Cys Arg Leu Ala Gln Thr Cys Lys Leu Leu Ser Gln His Cys
20 25 30

Cys Asp Pro Leu Gln Tyr Ile His Leu Asn Leu Gln Pro Tyr Trp Ala
35 40 45

<210> 76

<211> 44

<212> PRT

<213> Homo sapiens

<400> 76

Leu Pro Met Glu Val Leu Met Tyr Ile Phe Arg Trp Val Val Ser Ser
1 5 10 15

Asp Leu Asp Leu Arg Ser Leu Glu Gln Leu Ser Leu Val Cys Arg Gly
20 25 30

Phe Tyr Ile Cys Ala Arg Asp Pro Glu Ile Trp Arg
35 40

<210> 77

<211> 49

<212> PRT

<213> Homo sapiens

<400> 77

Leu Pro Pro Glu Ile Gln Ala Lys Phe Leu Cys Val Leu Glu Arg Thr
1 5 10 15

Cys Pro Ser Lys Glu Lys Ser Asn Ser Cys Arg Ile Leu Val Pro Ser
20 25 30

Tyr Arg Gln Lys Lys Asp Asp Met Leu Thr Arg Lys Ile Gln Ser Trp
35 40 45

Lys

<210> 78
 <211> 39
 <212> PRT
 <213> Homo sapiens

<400> 78
 Leu Pro His His Val Val Leu Gln Ile Phe Gln Tyr Leu Pro Leu Leu
 1 5 10 15

Asp Arg Ala Cys Ala Ser Ser Val Cys Arg Arg Trp Asn Glu Val Phe
 20 25 30

His Ile Ser Asp Leu Trp Arg
 35

<210> 79
 <211> 43
 <212> PRT
 <213> Homo sapiens

<400> 79
 Leu Trp Ala Trp Gly Glu Lys Gly Val Leu Ser Asn Ile Ser Ala Leu
 1 5 10 15

Thr Asp Leu Gly Gly Leu Asp Pro Val Trp Leu Val Cys Gly Ser Trp
 20 25 30

Arg Arg His Val Gly Ala Gly Leu Cys Trp Ala
 35 40

<210> 80
 <211> 59
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:
 Oligonucleotide

<400> 80
 agtagtaaca aaggtcaaag acagttgact gtatcgtaga ggaagccttc aattaagtt 59

<210> 81
 <211> 58
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:
 Oligonucleotide

<400> 81
 gagggttactt acttagagct cgaagtcctta cttacttagc tcaattctct tcacacca 58

<210> 82
 <211> 12
 <212> PRT
 <213> Homo sapiens

<400> 82

56/57

Cys Asp Gly Glu Lys Asp Thr Tyr Ser Tyr Leu Ala
1 5 10

<210> 83
<211> 25
<212> PRT
<213> Homo sapiens

<400> 83
Cys Glu Ser Ser Phe Ser Leu Asn Met Asn Phe Ser Ser Lys Arg Thr
1 5 10 15

Lys Phe Lys Ile Thr Thr Ser Met Gln
20 25

<210> 84
<211> 12
<212> PRT
<213> Homo sapiens

<400> 84
Cys Glu Glu Ala Gln Val Arg Lys Glu Asn Gln Trp
1 5 10

<210> 85
<211> 19
<212> PRT
<213> Homo sapiens

<400> 85
Asn Ala Gly Ser Val Glu Gln Thr Pro Lys Lys Pro Gly Leu Arg Arg
1 5 10 15

Arg Gln Thr

<210> 86
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 86
cctgggggat gttctca 17

<210> 87
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 87
ggcttcggg catttag 17

<210> 88
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 88
catctggcac gattcca

17

<210> 89
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 89
ccgctcatcg tatgaca

17

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
18 July 2002 (18.07.2002)

PCT

(10) International Publication Number
WO 02/055665 A3

(51) International Patent Classification⁷: **C12Q 1/00**,
G01N 33/53

(21) International Application Number: PCT/US02/00311

(22) International Filing Date: 7 January 2002 (07.01.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/260,179 5 January 2001 (05.01.2001) US

(71) Applicant: **NEW YORK UNIVERSITY** [US/US]; 70
Washington Square South, New York, NY 10012 (US).

(72) Inventor: **PAGANO, Michele**; One Washington Square
Village, New York, NY 10012 (US).

(74) Agents: **REIN, Barry, D.** et al.; Pennie & Edmonds LLP,
1155 Avenue of the Americas, New York, NY 10036 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,

CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN,
YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:
26 September 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 02/055665 A3

(54) Title: METHODS TO IDENTIFY COMPOUNDS USEFUL FOR THE TREATMENT OF PROLIFERATIVE AND DIFFERENTIATIVE DISORDERS

(57) Abstract: The present invention relates to the discovery, identification and characterization of nucleotides that encode novel substrate-targeting subunits of ubiquitin ligases. The invention encompasses nucleotides encoding novel substrate-targeting subunits of ubiquitin ligases: FBP1, FBP2, FBP3, FBP4, FBP5, FBP6, FBP7, FBP8, FBP9, FBP10, FBP11, FBP12, FBP13, FBP14, FBP15, FBP16, FBP17, FBP18, FBP19, FBP20, FBP21, FBP22, FBP23, FBP24, and FBP25, transgenic mice, knock-out mice, host cell expression systems and proteins encoded by the nucleotides of the present invention. The present invention relates to screening assays that use the novel substrate-targeting subunits to identify potential therapeutic agents such as small molecules, compounds or derivatives and analogues of the novel ubiquitin ligases which modulates activity of the novel ubiquitin ligases for the treatment of proliferative and differentiative disorders, such as cancer, major opportunistic infections, immune disorders, certain cardiovascular diseases, and inflammatory disorders. The invention further encompasses therapeutic protocols and pharmaceutical compositions designed to target ubiquitin ligases and their substrates for the treatment of proliferative disorders.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/00311

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C12Q 1/00; G01N 33/53
US CL : 435/4, 7.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
U.S. : 435/4, 7.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
WEST, MEDLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 00/12679 A1 (CHIAUR et al.) 09 March 2000 (09.03.00), entire article, especially page 40, lines 24-35, pages 46-47, page 95, and page 13, lines 16-35.	1-9
Y	WO 00/75184 A1 (ZHANG et al) 14 December 2000 (14.12.2000), entire article, especially pages 23, 28, 31, and 48-49.	1-9

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T"

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X"

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y"

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"Z"

document member of the same patent family

Date of the actual completion of the international search

21 May 2002 (21.05.2002)

Date of mailing of the international search report

02 JUL 2002

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

Telicia D. Roberts for
Gary B. Nickol Ph.D.

Telephone No. 703-308-0196